



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 04:36 AM UTC

PDB ID : 3SYP / pdb_00003syp
Title : Crystal structure of the G protein-gated inward rectifier K⁺ channel GIRK2 (Kir3.2) R201A mutant
Authors : Whorton, M.R.; MacKinnon, R.
Deposited on : 2011-07-18
Resolution : 3.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1816 (3.14-3.10)
Clashscore	190562	1906 (3.14-3.10)
Ramachandran outliers	187476	1802 (3.14-3.10)
Sidechain outliers	187428	1802 (3.14-3.10)
RSRZ outliers	180081	1816 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	17% 68% 20% 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G protein-activated inward rectifier potassium channel 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	305	Total	C	N	O	S	0	0	0
			2311	1497	376	426	12			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	MET	-	expression tag	UNP P48542
A	51	ALA	-	expression tag	UNP P48542
A	201	ALA	ARG	engineered mutation	UNP P48542
A	260	THR	SER	SEE REMARK 999	UNP P48542
A	313	MET	ILE	SEE REMARK 999	UNP P48542
A	344	LEU	MET	SEE REMARK 999	UNP P48542
A	381	SER	-	expression tag	UNP P48542
A	382	ASN	-	expression tag	UNP P48542
A	383	SER	-	expression tag	UNP P48542
A	384	LEU	-	expression tag	UNP P48542
A	385	GLU	-	expression tag	UNP P48542
A	386	VAL	-	expression tag	UNP P48542
A	387	LEU	-	expression tag	UNP P48542
A	388	PHE	-	expression tag	UNP P48542
A	389	GLN	-	expression tag	UNP P48542


- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	K	0	0
			5	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: G protein-activated inward rectifier potassium channel 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.43Å 85.43Å 178.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.36 – 3.12 42.36 – 3.12	Depositor EDS
% Data completeness (in resolution range)	96.7 (42.36-3.12) 96.9 (42.36-3.12)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.262 , 0.296 0.257 , 0.286	Depositor DCC
R_{free} test set	838 reflections (6.95%)	wwPDB-VP
Wilson B-factor (Å ²)	91.2	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2316	wwPDB-VP
Average B, all atoms (Å ²)	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2361	0.93	3/3217 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	255	ILE	CA-C-N	6.16	127.55	119.84
1	A	255	ILE	C-N-CA	6.16	127.55	119.84
1	A	233	HIS	N-CA-C	5.75	117.71	109.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2311	0	2195	38	1
2	A	5	0	0	0	0
All	All	2316	0	2195	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:PHE:O	1:A:190:CYS:HB2	1.77	0.85
1:A:243:LEU:HB2	1:A:262:ILE:HD11	1.70	0.72
1:A:293:GLU:HG2	1:A:370:LYS:HD3	1.76	0.67
1:A:276:VAL:HG13	1:A:277:SER:H	1.63	0.63
1:A:208:SER:HB3	1:A:225:ARG:HB3	1.79	0.62
1:A:233:HIS:ND1	1:A:234:ILE:HG22	2.15	0.62
1:A:93:PHE:O	1:A:97:ILE:HG12	2.03	0.58
1:A:329:THR:HA	1:A:332:ILE:HD12	1.89	0.54
1:A:234:ILE:HD11	1:A:312:GLY:HA3	1.90	0.53
1:A:225:ARG:HG2	1:A:356:PHE:CE1	2.44	0.52
1:A:295:SER:H	1:A:298:GLN:NE2	2.07	0.52
1:A:58:TYR:CE1	1:A:257:LEU:HD21	2.45	0.52
1:A:99:VAL:O	1:A:103:THR:HG23	2.11	0.51
1:A:58:TYR:CZ	1:A:257:LEU:CD2	2.96	0.49
1:A:108:PHE:O	1:A:112:ILE:HG12	2.14	0.47
1:A:304:LEU:C	1:A:304:LEU:HD12	2.39	0.47
1:A:207:PHE:CE2	1:A:308:VAL:HG12	2.49	0.47
1:A:336:TYR:HA	1:A:361:GLU:HA	1.98	0.46
1:A:209:THR:HB	1:A:357:HIS:ND1	2.30	0.46
1:A:202:ALA:HB2	1:A:321:CYS:HB2	1.98	0.46
1:A:313:MET:HG3	1:A:318:GLY:O	2.17	0.45
1:A:205:LEU:HD23	1:A:228:ASP:HB2	1.98	0.45
1:A:296:LYS:HG3	1:A:334:TRP:CG	2.52	0.45
1:A:322:GLN:OE1	1:A:324:ARG:NH1	2.38	0.45
1:A:150:GLU:O	1:A:155:ILE:N	2.50	0.44
1:A:104:VAL:HA	1:A:107:LEU:HD12	2.00	0.44
1:A:337:ARG:HD2	1:A:360:TYR:OH	2.18	0.44
1:A:83:PHE:HZ	1:A:193:VAL:HG21	1.84	0.43
1:A:106:TRP:HE1	1:A:152:GLU:CD	2.26	0.43
1:A:136:THR:HB	1:A:161:VAL:H	1.83	0.43
1:A:298:GLN:O	1:A:299:LEU:C	2.63	0.41
1:A:60:ARG:NH1	1:A:64:LYS:HD3	2.36	0.41
1:A:97:ILE:O	1:A:101:VAL:HG23	2.21	0.41
1:A:328:ILE:O	1:A:329:THR:C	2.64	0.40
1:A:243:LEU:O	1:A:259:GLN:HA	2.21	0.40
1:A:235:VAL:HG22	1:A:315:GLU:OE1	2.21	0.40
1:A:239:ILE:HG13	1:A:240:ARG:N	2.36	0.40
1:A:299:LEU:HB3	1:A:300:PRO:HD3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:GLN:O	1:A:349:TYR:OH[4_545]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	299/340 (88%)	279 (93%)	19 (6%)	1 (0%)	36 65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	234	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	236/305 (77%)	224 (95%)	12 (5%)	21 50

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ILE
1	A	145	PHE
1	A	182	ILE
1	A	209	THR
1	A	226	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	234	ILE
1	A	239	ILE
1	A	243	LEU
1	A	276	VAL
1	A	294	ILE
1	A	358	GLU
1	A	380	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	GLN
1	A	298	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/340 (89%)	0.95	59 (19%) 3 2	59, 99, 234, 298	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	LEU	5.5
1	A	181	SER	5.2
1	A	170	ILE	5.0
1	A	164	ASP	4.7
1	A	78	TYR	4.5
1	A	162	ILE	4.5
1	A	154	THR	4.3
1	A	166	CYS	4.3
1	A	79	LEU	4.1
1	A	83	PHE	4.1
1	A	163	THR	4.0
1	A	184	ASN	4.0
1	A	135	VAL	3.9
1	A	134	CYS	3.9
1	A	232	SER	3.8
1	A	153	THR	3.8
1	A	137	ASN	3.7
1	A	174	LEU	3.7
1	A	157	TYR	3.6
1	A	160	ARG	3.6
1	A	175	ILE	3.5
1	A	172	LEU	3.5
1	A	161	VAL	3.4
1	A	136	THR	3.3
1	A	363	SER	3.2
1	A	116	ILE	3.2
1	A	167	PRO	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	169	GLY	3.1
1	A	107	LEU	3.0
1	A	139	ASN	3.0
1	A	155	ILE	3.0
1	A	117	ALA	2.9
1	A	236	GLU	2.8
1	A	276	VAL	2.8
1	A	141	PHE	2.8
1	A	186	PHE	2.8
1	A	76	TYR	2.8
1	A	346	ASP	2.7
1	A	177	SER	2.6
1	A	82	ILE	2.6
1	A	111	MET	2.6
1	A	115	LEU	2.6
1	A	171	ILE	2.5
1	A	96	LEU	2.5
1	A	188	VAL	2.5
1	A	158	GLY	2.4
1	A	67	VAL	2.4
1	A	266	TYR	2.4
1	A	183	VAL	2.4
1	A	185	ALA	2.3
1	A	77	ARG	2.2
1	A	113	TRP	2.2
1	A	69	HIS	2.2
1	A	114	TRP	2.1
1	A	235	VAL	2.1
1	A	178	VAL	2.1
1	A	349	TYR	2.1
1	A	142	VAL	2.1
1	A	168	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	K	A	501	1/1	0.80	0.20	165,165,165,165	1
2	K	A	502	1/1	0.85	0.20	350,350,350,350	1
2	K	A	504	1/1	0.91	0.18	171,171,171,171	1
2	K	A	505	1/1	0.92	0.18	123,123,123,123	0
2	K	A	503	1/1	0.97	0.25	107,107,107,107	1

6.5 Other polymers [i](#)

There are no such residues in this entry.