



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 05:32 PM UTC

PDB ID : 6SY2 / pdb_00006sy2
BMRB ID : 34437
Title : Structure of the BRK domain of the SWI/SNF chromatin remodelling complex subunit BRG1 reveals a potential role in protein-protein interactions
Authors : Allen, M.D.; Bycoft, M.; Zinzalla, G.
Deposited on : 2019-09-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	49	57% 16% . 24%

2 Ensemble composition and analysis

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *none*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:615-A:651 (37)	0.27	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 4, 6, 8, 9, 10, 18, 20
2	3, 5, 7, 11, 13, 16
3	2, 14, 15, 17, 19
Single-model clusters	12

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 734 atoms, of which 364 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transcription activator BRG1.


Mol	Chain	Residues	Atoms						Trace
1	A	49	Total	C	H	N	O	S	0
			734	230	364	61	77	2	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transcription activator BRG1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Transcription activator BRG1




4.2.2 Score per residue for model 2

- Molecule 1: Transcription activator BRG1




4.2.3 Score per residue for model 3

- Molecule 1: Transcription activator BRG1




4.2.4 Score per residue for model 4

- Molecule 1: Transcription activator BRG1




4.2.5 Score per residue for model 5

- Molecule 1: Transcription activator BRG1




4.2.6 Score per residue for model 6

- Molecule 1: Transcription activator BRG1




4.2.7 Score per residue for model 7

- Molecule 1: Transcription activator BRG1




4.2.8 Score per residue for model 8

- Molecule 1: Transcription activator BRG1



4.2.9 Score per residue for model 9

- Molecule 1: Transcription activator BRG1




4.2.10 Score per residue for model 10 (medoid)

- Molecule 1: Transcription activator BRG1



4.2.11 Score per residue for model 11

- Molecule 1: Transcription activator BRG1




4.2.12 Score per residue for model 12

- Molecule 1: Transcription activator BRG1




4.2.13 Score per residue for model 13

- Molecule 1: Transcription activator BRG1




4.2.14 Score per residue for model 14

- Molecule 1: Transcription activator BRG1




4.2.15 Score per residue for model 15

- Molecule 1: Transcription activator BRG1




4.2.16 Score per residue for model 16

- Molecule 1: Transcription activator BRG1




4.2.17 Score per residue for model 17

- Molecule 1: Transcription activator BRG1




4.2.18 Score per residue for model 18

- Molecule 1: Transcription activator BRG1




4.2.19 Score per residue for model 19

- Molecule 1: Transcription activator BRG1



4.2.20 Score per residue for model 20

- Molecule 1: Transcription activator BRG1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	558
Number of shifts mapped to atoms	558
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	280	286	285	7±3
All	All	5600	5720	5700	137

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:620:ILE:HD13	1:A:627:ILE:CG1	0.81	2.06	18	9
1:A:620:ILE:HD13	1:A:627:ILE:HG12	0.80	1.54	4	10
1:A:620:ILE:HD12	1:A:626:LYS:O	0.72	1.85	4	7
1:A:617:VAL:CG1	1:A:651:VAL:HG13	0.64	2.23	18	3
1:A:620:ILE:HD13	1:A:627:ILE:HG13	0.56	1.77	18	4
1:A:622:VAL:HG23	1:A:648:GLY:O	0.55	2.01	12	4
1:A:619:VAL:HG23	1:A:649:TYR:CD1	0.54	2.37	19	3
1:A:620:ILE:HD13	1:A:620:ILE:C	0.53	2.28	6	3
1:A:619:VAL:O	1:A:627:ILE:HG23	0.53	2.03	16	1
1:A:621:HIS:CD2	1:A:649:TYR:CD1	0.52	2.98	18	5
1:A:621:HIS:CG	1:A:649:TYR:CE1	0.51	2.99	4	10
1:A:634:PRO:HG2	1:A:639:LEU:HD12	0.51	1.83	6	7
1:A:621:HIS:CD2	1:A:649:TYR:CD2	0.50	3.00	7	1
1:A:629:THR:OG1	1:A:631:THR:HG22	0.50	2.05	16	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:618:LYS:O	1:A:619:VAL:HG13	0.48	2.09	5	11
1:A:621:HIS:CG	1:A:649:TYR:CE2	0.47	3.03	20	3
1:A:628:LEU:HD12	1:A:649:TYR:CE1	0.47	2.45	7	2
1:A:628:LEU:CD1	1:A:649:TYR:CE1	0.47	2.98	3	5
1:A:628:LEU:CD1	1:A:649:TYR:CE2	0.46	2.98	16	10
1:A:621:HIS:ND1	1:A:649:TYR:CE2	0.46	2.83	6	5
1:A:628:LEU:HD12	1:A:649:TYR:CE2	0.46	2.46	9	5
1:A:621:HIS:ND1	1:A:649:TYR:CE1	0.45	2.84	10	3
1:A:619:VAL:HG12	1:A:651:VAL:HA	0.45	1.89	15	1
1:A:619:VAL:CG2	1:A:649:TYR:CD1	0.45	3.00	19	1
1:A:621:HIS:CD2	1:A:649:TYR:CE2	0.44	3.05	7	1
1:A:619:VAL:HA	1:A:650:GLU:O	0.44	2.12	17	4
1:A:642:TRP:CD2	1:A:646:ASN:ND2	0.44	2.86	19	1
1:A:617:VAL:O	1:A:633:ALA:HB1	0.43	2.13	2	3
1:A:621:HIS:HB2	1:A:649:TYR:CZ	0.43	2.48	18	9
1:A:620:ILE:CD1	1:A:627:ILE:CG1	0.43	2.97	7	2
1:A:619:VAL:HG23	1:A:649:TYR:CE1	0.41	2.50	19	1
1:A:617:VAL:HG12	1:A:651:VAL:HG13	0.41	1.90	18	1
1:A:619:VAL:CG2	1:A:628:LEU:HD12	0.40	2.47	19	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	37/49 (76%)	36±1 (98±2%)	1±1 (2±2%)	0±0 (0±0%)	100	100
All	All	740/980 (76%)	722 (98%)	18 (2%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	30/41 (73%)	24±1 (80±4%)	6±1 (20±4%)	3	33
All	All	600/820 (73%)	483 (80%)	117 (20%)	3	33

All 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	629	THR	20
1	A	618	LYS	19
1	A	644	GLU	10
1	A	626	LYS	10
1	A	645	MET	9
1	A	638	GLN	9
1	A	620	ILE	7
1	A	623	GLU	6
1	A	631	THR	6
1	A	632	ASP	6
1	A	640	GLU	5
1	A	635	LYS	3
1	A	627	ILE	3
1	A	624	SER	2
1	A	643	LEU	2

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 89% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch_output*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	558
Number of shifts mapped to atoms	558
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	48	-0.40 ± 0.28	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	45	-0.56 ± 0.15	Should be applied
$^{13}\text{C}'$	48	-0.18 ± 0.35	None needed (< 0.5 ppm)
^{15}N	44	0.10 ± 0.83	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 437 atoms were assigned a chemical shift out of a possible 491. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	179/183 (98%)	74/75 (99%)	72/74 (97%)	33/34 (97%)
Sidechain	246/279 (88%)	184/184 (100%)	60/90 (67%)	2/5 (40%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	12/29 (41%)	12/14 (86%)	0/12 (0%)	0/3 (0%)
Overall	437/491 (89%)	270/273 (99%)	132/176 (75%)	35/42 (83%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 89%, i.e. 557 atoms were assigned a chemical shift out of a possible 627. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	236/241 (98%)	96/98 (98%)	96/98 (98%)	44/45 (98%)
Sidechain	309/357 (87%)	229/233 (98%)	76/115 (66%)	4/9 (44%)
Aromatic	12/29 (41%)	12/14 (86%)	0/12 (0%)	0/3 (0%)
Overall	557/627 (89%)	337/345 (98%)	172/225 (76%)	48/57 (84%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	646	ASN	HB2	1.20	1.27 – 4.34	-5.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

