



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 12:54 AM UTC

PDB ID : 8SYD / pdb_00008syd
Title : X-ray crystal structure of UDP-2,3-diacetamido-2,3-dideoxy-glucuronic acid-2-epimerase from *Thermus thermophilus* strain HB27, D98N variant in the presence of UDP-2,3-diacetamido-2,3-dideoxy-glucuronic acid and UDP-N-acetylglucosamine at pH 6
Authors : Kroft, C.W.; Thoden, J.B.; Holden, H.M.
Deposited on : 2023-05-25
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	% 82% 10% 6%
1	B	385	% 84% 10% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-2,3-diacetamido-2,3-dideoxy-glucuronic acid-2-epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	0
			2833	1798	506	520	9			
1	B	371	Total	C	N	O	S	0	2	0
			2926	1858	525	533	10			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP Q72KY0
A	-18	GLY	-	expression tag	UNP Q72KY0
A	-17	SER	-	expression tag	UNP Q72KY0
A	-16	SER	-	expression tag	UNP Q72KY0
A	-15	HIS	-	expression tag	UNP Q72KY0
A	-14	HIS	-	expression tag	UNP Q72KY0
A	-13	HIS	-	expression tag	UNP Q72KY0
A	-12	HIS	-	expression tag	UNP Q72KY0
A	-11	HIS	-	expression tag	UNP Q72KY0
A	-10	HIS	-	expression tag	UNP Q72KY0
A	-9	SER	-	expression tag	UNP Q72KY0
A	-8	SER	-	expression tag	UNP Q72KY0
A	-7	GLU	-	expression tag	UNP Q72KY0
A	-6	ASN	-	expression tag	UNP Q72KY0
A	-5	LEU	-	expression tag	UNP Q72KY0
A	-4	TYR	-	expression tag	UNP Q72KY0
A	-3	PHE	-	expression tag	UNP Q72KY0
A	-2	GLN	-	expression tag	UNP Q72KY0
A	-1	GLY	-	expression tag	UNP Q72KY0
A	0	HIS	-	expression tag	UNP Q72KY0
A	98	ASN	ASP	engineered mutation	UNP Q72KY0
B	-19	MET	-	initiating methionine	UNP Q72KY0
B	-18	GLY	-	expression tag	UNP Q72KY0
B	-17	SER	-	expression tag	UNP Q72KY0
B	-16	SER	-	expression tag	UNP Q72KY0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP Q72KY0
B	-14	HIS	-	expression tag	UNP Q72KY0
B	-13	HIS	-	expression tag	UNP Q72KY0
B	-12	HIS	-	expression tag	UNP Q72KY0
B	-11	HIS	-	expression tag	UNP Q72KY0
B	-10	HIS	-	expression tag	UNP Q72KY0
B	-9	SER	-	expression tag	UNP Q72KY0
B	-8	SER	-	expression tag	UNP Q72KY0
B	-7	GLU	-	expression tag	UNP Q72KY0
B	-6	ASN	-	expression tag	UNP Q72KY0
B	-5	LEU	-	expression tag	UNP Q72KY0
B	-4	TYR	-	expression tag	UNP Q72KY0
B	-3	PHE	-	expression tag	UNP Q72KY0
B	-2	GLN	-	expression tag	UNP Q72KY0
B	-1	GLY	-	expression tag	UNP Q72KY0
B	0	HIS	-	expression tag	UNP Q72KY0
B	98	ASN	ASP	engineered mutation	UNP Q72KY0

- Molecule 2 is (2 {S},3 {S},4 {R},5 {R},6 {R})-4,5-diacetamido-6-[[[(2 {R},3 {S},4 {R},5 {R})-5-[2,4-bis(oxidanylidene)pyrimidin-1-yl]-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-oxidanyl-phosphoryl]oxy-3-oxidanyl-oxane-2-carboxylic acid (CCD ID: MJL) (formula: C₁₉H₂₈N₄O₁₈P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			43	19	4	18	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			43	19	4	18	2		

- Molecule 3 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (CCD ID: UD1) (formula: C₁₇H₂₇N₃O₁₇P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
3	B	1	Total	C	N	O	P	0	0
			39	17	3	17	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total	O	0	0
			137	137		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	137	Total 137	O 137	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-2,3-diacetamido-2,3-dideoxy-glucuronic acid-2-epimerase



- Molecule 1: UDP-2,3-diacetamido-2,3-dideoxy-glucuronic acid-2-epimerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.27Å 128.44Å 58.13Å 90.00° 114.06° 90.00°	Depositor
Resolution (Å)	40.12 – 2.20 40.12 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.12-2.20) 99.5 (40.12-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0403	Depositor
R, R_{free}	0.172 , 0.229 0.181 , 0.233	Depositor DCC
R_{free} test set	1822 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.047 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6199	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UD1, CL, MJL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2889	1.10	3/3918 (0.1%)
1	B	0.64	0/2990	1.11	6/4051 (0.1%)
All	All	0.64	0/5879	1.10	9/7969 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
All	All	0	4

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	89	LYS	CB-CA-C	7.05	117.14	111.00
1	A	52	GLU	CB-CA-C	6.34	121.32	110.86
1	B	322	GLN	CB-CA-C	6.14	119.68	109.56
1	A	243	THR	CA-CB-OG1	-6.08	100.48	109.60
1	B	151	GLU	CB-CG-CD	5.87	122.58	112.60
1	B	176	ASP	CA-CB-CG	5.80	118.40	112.60
1	B	361	VAL	N-CA-CB	5.17	116.25	110.51
1	A	144	ASP	CA-CB-CG	-5.08	107.52	112.60
1	B	132	GLU	CB-CG-CD	5.01	121.11	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ARG	Sidechain
1	B	128	ARG	Sidechain
1	B	187	ARG	Sidechain
1	B	336	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2833	0	2853	22	0
1	B	2926	0	2948	29	0
2	A	43	0	0	1	0
2	B	43	0	0	0	0
3	A	39	0	25	0	0
3	B	39	0	25	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	137	0	0	2	0
5	B	137	0	0	2	0
All	All	6199	0	5851	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:323:ASN:HD21	1:B:326:ASP:CG	1.80	0.90
1:A:272:MET:HA	1:A:272:MET:HE2	1.68	0.75
1:A:33:GLU:O	5:A:501:HOH:O	2.03	0.74
1:B:244:ARG:HG3	1:B:245:LYS:N	2.04	0.70
1:A:216:GLN:OE1	5:A:502:HOH:O	2.12	0.68
1:B:352:GLN:HE21	1:B:352:GLN:H	1.41	0.67
1:A:132:GLU:O	1:A:136:ARG:HG2	1.98	0.63
1:B:205:ALA:HA	1:B:281:VAL:O	2.01	0.60
1:B:323:ASN:ND2	1:B:326:ASP:CG	2.57	0.59
1:A:205:ALA:HA	1:A:281:VAL:O	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:332:LEU:HD23	1:B:335:MET:HE1	1.86	0.58
1:B:352:GLN:HE21	1:B:352:GLN:N	2.00	0.58
1:A:146:LEU:CD1	1:A:162:ILE:HG21	2.34	0.58
1:B:332:LEU:HA	1:B:335:MET:HE3	1.85	0.57
1:B:200:LYS:CE	5:B:579:HOH:O	2.53	0.57
1:B:300:THR:HB	1:B:318:LEU:HD23	1.86	0.57
1:B:98:ASN:ND2	1:B:132:GLU:HB3	2.20	0.56
1:B:200:LYS:HE2	5:B:579:HOH:O	2.05	0.56
1:B:146:LEU:HD22	1:B:157:LEU:HD22	1.89	0.54
1:B:225:ALA:O	1:B:229:VAL:HG23	2.08	0.54
1:B:130:MET:HE2	1:B:132:GLU:HG2	1.89	0.53
1:B:98:ASN:HD22	1:B:132:GLU:HB3	1.74	0.53
1:B:94:LEU:HD23	1:B:117:ALA:HB3	1.90	0.53
1:B:271:VAL:O	1:B:275:LYS:HB2	2.09	0.53
1:A:94:LEU:HD12	1:A:117:ALA:HB3	1.92	0.52
1:A:11:ALA:HB1	1:A:41:HIS:CE1	2.45	0.51
1:A:216:GLN:HG3	1:A:250:PHE:CG	2.46	0.51
1:A:272:MET:HE2	1:A:272:MET:CA	2.40	0.50
1:B:332:LEU:HD23	1:B:335:MET:CE	2.41	0.50
1:A:97:GLY:HA3	2:A:401:MJL:C6'	2.42	0.49
1:B:282:THR:O	1:B:300:THR:HA	2.13	0.48
1:A:206:THR:O	1:A:282:THR:HA	2.15	0.47
1:B:5:ILE:HD12	1:B:31:VAL:HG13	1.96	0.47
1:A:163:PRO:HG2	1:A:166:ARG:HD3	1.97	0.46
1:B:41:HIS:CD2	3:B:402:UD1:H5	2.50	0.46
1:A:270:MET:O	1:A:274:GLU:HG3	2.15	0.46
1:A:272:MET:HA	1:A:272:MET:CE	2.43	0.45
1:B:189:SER:HB3	1:B:272[B]:MET:HG3	1.99	0.45
1:B:189:SER:HB3	1:B:272[A]:MET:HG3	1.98	0.45
1:A:94:LEU:CD1	1:A:117:ALA:HB3	2.46	0.44
1:B:271:VAL:HG12	1:B:272[B]:MET:HE2	2.00	0.44
1:A:165:ASN:HD22	1:A:165:ASN:C	2.26	0.43
1:A:30:GLY:O	1:A:31:VAL:HG23	2.18	0.43
1:B:206:THR:O	1:B:282:THR:HA	2.19	0.43
1:B:98:ASN:ND2	1:B:132:GLU:CB	2.81	0.43
1:A:282:THR:O	1:A:300:THR:HA	2.19	0.43
1:A:95:VAL:HB	1:A:102:THR:HG23	2.00	0.42
1:A:88:GLU:O	1:A:89:LYS:C	2.62	0.42
1:A:349:ARG:HB3	1:A:352:GLN:HG2	2.02	0.42
1:B:12:ARG:HB3	1:B:13:PRO:CD	2.50	0.41
1:B:363:ILE:O	1:B:364:ARG:C	2.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/385 (93%)	344 (96%)	13 (4%)	1 (0%)	36	42
1	B	371/385 (96%)	354 (95%)	17 (5%)	0	100	100
All	All	729/770 (95%)	698 (96%)	30 (4%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/321 (92%)	282 (95%)	15 (5%)	21	27
1	B	306/321 (95%)	290 (95%)	16 (5%)	21	26
All	All	603/642 (94%)	572 (95%)	31 (5%)	22	27

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ARG
1	A	58	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	LEU
1	A	98	ASN
1	A	128	ARG
1	A	132	GLU
1	A	165	ASN
1	A	198	GLN
1	A	233	VAL
1	A	272	MET
1	A	303	GLU
1	A	312	LYS
1	A	335	MET
1	A	338	LYS
1	A	352	GLN
1	B	0	HIS
1	B	26	ARG
1	B	32	ARG
1	B	87	LYS
1	B	89	LYS
1	B	132	GLU
1	B	151	GLU
1	B	179	LEU
1	B	244	ARG
1	B	272[A]	MET
1	B	272[B]	MET
1	B	281	VAL
1	B	323	ASN
1	B	341	GLU
1	B	352	GLN
1	B	361	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN
1	A	168	HIS
1	A	180	HIS
1	A	198	GLN
1	B	198	GLN
1	B	216	GLN
1	B	323	ASN
1	B	333	HIS
1	B	352	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MJL	A	401	-	43,45,45	1.55	4 (9%)	60,68,68	1.66	11 (18%)
3	UD1	A	402	-	40,41,41	0.54	0	59,62,62	0.88	2 (3%)
3	UD1	B	402	-	40,41,41	0.49	0	59,62,62	0.89	2 (3%)
2	MJL	B	401	-	43,45,45	1.38	7 (16%)	60,68,68	1.38	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MJL	A	401	-	-	4/32/69/69	0/3/3/3
3	UD1	A	402	-	-	4/26/63/63	0/3/3/3
3	UD1	B	402	-	-	4/26/63/63	0/3/3/3
2	MJL	B	401	-	-	5/32/69/69	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	MJL	C5-C4	-6.65	1.29	1.43
2	B	401	MJL	C5-C4	-4.50	1.33	1.43
2	B	401	MJL	PA-O3A	3.67	1.63	1.59
2	A	401	MJL	C2-N3	-3.48	1.31	1.38
2	B	401	MJL	O3C-C3C	2.53	1.49	1.43
2	B	401	MJL	C3'-N3'	2.41	1.49	1.45
2	A	401	MJL	C6-C5	-2.36	1.29	1.35
2	B	401	MJL	C2-N1	2.30	1.42	1.38
2	A	401	MJL	C4-N3	-2.29	1.34	1.38
2	B	401	MJL	O4C-C4C	-2.19	1.40	1.45
2	B	401	MJL	O2-C2	2.01	1.26	1.23

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	MJL	C4-N3-C2	-6.35	118.73	126.61
2	B	401	MJL	C4-N3-C2	-4.85	120.60	126.61
2	A	401	MJL	C5-C4-N3	4.55	121.18	114.80
2	B	401	MJL	C5-C4-N3	3.70	119.98	114.80
3	B	402	UD1	C1'-C2'-N2'	3.67	117.07	110.92
2	A	401	MJL	N3-C2-N1	3.53	119.49	114.89
2	A	401	MJL	O5'-C1'-O3B	-3.21	107.17	111.36
2	B	401	MJL	N3-C2-N1	2.73	118.44	114.89
2	A	401	MJL	O3C-C3C-C4C	-2.53	103.81	111.08
2	A	401	MJL	C6-N1-C2	-2.46	118.00	121.00
2	B	401	MJL	C6-N1-C2	-2.45	118.01	121.00
3	A	402	UD1	C1'-C2'-N2'	-2.43	106.84	110.92
2	A	401	MJL	O2-C2-N3	-2.39	117.08	121.49
2	B	401	MJL	O4C-C4C-C3C	2.31	109.73	105.15
3	B	402	UD1	O3B-C3B-C2B	2.29	119.16	111.82
2	A	401	MJL	C2C-C3C-C4C	2.25	106.96	102.61
3	A	402	UD1	O2A-PA-O1A	2.21	122.74	112.44
2	A	401	MJL	O4-C4-N3	-2.17	116.12	119.27
2	A	401	MJL	PB-O3B-C1'	2.17	130.05	121.21
2	B	401	MJL	C2C-C1C-N1	2.07	119.01	113.25
2	A	401	MJL	C6-C5-C4	2.06	122.17	119.53

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	MJL	C5C-O5C-PA-O1A
2	A	401	MJL	C5C-O5C-PA-O2A
2	A	401	MJL	C5C-O5C-PA-O3A
2	A	401	MJL	O5'-C1'-O3B-PB
2	B	401	MJL	C5C-O5C-PA-O1A
2	B	401	MJL	C5C-O5C-PA-O3A
2	B	401	MJL	O5'-C1'-O3B-PB
3	A	402	UD1	C4'-C5'-C6'-O6'
3	A	402	UD1	O5'-C5'-C6'-O6'
3	B	402	UD1	C4'-C5'-C6'-O6'
3	B	402	UD1	O5'-C5'-C6'-O6'
3	A	402	UD1	PB-O3A-PA-O5B
3	B	402	UD1	PB-O3A-PA-O5B
2	B	401	MJL	C5C-O5C-PA-O2A
3	A	402	UD1	PB-O3A-PA-O1A
3	B	402	UD1	PB-O3A-PA-O1A
2	B	401	MJL	C1'-O3B-PB-O2B


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	MJL	1	0
3	B	402	UD1	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/385 (93%)	-0.43	3 (0%) 82 80	7, 14, 34, 62	0
1	B	371/385 (96%)	-0.42	4 (1%) 78 76	6, 13, 35, 69	2 (0%)
All	All	731/770 (94%)	-0.43	7 (0%) 79 77	6, 13, 35, 69	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	0	HIS	4.3
1	A	2	TRP	2.7
1	A	1	MET	2.6
1	A	29	PRO	2.3
1	B	364	ARG	2.3
1	B	363	ILE	2.1
1	B	2	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MJL	A	401	43/43	0.98	0.04	7,8,9,9	0
2	MJL	B	401	43/43	0.98	0.05	5,6,7,7	0
3	UD1	A	402	39/39	0.98	0.05	7,9,13,13	0
3	UD1	B	402	39/39	0.98	0.04	7,9,10,11	0
4	CL	B	403	1/1	0.98	0.04	11,11,11,11	0
4	CL	A	403	1/1	0.99	0.04	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around MJL B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UD1 A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.