



## wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 06:50 AM UTC

PDB ID : 7T0O / pdb\_00007t0o  
EMDB ID : EMD-25581  
Title : cryoEM reconstruction of the HIV gp140 in complex with the extracellular domains of CD4 and the adnectin domain of Combnectin. The gp140 and CD4 coordinates from entry 6EDU were rigid body fitted to the EM map along with the crystal structure of CD4+adnectin  
Authors : Concha, N.O.; William, S.P.; Wenzel, D.L.  
Deposited on : 2021-11-30  
Resolution : 8.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49



# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 8.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-----------------------------|----------------------------------------------------------|
| Clashscore            | 229148                      | 23984                       | -                                                        |
| Ramachandran outliers | 224038                      | 23583                       | -                                                        |
| Sidechain outliers    | 223484                      | 23102                       | -                                                        |
| Q-score               | -                           | 25397                       | 282 ( 8.20 - 9.20 )                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain                                                         |
|-----|-------|--------|--------------------------------------------------------------------------|
| 1   | D     | 650    | <div> <div>12%</div> <div>40%</div> <div>14%</div> <div>45%</div> </div> |
| 1   | G     | 650    | <div> <div>13%</div> <div>38%</div> <div>15%</div> <div>46%</div> </div> |
| 1   | I     | 650    | <div> <div>12%</div> <div>39%</div> <div>14%</div> <div>45%</div> </div> |
| 2   | B     | 39     | <div> <div>10%</div> <div>77%</div> <div>13%</div> <div>10%</div> </div> |

Continued on next page...

*Continued from previous page...*

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 3   | F     | 38     |                  |
| 3   | J     | 38     |                  |
| 4   | H     | 370    |                  |
| 4   | L     | 370    |                  |
| 4   | M     | 370    |                  |
| 5   | O     | 108    |                  |
| 5   | P     | 108    |                  |
| 5   | Q     | 108    |                  |
| 6   | K     | 35     |                  |
| 6   | N     | 35     |                  |
| 6   | R     | 35     |                  |

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 19033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BG505 SOSIP.664 gp140.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1   | D     | 360      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2772  | 1734 | 487 | 527 | 24 |         |       |
| 1   | G     | 353      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2731  | 1713 | 478 | 516 | 24 |         |       |
| 1   | I     | 356      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2749  | 1722 | 482 | 521 | 24 |         |       |

- Molecule 2 is a protein called BG505 SOSIP.664 gp140.

| Mol | Chain | Residues | Atoms |     |    |    | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|-------|
| 2   | B     | 35       | Total | C   | N  | O  | 0       | 0     |
|     |       |          | 175   | 105 | 35 | 35 |         |       |

- Molecule 3 is a protein called BG505 SOSIP.664 gp140.

| Mol | Chain | Residues | Atoms |     |    |    | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|-------|
| 3   | F     | 37       | Total | C   | N  | O  | 0       | 0     |
|     |       |          | 185   | 111 | 37 | 37 |         |       |
| 3   | J     | 37       | Total | C   | N  | O  | 0       | 0     |
|     |       |          | 185   | 111 | 37 | 37 |         |       |

- Molecule 4 is a protein called T-cell surface glycoprotein CD4.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 4   | H     | 356      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2544  | 1617 | 419 | 498 | 10 |         |       |
| 4   | L     | 357      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2553  | 1613 | 420 | 510 | 10 |         |       |
| 4   | M     | 353      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2519  | 1592 | 413 | 504 | 10 |         |       |

There are 18 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment        | Reference  |
|-------|---------|----------|--------|----------------|------------|
| H     | 365     | HIS      | -      | expression tag | UNP P01730 |
| H     | 366     | HIS      | -      | expression tag | UNP P01730 |
| H     | 367     | HIS      | -      | expression tag | UNP P01730 |
| H     | 368     | HIS      | -      | expression tag | UNP P01730 |
| H     | 369     | HIS      | -      | expression tag | UNP P01730 |
| H     | 370     | HIS      | -      | expression tag | UNP P01730 |
| L     | 365     | HIS      | -      | expression tag | UNP P01730 |
| L     | 366     | HIS      | -      | expression tag | UNP P01730 |
| L     | 367     | HIS      | -      | expression tag | UNP P01730 |
| L     | 368     | HIS      | -      | expression tag | UNP P01730 |
| L     | 369     | HIS      | -      | expression tag | UNP P01730 |
| L     | 370     | HIS      | -      | expression tag | UNP P01730 |
| M     | 365     | HIS      | -      | expression tag | UNP P01730 |
| M     | 366     | HIS      | -      | expression tag | UNP P01730 |
| M     | 367     | HIS      | -      | expression tag | UNP P01730 |
| M     | 368     | HIS      | -      | expression tag | UNP P01730 |
| M     | 369     | HIS      | -      | expression tag | UNP P01730 |
| M     | 370     | HIS      | -      | expression tag | UNP P01730 |

- Molecule 5 is a protein called Adnectin.


| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 5   | O     | 95       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 711   | 457 | 118 | 135 | 1 |         |       |
| 5   | P     | 95       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 707   | 455 | 117 | 134 | 1 |         |       |
| 5   | Q     | 95       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 707   | 455 | 117 | 134 | 1 |         |       |

- Molecule 6 is a protein called BG505 SOSIP.664 gp140.


| Mol | Chain | Residues | Atoms |    |    |    | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---------|-------|
| 6   | K     | 33       | Total | C  | N  | O  | 0       | 0     |
|     |       |          | 165   | 99 | 33 | 33 |         |       |
| 6   | N     | 33       | Total | C  | N  | O  | 0       | 0     |
|     |       |          | 165   | 99 | 33 | 33 |         |       |
| 6   | R     | 33       | Total | C  | N  | O  | 0       | 0     |
|     |       |          | 165   | 99 | 33 | 33 |         |       |











- Molecule 2: BG505 SOSIP.664 gp140





- Molecule 3: BG505 SOSIP.664 gp140




- Molecule 3: BG505 SOSIP.664 gp140




- Molecule 4: T-cell surface glycoprotein CD4







• Molecule 4: T-cell surface glycoprotein CD4




• Molecule 4: T-cell surface glycoprotein CD4








- Molecule 5: Adnectin




- Molecule 5: Adnectin



- Molecule 5: Adnectin




- Molecule 6: BG505 SOSIP.664 gp140




- Molecule 6: BG505 SOSIP.664 gp140





- Molecule 6: BG505 SOSIP.664 gp140



## 4 Experimental information

| Property                             | Value                                   | Source    |
|--------------------------------------|-----------------------------------------|-----------|
| EM reconstruction method             | SINGLE PARTICLE                         | Depositor |
| Imposed symmetry                     | POINT, C3                               | Depositor |
| Number of particles used             | 208485                                  | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF                       | Depositor |
| CTF correction method                | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope                           | FEI TITAN KRIOS                         | Depositor |
| Voltage (kV)                         | 300                                     | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 1.022                                   | Depositor |
| Minimum defocus (nm)                 | 1400                                    | Depositor |
| Maximum defocus (nm)                 | 3400                                    | Depositor |
| Magnification                        | Not provided                            |           |
| Image detector                       | FEI FALCON III (4k x 4k)                | Depositor |
| Maximum map value                    | 28.005                                  | Depositor |
| Minimum map value                    | -5.139                                  | Depositor |
| Average map value                    | 0.625                                   | Depositor |
| Map value standard deviation         | 2.529                                   | Depositor |
| Recommended contour level            | 4.42                                    | Depositor |
| Map size (Å)                         | 243.7776, 229.00319, 123.119995         | wwPDB     |
| Map dimensions                       | 50, 93, 99                              | wwPDB     |
| Map angles (°)                       | 90.0, 90.0, 90.0                        | wwPDB     |
| Pixel spacing (Å)                    | 2.4624, 2.4624, 2.4624                  | Depositor |

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |         | Bond angles |                 |
|-----|-------|--------------|---------|-------------|-----------------|
|     |       | RMSZ         | # Z  >5 | RMSZ        | # Z  >5         |
| 1   | D     | 0.47         | 0/2828  | 0.85        | 5/3845 (0.1%)   |
| 1   | G     | 0.47         | 0/2785  | 0.84        | 5/3784 (0.1%)   |
| 1   | I     | 0.47         | 0/2804  | 0.85        | 5/3812 (0.1%)   |
| 4   | H     | 0.23         | 0/2589  | 0.54        | 0/3544          |
| 4   | L     | 0.23         | 0/2595  | 0.55        | 0/3551          |
| 4   | M     | 0.23         | 0/2559  | 0.55        | 0/3500          |
| 5   | O     | 0.22         | 0/731   | 0.60        | 0/1004          |
| 5   | P     | 0.20         | 0/727   | 0.52        | 0/999           |
| 5   | Q     | 0.20         | 0/727   | 0.52        | 0/999           |
| All | All   | 0.36         | 0/18345 | 0.70        | 15/25038 (0.1%) |

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms   | Z     | Observed(°) | Ideal(°) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 1   | D     | 240 | PRO  | CA-N-CD | -9.30 | 98.98       | 112.00   |
| 1   | G     | 240 | PRO  | CA-N-CD | -9.28 | 99.01       | 112.00   |
| 1   | I     | 240 | PRO  | CA-N-CD | -9.27 | 99.02       | 112.00   |
| 1   | G     | 417 | PRO  | CA-N-CD | -9.00 | 99.40       | 112.00   |
| 1   | D     | 291 | PRO  | CA-N-CD | -8.99 | 99.41       | 112.00   |

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | D     | 2772  | 0        | 2690     | 102     | 0            |
| 1   | G     | 2731  | 0        | 2658     | 100     | 0            |
| 1   | I     | 2749  | 0        | 2670     | 125     | 0            |
| 2   | B     | 175   | 0        | 37       | 7       | 0            |
| 3   | F     | 185   | 0        | 39       | 2       | 0            |
| 3   | J     | 185   | 0        | 39       | 30      | 0            |
| 4   | H     | 2544  | 0        | 2381     | 76      | 0            |
| 4   | L     | 2553  | 0        | 2386     | 73      | 0            |
| 4   | M     | 2519  | 0        | 2353     | 76      | 0            |
| 5   | O     | 711   | 0        | 674      | 15      | 0            |
| 5   | P     | 707   | 0        | 668      | 17      | 0            |
| 5   | Q     | 707   | 0        | 668      | 17      | 0            |
| 6   | K     | 165   | 0        | 36       | 0       | 0            |
| 6   | N     | 165   | 0        | 36       | 0       | 0            |
| 6   | R     | 165   | 0        | 36       | 0       | 0            |
| All | All   | 19033 | 0        | 17371    | 587     | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 587 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1          | Atom-2        | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|---------------|--------------------------|-------------------|
| 1:I:61:TYR:CB   | 3:J:1:UNK:H   | 1.15                     | 1.57              |
| 1:I:61:TYR:HB2  | 3:J:1:UNK:N   | 1.34                     | 1.43              |
| 1:D:221:ALA:HB1 | 3:J:33:UNK:CB | 1.54                     | 1.34              |
| 1:I:61:TYR:CD2  | 3:J:1:UNK:N   | 2.05                     | 1.24              |
| 1:I:61:TYR:CG   | 3:J:1:UNK:N   | 2.07                     | 1.22              |

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Favoured   | Allowed  | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 1   | D     | 352/650 (54%)   | 303 (86%)  | 41 (12%) | 8 (2%)   | 5           | 28  |
| 1   | G     | 341/650 (52%)   | 295 (86%)  | 39 (11%) | 7 (2%)   | 5           | 30  |
| 1   | I     | 346/650 (53%)   | 298 (86%)  | 40 (12%) | 8 (2%)   | 5           | 28  |
| 4   | H     | 352/370 (95%)   | 333 (95%)  | 17 (5%)  | 2 (1%)   | 21          | 59  |
| 4   | L     | 353/370 (95%)   | 335 (95%)  | 18 (5%)  | 0        | 100         | 100 |
| 4   | M     | 347/370 (94%)   | 329 (95%)  | 18 (5%)  | 0        | 100         | 100 |
| 5   | O     | 93/108 (86%)    | 87 (94%)   | 5 (5%)   | 1 (1%)   | 11          | 46  |
| 5   | P     | 93/108 (86%)    | 87 (94%)   | 5 (5%)   | 1 (1%)   | 11          | 46  |
| 5   | Q     | 93/108 (86%)    | 87 (94%)   | 5 (5%)   | 1 (1%)   | 11          | 46  |
| All | All   | 2370/3384 (70%) | 2154 (91%) | 188 (8%) | 28 (1%)  | 13          | 44  |

5 of 28 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | D     | 240 | PRO  |
| 1   | D     | 301 | ASN  |
| 1   | D     | 429 | ARG  |
| 1   | G     | 240 | PRO  |
| 1   | G     | 301 | ASN  |

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Rotameric  | Outliers | Percentiles |     |
|-----|-------|---------------|------------|----------|-------------|-----|
| 1   | D     | 312/568 (55%) | 312 (100%) | 0        | 100         | 100 |
| 1   | G     | 309/568 (54%) | 309 (100%) | 0        | 100         | 100 |
| 1   | I     | 310/568 (55%) | 310 (100%) | 0        | 100         | 100 |
| 4   | H     | 259/331 (78%) | 259 (100%) | 0        | 100         | 100 |
| 4   | L     | 263/331 (80%) | 263 (100%) | 0        | 100         | 100 |
| 4   | M     | 259/331 (78%) | 259 (100%) | 0        | 100         | 100 |
| 5   | O     | 72/91 (79%)   | 72 (100%)  | 0        | 100         | 100 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Analysed        | Rotameric   | Outliers | Percentiles |     |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 5   | P     | 71/91 (78%)     | 71 (100%)   | 0        | 100         | 100 |
| 5   | Q     | 71/91 (78%)     | 71 (100%)   | 0        | 100         | 100 |
| All | All   | 1926/2970 (65%) | 1926 (100%) | 0        | 100         | 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 4   | L     | 103 | ASN  |
| 4   | L     | 163 | GLN  |
| 4   | M     | 163 | GLN  |
| 1   | G     | 356 | ASN  |
| 1   | G     | 103 | GLN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-25581. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 4.42. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_25581\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 441 nm<sup>3</sup>; this corresponds to an approximate mass of 398 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.


## 7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.115 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


| Resolution estimate (Å)   | Estimation criterion (FSC cut-off) |      |          |
|---------------------------|------------------------------------|------|----------|
|                           | 0.143                              | 0.5  | Half-bit |
| Reported by author        | 8.70                               | -    | -        |
| Author-provided FSC curve | 8.69                               | 9.60 | 8.89     |
| Unmasked-calculated*      | -                                  | -    | -        |

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-25581 and PDB model 7T0O. Per-residue inclusion information can be found in section [3](#) on page [6](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 4.42 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (4.42).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 64% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (4.42) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion                                                                             | Q-score                                                                                    |
|-------|--------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| All   |  0.6190   |  0.0370   |
| B     |  0.8690   |  0.0560   |
| D     |  0.7570   |  0.0340   |
| F     |  0.8860   |  0.0490   |
| G     |  0.7440   |  0.0300   |
| H     |  0.5280   |  0.0400   |
| I     |  0.7500   |  0.0330   |
| J     |  0.8760   |  0.0240   |
| K     |  0.9090   |  0.2490   |
| L     |  0.4780   |  0.0290   |
| M     |  0.4760   |  0.0260   |
| N     |  0.7330   |  0.1970   |
| O     |  0.4820   |  0.0340   |
| P     |  0.4090   |  0.0240   |
| Q     |  0.4370  |  0.0400  |
| R     |  0.8000 |  0.2010 |

