



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 08:44 PM UTC

PDB ID : 6T1F / pdb_00006t1f
Title : Crystal structure of the C-terminally truncated chromosome-partitioning protein ParB from *Caulobacter crescentus* complexed to the centromeric parS site
Authors : Jalal, A.S.B.; Pastrana, C.L.; Tran, N.T.; Stevenson, C.E.M.; Lawson, D.M.; Moreno-Herrero, F.; Le, T.B.K.
Deposited on : 2019-10-04
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	2% 61% 11% 26%
1	B	257	3% 63% 10% 26%
1	C	257	0% 63% 12% 24%
1	D	257	9% 64% 8% 27%
2	E	22	77% 23%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	22	 77% 23%
2	G	22	 73% 27%
2	H	22	 82% 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromosome-partitioning protein ParB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1376	858	251	264	3			
1	B	190	Total	C	N	O	S	0	0	0
			1312	820	234	256	2			
1	C	195	Total	C	N	O	S	0	0	0
			1437	899	263	272	3			
1	D	187	Total	C	N	O	S	0	0	0
			1246	772	232	240	2			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	255	LYS	-	expression tag	UNP B8GW30
A	256	LEU	-	expression tag	UNP B8GW30
A	257	ALA	-	expression tag	UNP B8GW30
A	258	ALA	-	expression tag	UNP B8GW30
A	259	ALA	-	expression tag	UNP B8GW30
A	260	LEU	-	expression tag	UNP B8GW30
A	261	GLU	-	expression tag	UNP B8GW30
A	262	HIS	-	expression tag	UNP B8GW30
A	263	HIS	-	expression tag	UNP B8GW30
A	264	HIS	-	expression tag	UNP B8GW30
A	265	HIS	-	expression tag	UNP B8GW30
A	266	HIS	-	expression tag	UNP B8GW30
A	267	HIS	-	expression tag	UNP B8GW30
B	255	LYS	-	expression tag	UNP B8GW30
B	256	LEU	-	expression tag	UNP B8GW30
B	257	ALA	-	expression tag	UNP B8GW30
B	258	ALA	-	expression tag	UNP B8GW30
B	259	ALA	-	expression tag	UNP B8GW30
B	260	LEU	-	expression tag	UNP B8GW30
B	261	GLU	-	expression tag	UNP B8GW30
B	262	HIS	-	expression tag	UNP B8GW30

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	263	HIS	-	expression tag	UNP B8GW30
B	264	HIS	-	expression tag	UNP B8GW30
B	265	HIS	-	expression tag	UNP B8GW30
B	266	HIS	-	expression tag	UNP B8GW30
B	267	HIS	-	expression tag	UNP B8GW30
C	255	LYS	-	expression tag	UNP B8GW30
C	256	LEU	-	expression tag	UNP B8GW30
C	257	ALA	-	expression tag	UNP B8GW30
C	258	ALA	-	expression tag	UNP B8GW30
C	259	ALA	-	expression tag	UNP B8GW30
C	260	LEU	-	expression tag	UNP B8GW30
C	261	GLU	-	expression tag	UNP B8GW30
C	262	HIS	-	expression tag	UNP B8GW30
C	263	HIS	-	expression tag	UNP B8GW30
C	264	HIS	-	expression tag	UNP B8GW30
C	265	HIS	-	expression tag	UNP B8GW30
C	266	HIS	-	expression tag	UNP B8GW30
C	267	HIS	-	expression tag	UNP B8GW30
D	255	LYS	-	expression tag	UNP B8GW30
D	256	LEU	-	expression tag	UNP B8GW30
D	257	ALA	-	expression tag	UNP B8GW30
D	258	ALA	-	expression tag	UNP B8GW30
D	259	ALA	-	expression tag	UNP B8GW30
D	260	LEU	-	expression tag	UNP B8GW30
D	261	GLU	-	expression tag	UNP B8GW30
D	262	HIS	-	expression tag	UNP B8GW30
D	263	HIS	-	expression tag	UNP B8GW30
D	264	HIS	-	expression tag	UNP B8GW30
D	265	HIS	-	expression tag	UNP B8GW30
D	266	HIS	-	expression tag	UNP B8GW30
D	267	HIS	-	expression tag	UNP B8GW30

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*AP*TP*GP*TP*TP*TP*CP*AP*CP*GP*TP*GP*AP*AP*AP*CP*AP*TP*CP*C)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	22	Total	C	N	O	P	0	0	0
			448	215	82	130	21			
2	F	22	Total	C	N	O	P	0	0	0
			448	215	82	130	21			
2	G	22	Total	C	N	O	P	0	0	0
			448	215	82	130	21			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	22	Total 448	C 215	N 82	O 130	P 21	0	0	0


- Molecule 1: Chromosome-partitioning protein ParB




- Molecule 2: DNA (5'-D(*GP*GP*AP*TP*GP*TP*TP*TP*CP*AP*CP*GP*TP*GP*AP*AP*AP*CP*AP*TP*CP*C)-3')




- Molecule 2: DNA (5'-D(*GP*GP*AP*TP*GP*TP*TP*TP*CP*AP*CP*GP*TP*GP*AP*AP*AP*CP*AP*TP*CP*C)-3')



- Molecule 2: DNA (5'-D(*GP*GP*AP*TP*GP*TP*TP*TP*CP*AP*CP*GP*TP*GP*AP*AP*AP*CP*AP*TP*CP*C)-3')



- Molecule 2: DNA (5'-D(*GP*GP*AP*TP*GP*TP*TP*TP*CP*AP*CP*GP*TP*GP*AP*AP*AP*CP*AP*TP*CP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.25Å 172.93Å 72.85Å 90.00° 90.54° 90.00°	Depositor
Resolution (Å)	72.96 – 2.90 72.96 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (72.96-2.90) 99.6 (72.96-2.90)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.239 , 0.263 0.241 , 0.265	Depositor DCC
R_{free} test set	1466 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	78.6	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.108 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7163	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/1391	1.03	0/1892
1	B	0.65	0/1329	1.03	0/1820
1	C	0.65	0/1454	1.03	0/1975
1	D	0.66	0/1261	1.03	0/1731
2	E	0.41	0/502	0.87	1/773 (0.1%)
2	F	0.45	0/502	0.86	0/773
2	G	0.44	0/502	0.90	1/773 (0.1%)
2	H	0.42	0/502	0.86	0/773
All	All	0.60	0/7443	0.99	2/10510 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
1	C	0	2
1	D	0	4
All	All	0	12

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	DG	C2'-C3'-O3'	-5.53	103.21	111.50
2	E	2	DG	C2'-C3'-O3'	-5.05	103.93	111.50

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	139	ARG	Sidechain
1	A	160	ARG	Sidechain
1	A	173	ARG	Sidechain
1	A	54	ARG	Sidechain
1	B	104	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1376	0	1317	15	0
1	B	1312	0	1171	17	0
1	C	1437	0	1418	24	0
1	D	1246	0	1077	14	0
2	E	448	0	250	4	0
2	F	448	0	250	4	0
2	G	448	0	250	5	0
2	H	448	0	250	4	0
All	All	7163	0	5983	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:51:ILE:HG22	1:C:95:GLU:HA	1.46	0.92
1:B:199:THR:HG22	1:B:202:HIS:ND1	1.88	0.89
1:D:199:THR:HG22	1:D:202:HIS:ND1	1.87	0.88
1:B:219:ILE:HG12	1:B:229:THR:HG22	1.59	0.83
1:D:219:ILE:HG12	1:D:229:THR:HG22	1.67	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/257 (73%)	177 (95%)	10 (5%)	0	100	100
1	B	186/257 (72%)	179 (96%)	7 (4%)	0	100	100
1	C	193/257 (75%)	182 (94%)	11 (6%)	0	100	100
1	D	183/257 (71%)	175 (96%)	8 (4%)	0	100	100
All	All	749/1028 (73%)	713 (95%)	36 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/203 (62%)	115 (91%)	11 (9%)	9	30
1	B	108/203 (53%)	99 (92%)	9 (8%)	10	32
1	C	137/203 (68%)	133 (97%)	4 (3%)	37	71
1	D	95/203 (47%)	89 (94%)	6 (6%)	16	45
All	All	466/812 (57%)	436 (94%)	30 (6%)	16	44

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	154	LEU
1	D	164	ASN
1	B	193	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	227	ARG
1	D	54	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	136	ASN
1	D	55	ASN
1	D	82	GLN
1	A	167	GLN
1	A	73	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/257 (74%)	0.29	4 (2%) 63 54	64, 92, 126, 147	0
1	B	190/257 (73%)	0.55	7 (3%) 45 37	67, 100, 130, 136	0
1	C	195/257 (75%)	0.22	2 (1%) 79 73	59, 85, 123, 140	0
1	D	187/257 (72%)	1.01	22 (11%) 9 7	88, 115, 138, 149	0
2	E	22/22 (100%)	-0.59	0 100 100	58, 68, 89, 92	0
2	F	22/22 (100%)	-0.47	0 100 100	59, 69, 84, 90	0
2	G	22/22 (100%)	-0.31	0 100 100	62, 71, 93, 98	0
2	H	22/22 (100%)	-0.24	0 100 100	56, 75, 92, 104	0
All	All	851/1116 (76%)	0.42	35 (4%) 41 33	56, 95, 132, 149	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	77	GLU	4.3
1	D	207	ALA	3.8
1	D	107	ALA	3.6
1	B	81	LEU	3.2
1	D	151	TYR	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.