



wwPDB EM Validation Summary Report ⓘ

Mar 28, 2026 – 04:00 AM UTC

PDB ID : 8T1K / pdb_00008t1k
EMDB ID : EMD-40970
Title : DSBU crosslinked nNOS-CaM oxygenase homodimer
Authors : Lee, K.; Pospiech, T.H.; Southworth, D.
Deposited on : 2023-06-02
Resolution : 3.14 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14483 (2.64 - 3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1429	 28% 71%
1	B	1429	 27% 71%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	419	Total	C	N	O	S	0	0
			3414	2181	588	624	21		
1	B	419	Total	C	N	O	S	0	0
			3414	2181	588	624	21		

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
2	A	1	Total	Zn	0
			1	1	

- Molecule 3 is ARGinine (CCD ID: ARG) (formula: C₆H₁₅N₄O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	N	O	0
			12	6	4	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
3	B	1	12	6	4	2	0

- Molecule 4 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	A	1	17	9	5	3	0
4	B	1	17	9	5	3	0


- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
5	A	1	Total 43	C 34	Fe 1	N 4	O 4	0
5	B	1	Total 43	C 34	Fe 1	N 4	O 4	0

[illegible]

- Molecule 1: Nitric oxide synthase 1

Chain B:  27% 71%

[illegible]

PHE	ILE	SER	ARG	LEU	ARG	ASP	ASP	GLN	TRP	ARG	GLY	LYS	LYS	ASP	THR	ASP
SER	LEU	ARG	VAL	GLN	GLY	ASP	GLY	GLN	GLN	GLY	ASN	GLY	GLY	ASN	LEU	ASP
LEU	LEU	GLN	GLN	GLN	PRO	PRO	GLY	ASN	ASN	ASN	SER	SER	TRP	SER	GLU	LEU
ARG	ARG	ASP	ASP	PHE	ASP	PHE	PRO	THR	THR	THR	LEU	LEU	LEU	LYS	GLU	GLU
ASP	ASP	VAL	VAL	VAL	VAL	VAL	MET	MET	MET	MET	PRO	GLN	ARG	ARG	LEU	LEU
ASN	ASN	GLN	GLN	HIS	HIS	HIS	GLY	GLY	GLY	GLY	PRO	GLN	TYR	LYS	GLY	GLY
ARG	ARG	GLU	GLN	GLN	VAL	VAL	LEU	VAL	VAL	VAL	THR	ILE	PRO	ARG	GLY	GLU
TYR	TYR	GLN	GLN	GLY	CYS	CYS	GLY	GLY	GLY	GLY	ASP	GLY	GLY	GLY	ARG	ARG
HIS	HIS	LEU	ALA	ALA	MET	MET	SER	GLU	GLU	GLU	PHE	PHE	ASP	ILE	LEU	ILE
GLU	GLU	ASP	GLU	GLU	ASN	ASN	SER	PHE	PHE	PHE	ALA	GLN	HIS	THR	LEU	LEU
ASP	ASP	ILE	ILE	ILE	SER	PRO	TRP	PRO	PRO	PRO	GLY	GLY	LEU	TYR	LYS	LYS
PHE	PHE	GLY	VAL	VAL	VAL	CYS	LEU	SER	ILE	ILE	ASN	LYS	VAL	ALA	ARG	ARG
VAL	VAL	THR	THR	THR	ILE	PRO	ARG	GLN	GLN	GLN	THR	TTR	PRO	ALA	GLY	GLY
THR	THR	LEU	ALA	ALA	VAL	VAL	ILE	ILE	ILE	ILE	LEU	LEU	PRO	ASP	ASP	ASP
LEU	LEU	ARG	TYR	TYR	GLY	GLY	ASP	THR	THR	THR	GLY	THR	GLY	LEU	LEU	LEU
ARG	ARG	VAL	VAL	VAL	CYS	CYS	VAL	LEU	LEU	LEU	ASP	PRO	GLY	GLY	GLY	GLY
SER	SER	GLY	GLY	GLY	ARG	GLY	GLY	LEU	LEU	LEU	LEU	GLN	LEU	LEU	LEU	LEU
GLU	GLU	ASP	ASP	ASP	ILE	ILE	GLY	SER	THR	THR	ASN	ASN	SER	SER	GLU	GLU
ALA	ALA	TYR	TYR	TYR	PHE	PHE	VAL	LEU	LEU	LEU	ALA	ALA	ALA	ALA	ALA	ALA
PHE	PHE	GLU	GLU	GLU	GLU	GLU	ARG	SER	SER	SER	ALA	ALA	ALA	ALA	ALA	ALA
ILE	ILE	GLU	GLU	GLU	THR	THR	GLY	THR	THR	THR	PRO	PRO	PRO	PRO	PRO	PRO
GLU	GLU	VAL	VAL	VAL	ASP	ASP	GLY	GLY	GLY	GLY	ASN	ASN	ASN	ASN	ASN	ASN
SER	SER	VAL	VAL	VAL	ILE	ILE	VAL	LEU	LEU	LEU	GLY	GLY	GLY	GLY	GLY	GLY
LYS	LYS	GLY	GLY	GLY	GLY	GLY	VAL	SER	SER	SER	ILE	ILE	ILE	ILE	ILE	ILE
LYS	LYS	ASP	ASP	ASP	ASP	ASP	ARG	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
ASP	ASP	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
ASP	ASP	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL
PHE	PHE	THR	THR	THR	THR	THR	VAL	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
SER	SER	GLN	GLN	GLN	GLN	GLN	GLY	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL
SER	SER	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY
SER	SER	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR					

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	294924	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	62.00	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	8.752	Depositor
Minimum map value	-6.717	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.141	Depositor
Recommended contour level	1.8	Depositor
Map size ($\text{\AA}$)	310.08, 310.08, 310.08	wwPDB
Map dimensions	340, 340, 340	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.91199994, 0.91199994, 0.91199994	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, H4B, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/3511	0.97	12/4764 (0.3%)
1	B	0.66	2/3511 (0.1%)	0.99	12/4764 (0.3%)
All	All	0.64	2/7022 (0.0%)	0.98	24/9528 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	588	TYR	C-O	-7.97	1.14	1.23
1	B	588	TYR	CA-C	-7.94	1.42	1.52

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	588	TYR	CA-C-O	-8.14	111.77	120.80
1	B	354	LEU	N-CA-C	7.12	119.12	111.36
1	A	354	LEU	N-CA-C	7.06	119.06	111.36
1	A	712	ASN	N-CA-C	-6.04	105.56	113.17
1	B	712	ASN	N-CA-C	-5.99	105.62	113.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3414	0	3322	7	0
1	B	3414	0	3322	11	0
2	A	1	0	0	0	0
3	A	12	0	12	1	0
3	B	12	0	12	0	0
4	A	17	0	15	1	0
4	B	17	0	15	1	0
5	A	43	0	30	0	0
5	B	43	0	30	0	0
All	All	6973	0	6758	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:478:GLN:HE22	3:A:1502:ARG:C	2.19	0.51
1:B:478:GLN:OE1	1:B:481:ARG:NH2	2.47	0.48
1:B:359:LYS:NZ	1:B:363:ASP:OD2	2.43	0.47
1:A:715:VAL:O	1:A:716:TRP:HB2	2.16	0.46
1:B:462:PHE:HB3	1:B:463:PRO:HD2	1.98	0.45

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/1429 (29%)	407 (98%)	10 (2%)	0	100	100
1	B	417/1429 (29%)	405 (97%)	12 (3%)	0	100	100
All	All	834/2858 (29%)	812 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/1241 (30%)	375 (100%)	0	100	100
1	B	375/1241 (30%)	375 (100%)	0	100	100
All	All	750/2482 (30%)	750 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	478	GLN
1	A	525	GLN
1	A	527	ASN
1	B	527	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	H4B	A	1503	-	17,18,18	1.68	5 (29%)	14,26,26	1.90	3 (21%)
4	H4B	B	1502	-	17,18,18	1.49	4 (23%)	14,26,26	1.91	2 (14%)
3	ARG	B	1501	-	10,11,11	0.65	0	9,13,13	0.36	0
5	HEM	B	1503	-	50,50,50	1.56	8 (16%)	67,82,82	1.30	7 (10%)
5	HEM	A	1504	1	50,50,50	1.51	4 (8%)	67,82,82	1.16	8 (11%)
3	ARG	A	1502	-	10,11,11	0.61	0	9,13,13	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	H4B	A	1503	-	-	0/8/17/17	0/2/2/2
4	H4B	B	1502	-	-	0/8/17/17	0/2/2/2
3	ARG	B	1501	-	-	0/11/11/11	-
5	HEM	B	1503	-	-	3/14/54/54	-
5	HEM	A	1504	1	-	4/14/54/54	-
3	ARG	A	1502	-	-	4/11/11/11	-

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1504	HEM	FE-NC	6.74	2.17	1.95
5	B	1503	HEM	FE-NC	5.85	2.14	1.95
4	A	1503	H4B	C4A-N5	3.90	1.47	1.37
5	A	1504	HEM	FE-NA	3.16	2.05	1.95
5	B	1503	HEM	FE-NA	3.16	2.05	1.95

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1503	H4B	C2-N1-C8A	5.18	122.50	113.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1502	H4B	C2-N1-C8A	5.16	122.47	113.36
5	A	1504	HEM	CHD-C4C-NC	-3.61	120.52	124.45
5	B	1503	HEM	C3B-C2B-C1B	3.38	108.95	106.41
4	B	1502	H4B	N3-C2-N1	-3.34	117.21	123.32

There are no chirality outliers.

5 of 11 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	1502	ARG	O-C-CA-N
5	A	1504	HEM	C2B-C3B-CAB-CBB
5	A	1504	HEM	C4B-C3B-CAB-CBB
5	B	1503	HEM	C2B-C3B-CAB-CBB
3	A	1502	ARG	OXT-C-CA-N



There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1503	H4B	1	0
4	B	1502	H4B	1	0
3	A	1502	ARG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Ligand HEM A 1504



Ligand ARG A 1502



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-40970. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 170




Y Index: 170




Z Index: 170


6.2.2 Raw map



X Index: 170



Y Index: 170




Z Index: 170


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 170




Y Index: 171




Z Index: 196


6.3.2 Raw map



X Index: 180



Y Index: 173




Z Index: 196


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.8. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 26 nm³; this corresponds to an approximate mass of 24 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.318 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.318 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.14	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.37	3.73	3.41

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-40970 and PDB model 8T1K. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 1.8 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.8).

9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (1.8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7180	0.5640
A	0.7190	0.5640
B	0.7170	0.5640

