



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:10 PM UTC

PDB ID : 8T31 / pdb_00008t31
Title : Crystal structure of GABARAP in complex with the LIR of TP53INP2/DOR
Authors : Ali, M.G.H.; Wahba, H.M.; Cyr, N.; Omichinski, J.G.
Deposited on : 2023-06-07
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	91% 5% .
1	C	119	% 85% 13% .
1	E	119	% 84% 14% .
1	G	119	2% 80% 18% .
1	I	119	% 81% 15% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	15	 80% 13% 7%
2	D	15	 27% 33% 40%
2	F	15	 73% 13% 13%
2	H	15	 40% 60%
2	J	15	 20% 13% 67%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid receptor-associated protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	114	Total	C	N	O	S	0	0	0
			962	627	163	170	2			
1	C	116	Total	C	N	O	S	0	0	0
			978	638	165	173	2			
1	E	117	Total	C	N	O	S	0	0	0
			986	644	166	174	2			
1	G	116	Total	C	N	O	S	0	0	0
			978	638	165	173	2			
1	I	114	Total	C	N	O	S	0	0	0
			962	627	163	170	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP O95166
A	0	SER	-	expression tag	UNP O95166
C	-1	GLY	-	expression tag	UNP O95166
C	0	SER	-	expression tag	UNP O95166
E	-1	GLY	-	expression tag	UNP O95166
E	0	SER	-	expression tag	UNP O95166
G	-1	GLY	-	expression tag	UNP O95166
G	0	SER	-	expression tag	UNP O95166
I	-1	GLY	-	expression tag	UNP O95166
I	0	SER	-	expression tag	UNP O95166

- Molecule 2 is a protein called Tumor protein p53-inducible nuclear protein 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	0	0	0
			107	69	15	23			
2	D	9	Total	C	N	O	0	0	0
			74	49	10	15			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	13	Total	C	N	O	0	0	0
			99	65	14	20			
2	H	6	Total	C	N	O	0	0	0
			50	35	7	8			
2	J	5	Total	C	N	O	0	0	0
			46	33	6	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	29	GLY	-	expression tag	UNP Q8IXH6
B	30	SER	-	expression tag	UNP Q8IXH6
D	29	GLY	-	expression tag	UNP Q8IXH6
D	30	SER	-	expression tag	UNP Q8IXH6
F	29	GLY	-	expression tag	UNP Q8IXH6
F	30	SER	-	expression tag	UNP Q8IXH6
H	29	GLY	-	expression tag	UNP Q8IXH6
H	30	SER	-	expression tag	UNP Q8IXH6
J	29	GLY	-	expression tag	UNP Q8IXH6
J	30	SER	-	expression tag	UNP Q8IXH6

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total	O	0	0
			39	39		
3	B	7	Total	O	0	0
			7	7		
3	C	28	Total	O	0	0
			28	28		
3	E	22	Total	O	0	0
			22	22		
3	F	7	Total	O	0	0
			7	7		
3	G	14	Total	O	0	0
			14	14		
3	I	11	Total	O	0	0
			11	11		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-aminobutyric acid receptor-associated protein

Chain A: 



- Molecule 1: Gamma-aminobutyric acid receptor-associated protein

Chain C: 



- Molecule 1: Gamma-aminobutyric acid receptor-associated protein

Chain E: 



- Molecule 1: Gamma-aminobutyric acid receptor-associated protein

Chain G: 



- Molecule 1: Gamma-aminobutyric acid receptor-associated protein

Chain I: 





- Molecule 2: Tumor protein p53-inducible nuclear protein 2

Chain B:  80% 13% 7%




- Molecule 2: Tumor protein p53-inducible nuclear protein 2

Chain D:  27% 33% 40%




- Molecule 2: Tumor protein p53-inducible nuclear protein 2

Chain F:  73% 13% 13%




- Molecule 2: Tumor protein p53-inducible nuclear protein 2

Chain H:  40% 60%



- Molecule 2: Tumor protein p53-inducible nuclear protein 2

Chain J:  20% 13% 67%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.90Å 95.78Å 138.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.25 – 2.10 26.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	90.1 (26.25-2.10) 86.3 (26.25-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.75 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.197 , 0.264 0.197 , 0.263	Depositor DCC
R_{free} test set	2000 reflections (5.37%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5370	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/988	0.62	0/1329
1	C	0.40	0/1005	0.63	0/1352
1	E	0.40	0/1013	0.71	0/1363
1	G	0.33	0/1005	0.55	0/1352
1	I	0.32	0/988	0.56	0/1329
2	B	0.39	0/109	0.63	0/149
2	D	0.30	0/75	0.59	0/102
2	F	0.31	0/101	0.60	0/138
2	H	0.29	0/51	0.40	0/69
2	J	0.26	0/47	0.45	0/64
All	All	0.37	0/5382	0.62	0/7247

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	962	0	965	5	0
1	C	978	0	977	13	1
1	E	986	0	988	13	0
1	G	978	0	977	11	1
1	I	962	0	965	12	0
2	B	107	0	98	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	74	0	68	5	0
2	F	99	0	94	2	0
2	H	50	0	49	0	0
2	J	46	0	46	2	0
3	A	39	0	0	2	0
3	B	7	0	0	0	0
3	C	28	0	0	1	0
3	E	22	0	0	1	0
3	F	7	0	0	2	0
3	G	14	0	0	0	0
3	I	11	0	0	2	0
All	All	5370	0	5227	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:39:ASP:OD1	3:F:101:HOH:O	1.92	0.85
1:E:88:SER:OG	3:E:201:HOH:O	2.06	0.73
1:E:86:PRO:HG2	1:E:89:ALA:HB2	1.73	0.70
1:E:65:ARG:NH1	1:E:74:ASP:O	2.25	0.67
3:A:207:HOH:O	1:C:1:MET:SD	2.53	0.66

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:112:GLU:OE2	1:G:88:SER:OG[2_554]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/119 (94%)	111 (99%)	1 (1%)	0	100	100
1	C	114/119 (96%)	114 (100%)	0	0	100	100
1	E	115/119 (97%)	115 (100%)	0	0	100	100
1	G	114/119 (96%)	113 (99%)	1 (1%)	0	100	100
1	I	112/119 (94%)	111 (99%)	1 (1%)	0	100	100
2	B	12/15 (80%)	12 (100%)	0	0	100	100
2	D	7/15 (47%)	7 (100%)	0	0	100	100
2	F	11/15 (73%)	11 (100%)	0	0	100	100
2	H	4/15 (27%)	4 (100%)	0	0	100	100
2	J	3/15 (20%)	3 (100%)	0	0	100	100
All	All	604/670 (90%)	601 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/107 (97%)	104 (100%)	0	100	100
1	C	105/107 (98%)	105 (100%)	0	100	100
1	E	106/107 (99%)	104 (98%)	2 (2%)	50	58
1	G	105/107 (98%)	104 (99%)	1 (1%)	68	76
1	I	104/107 (97%)	102 (98%)	2 (2%)	50	58
2	B	12/13 (92%)	12 (100%)	0	100	100
2	D	8/13 (62%)	8 (100%)	0	100	100
2	F	11/13 (85%)	11 (100%)	0	100	100
2	H	5/13 (38%)	5 (100%)	0	100	100
2	J	5/13 (38%)	4 (80%)	1 (20%)	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	565/600 (94%)	559 (99%)	6 (1%)	65 74

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	40	ARG
1	I	88	SER
2	J	37	ILE
1	E	94	LEU
1	E	51	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	96	GLN
1	G	9	HIS
1	I	9	HIS
1	G	69	HIS
1	C	9	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	114/119 (95%)	-0.37	0 100 100	28, 42, 71, 84	0
1	C	116/119 (97%)	-0.01	1 (0%) 81 83	37, 53, 78, 93	0
1	E	117/119 (98%)	-0.14	1 (0%) 81 83	31, 47, 76, 96	0
1	G	116/119 (97%)	0.18	2 (1%) 69 71	42, 64, 94, 111	0
1	I	114/119 (95%)	0.16	1 (0%) 81 83	42, 65, 87, 103	0
2	B	14/15 (93%)	-0.31	0 100 100	35, 47, 63, 75	0
2	D	9/15 (60%)	0.57	0 100 100	55, 64, 79, 87	0
2	F	13/15 (86%)	0.09	0 100 100	38, 50, 68, 80	0
2	H	6/15 (40%)	0.95	0 100 100	67, 73, 76, 76	0
2	J	5/15 (33%)	0.90	0 100 100	74, 76, 84, 99	0
All	All	624/670 (93%)	-0.01	5 (0%) 82 84	28, 55, 85, 111	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	115	TYR	3.8
1	I	61	TYR	3.4
1	E	68	ILE	3.3
1	C	1	MET	2.2
1	G	1	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.