



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 28, 2026 – 08:27 AM UTC

PDB ID : 1T4L / pdb_00001t4l
Title : Solution structure of double-stranded RNA binding domain of *S. cerevisiae* RNase III (Rnt1p) in complex with the 5' terminal RNA hairpin of snR47 precursor
Authors : Wu, H.; Henras, A.; Chanfreau, G.; Feigon, J.
Deposited on : 2004-04-29

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	32	 88% 12%
2	B	90	 9% 49% 29% • 12%

2 Ensemble composition and analysis

This entry contains 15 models. Model 5 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:367-B:445 (79)	0.76	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 5, 6, 12
2	2, 7, 10, 11, 15
3	4, 8, 9, 13
Single-model clusters	14

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2437 atoms, of which 1067 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5' terminal hairpin of snR47 precursor.

Mol	Chain	Residues	Atoms						Trace
1	A	32	Total	C	H	N	O	P	0
			1028	306	346	124	221	31	

- Molecule 2 is a protein called Ribonuclease III.

Mol	Chain	Residues	Atoms						Trace
2	B	90	Total	C	H	N	O	S	0
			1409	429	721	129	127	3	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	364	GLY	-	cloning artifact	UNP Q02555
B	365	SER	-	cloning artifact	UNP Q02555

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5' terminal hairpin of snR47 precursor

Chain A: 

G1 G2 G3 G4 G5 G6 G7 G8 G9 G10 G11 G12 G13 G14 A15 A16 A17 A18 A19 A20 A21 A22 G23 G24 G25 G26 G27 U28 A29 U30 C31 U32 C32

- Molecule 2: Ribonuclease III

Chain B: 


G364 G365 G366 G367 G368 G369 G370 G371 G372 G373 G374 G375 G376 G377 G378 G379 G380 A381 A382 A383 A384 A385 A386 A387 A388 A389 A390 A391 A392 A393 A394 A395 A396 A397 A398 A399 A400 A401 A402 A403 A404 A405 A406 A407 A408 A409 A410 A411 A412 A413 A414 A415 A416 A417 A418 A419 A420 A421 A422 A423

G424 G425 G426 G427 G428 G429 G430 G431 G432 G433 G434 G435 G436 G437 G438 G439 G440 G441 G442 G443 G444 G445 G446 G447 G448 G449 G450 G451 G452 G453

4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: 5' terminal hairpin of snR47 precursor

Chain A: 

G1 G2 G3 G4 G5 G6 G7 G8 G9 G10 G11 G12 G13 G14 A15 A16 A17 A18 A19 A20 A21 A22 G23 G24 G25 G26 G27 U28 A29 U30 C31 U32 C32

- Molecule 2: Ribonuclease III

Chain B: 

G364 G365 G366 G367 G368 G369 G370 G371 G372 G373 G374 G375 G376 G377 G378 G379 G380 A381 A382 A383 A384 A385 A386 A387 A388 A389 A390 A391 A392 A393 A394 A395 A396 A397 A398 A399 A400 A401 A402 A403 A404 A405 A406 A407 A408 A409 A410 A411 A412 A413 A414 A415 A416 A417 A418 A419 A420 A421 A422 A423 G424

I425	R426	A427	A428	E429	I430	A431	L432	R433	D434	K435	K436	I437	L438	D439	F440	Y441	A442	K443	Q444	R445	A446	A447	I448	P449	R450	S451	E452	S453
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	NIH

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	682	346	346	90±10
2	B	612	643	642	111±10
All	All	19410	14835	14820	2932

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 86.

5 of 908 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:390:VAL:HG23	2:B:401:ILE:HG23	1.04	1.29	6	14
2:B:406:VAL:HG13	2:B:412:LEU:HD21	1.02	1.20	15	3
2:B:380:TYR:CE2	2:B:383:LEU:HD12	1.00	1.91	9	2
1:A:30:C:O2'	1:A:31:U:H5'	0.95	1.61	11	15
2:B:377:LEU:O	2:B:378:ILE:HD13	0.93	1.62	8	12

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	79/90 (88%)	56±3 (70±3%)	16±2 (20±3%)	8±2 (10±3%)	1	9
All	All	1185/1350 (88%)	834 (70%)	233 (20%)	118 (10%)	1	9

5 of 23 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	420	ILE	15
2	B	391	LYS	11
2	B	397	ASP	10
2	B	381	ALA	10
2	B	370	ALA	9

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	64/72 (89%)	36±3 (56±5%)	28±3 (44±5%)	0	3
All	All	960/1080 (89%)	539 (56%)	421 (44%)	0	3

5 of 54 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	388	VAL	15
2	B	402	VAL	15
2	B	412	LEU	15
2	B	420	ILE	15
2	B	421	LYS	15

6.3.3 RNA [i](#)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	31/32 (97%)	4±0 (13±0%)	2±1 (6±3%)	0.13±0.00
All	All	466/480 (97%)	60 (13%)	30 (6%)	0.13

The overall RNA backbone suiteness is 0.11.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	15	A	15
1	A	16	G	15
1	A	17	A	15
1	A	18	A	15

5 of 8 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	16	G	11
1	A	17	A	8
1	A	28	A	3
1	A	25	G	2
1	A	3	G	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided