



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:27 PM UTC

PDB ID : 5T4G / pdb_00005t4g
Title : Crystal structure of BhGH81 in complex with laminarin
Authors : Pluvinage, B.; Boraston, A.B.
Deposited on : 2016-08-29
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	783	90% 6%
2	B	3	100%
3	C	2	50% 50%
4	D	10	100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycoside Hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	752	Total	C	N	O	S	5	15	0
			6142	3898	1040	1186	18			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	initiating methionine	UNP Q9KG76
A	8	GLY	-	expression tag	UNP Q9KG76
A	9	SER	-	expression tag	UNP Q9KG76
A	10	SER	-	expression tag	UNP Q9KG76
A	11	HIS	-	expression tag	UNP Q9KG76
A	12	HIS	-	expression tag	UNP Q9KG76
A	13	HIS	-	expression tag	UNP Q9KG76
A	14	HIS	-	expression tag	UNP Q9KG76
A	15	HIS	-	expression tag	UNP Q9KG76
A	16	HIS	-	expression tag	UNP Q9KG76
A	17	SER	-	expression tag	UNP Q9KG76
A	18	SER	-	expression tag	UNP Q9KG76
A	19	GLY	-	expression tag	UNP Q9KG76
A	20	LEU	-	expression tag	UNP Q9KG76
A	21	VAL	-	expression tag	UNP Q9KG76
A	22	PRO	-	expression tag	UNP Q9KG76
A	23	ARG	-	expression tag	UNP Q9KG76
A	24	GLY	-	expression tag	UNP Q9KG76
A	25	SER	-	expression tag	UNP Q9KG76
A	26	HIS	-	expression tag	UNP Q9KG76
A	27	MET	-	expression tag	UNP Q9KG76

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	10	Total	C	O	0	0	0
			111	60	51			

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	657	Total	O	0	0
			657	657		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.40Å 97.19Å 156.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.55 – 1.80 82.55 – 1.80	Depositor EDS
% Data completeness (in resolution range)	91.8 (82.55-1.80) 91.8 (82.55-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.162 , 0.177 0.162 , 0.177	Depositor DCC
R_{free} test set	4320 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	16.1	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7095	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO, GLC, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	1/6331 (0.0%)	0.80	3/8625 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	767	GLN	CG-CD	-14.07	1.16	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	767	GLN	CG-CD-NE2	13.00	135.89	116.40
1	A	767	GLN	CG-CD-OE1	-10.14	100.52	120.80
1	A	767	GLN	CB-CG-CD	6.53	123.70	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	98	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6142	0	5646	32	0
2	B	34	0	30	0	0
3	C	23	0	21	0	0
4	D	111	0	93	0	0
5	A	60	0	0	0	0
6	A	68	0	102	3	0
7	A	657	0	0	0	0
All	All	7095	0	5892	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28[B]:HIS:ND1	1:A:40[B]:GLU:HG3	1.82	0.95
1:A:103:ARG:HE	6:A:835:EDO:H11	1.56	0.71
1:A:324:LEU:HD11	1:A:338:ILE:HD12	1.81	0.63
1:A:324:LEU:HD11	1:A:338:ILE:CD1	2.31	0.60
1:A:108:HIS:HD2	1:A:136:THR:OG1	1.87	0.57
1:A:681:LEU:O	1:A:685:GLU:HG2	2.05	0.57
1:A:96:HIS:HD2	1:A:309:TYR:OH	1.92	0.53
1:A:213:SER:OG	1:A:219:HIS:HD2	1.93	0.52
1:A:171:GLN:HB2	1:A:188:GLU:HB3	1.93	0.51
1:A:452:LEU:HD22	1:A:461:ALA:HA	1.93	0.50
1:A:549:HIS:CE1	1:A:553:ASN:HD21	2.32	0.47
1:A:313:TYR:CD1	1:A:338:ILE:HG12	2.50	0.47
1:A:102:HIS:HE1	1:A:176:TYR:OH	1.97	0.46
1:A:105:GLU:OE2	1:A:108:HIS:HE1	1.98	0.46
1:A:28[B]:HIS:CG	1:A:40[B]:GLU:HG3	2.49	0.46
1:A:80:TRP:CZ2	1:A:177:GLY:HA2	2.51	0.46
1:A:410:GLU:HB2	6:A:836:EDO:H22	1.99	0.45
1:A:107:LEU:HD22	1:A:174:ILE:HG21	2.00	0.44
1:A:605:ASN:HD21	1:A:610:MET:HE2	1.83	0.44
1:A:516:TYR:HB3	6:A:842:EDO:H22	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:SER:H	1:A:627:ASN:ND2	2.16	0.43
1:A:179:GLY:HA3	1:A:523:TYR:CE2	2.53	0.43
1:A:101:ARG:HG2	1:A:108:HIS:HB2	2.00	0.43
1:A:468:HIS:CE1	1:A:519:MET:HE2	2.53	0.43
1:A:682[B]:GLN:O	1:A:682[B]:GLN:CD	2.62	0.42
1:A:384:SER:OG	1:A:455:HIS:HD2	2.02	0.42
1:A:45:ASP:O	1:A:455:HIS:HE1	2.04	0.41
1:A:468:HIS:HB3	1:A:519:MET:HG3	2.03	0.41
1:A:468:HIS:ND1	1:A:519:MET:HE2	2.34	0.41
1:A:148:PHE:HA	1:A:167:ASN:HB2	2.03	0.41
1:A:476:LYS:HB2	1:A:553:ASN:ND2	2.36	0.40
1:A:540:ASN:HB3	1:A:606:TRP:CE3	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	765/783 (98%)	745 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	634/648 (98%)	632 (100%)	2 (0%)	86 86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	364	TYR
1	A	468	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	96	HIS
1	A	102	HIS
1	A	108	HIS
1	A	150	GLN
1	A	171	GLN
1	A	219	HIS
1	A	326	GLN
1	A	327	ASN
1	A	455	HIS
1	A	549	HIS
1	A	553	ASN
1	A	605	ASN
1	A	627	ASN
1	A	666	ASN
1	A	713	HIS
1	A	759	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2	12,12,12	0.89	1 (8%)	17,17,17	1.58	3 (17%)
2	BGC	B	2	2	11,11,12	1.13	1 (9%)	15,15,17	1.17	2 (13%)
2	BGC	B	3	2	11,11,12	0.40	0	15,15,17	1.03	1 (6%)
3	BGC	C	1	3	12,12,12	1.28	1 (8%)	17,17,17	0.86	0
3	BGC	C	2	3	11,11,12	0.39	0	15,15,17	0.83	0
4	BGC	D	1	4	12,12,12	0.70	0	17,17,17	1.00	2 (11%)
4	BGC	D	10	4	11,11,12	0.57	0	15,15,17	1.88	1 (6%)
4	BGC	D	2	4	11,11,12	0.54	0	15,15,17	0.98	1 (6%)
4	BGC	D	3	4	11,11,12	0.53	0	15,15,17	1.40	2 (13%)
4	BGC	D	4	4	11,11,12	0.43	0	15,15,17	0.85	1 (6%)
4	BGC	D	5	4	11,11,12	0.40	0	15,15,17	1.35	1 (6%)
4	BGC	D	6	4	11,11,12	0.44	0	15,15,17	1.48	1 (6%)
4	BGC	D	7	4	11,11,12	0.60	0	15,15,17	0.96	1 (6%)
4	BGC	D	8	4	11,11,12	0.56	0	15,15,17	1.03	1 (6%)
4	BGC	D	9	4	11,11,12	0.60	0	15,15,17	1.27	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1
2	BGC	B	3	2	-	0/2/19/22	0/1/1/1
3	BGC	C	1	3	-	0/2/22/22	0/1/1/1
3	BGC	C	2	3	-	0/2/19/22	0/1/1/1
4	BGC	D	1	4	-	2/2/22/22	0/1/1/1
4	BGC	D	10	4	-	1/2/19/22	0/1/1/1
4	BGC	D	2	4	-	0/2/19/22	0/1/1/1
4	BGC	D	3	4	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BGC	D	4	4	-	0/2/19/22	0/1/1/1
4	BGC	D	5	4	-	0/2/19/22	0/1/1/1
4	BGC	D	6	4	-	0/2/19/22	0/1/1/1
4	BGC	D	7	4	-	0/2/19/22	0/1/1/1
4	BGC	D	8	4	-	1/2/19/22	0/1/1/1
4	BGC	D	9	4	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	BGC	O3-C3	-3.93	1.33	1.43
2	B	2	BGC	O3-C3	-3.42	1.34	1.43
2	B	1	GLC	O1-C1	2.01	1.45	1.39

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	10	BGC	C1-O5-C5	6.75	121.23	112.19
4	D	6	BGC	C1-O5-C5	5.28	119.26	112.19
4	D	5	BGC	C1-O5-C5	4.82	118.65	112.19
4	D	9	BGC	C1-C2-C3	3.74	115.09	109.64
2	B	1	GLC	C1-O5-C5	3.51	120.45	113.65
2	B	3	BGC	C1-O5-C5	3.51	116.88	112.19
2	B	2	BGC	O3-C3-C2	-3.23	103.47	110.05
4	D	3	BGC	C1-O5-C5	3.22	116.50	112.19
4	D	3	BGC	O3-C3-C2	-3.20	103.52	110.05
2	B	1	GLC	C3-C4-C5	2.81	115.33	110.23
2	B	2	BGC	C1-O5-C5	2.61	115.69	112.19
4	D	7	BGC	C1-O5-C5	2.48	115.52	112.19
4	D	8	BGC	C3-C4-C5	2.39	114.56	110.23
4	D	2	BGC	C1-C2-C3	2.21	112.86	109.64
4	D	1	BGC	C3-C4-C5	2.20	114.22	110.23
4	D	1	BGC	C1-O5-C5	2.09	117.70	113.65
2	B	1	GLC	O1-C1-C2	2.07	114.99	108.98
4	D	4	BGC	C1-O5-C5	2.03	114.91	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1	BGC	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	D	10	BGC	O5-C5-C6-O6
4	D	8	BGC	C4-C5-C6-O6
4	D	1	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	A	842	-	3,3,3	0.47	0	2,2,2	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	836	-	3,3,3	0.35	0	2,2,2	0.41	0
6	EDO	A	838	-	3,3,3	0.38	0	2,2,2	0.57	0
6	EDO	A	829	-	3,3,3	0.44	0	2,2,2	0.62	0
5	PO4	A	824	-	4,4,4	0.91	0	6,6,6	0.48	0
5	PO4	A	817	-	4,4,4	1.06	0	6,6,6	0.72	0
5	PO4	A	820	-	4,4,4	0.99	0	6,6,6	0.50	0
6	EDO	A	831	-	3,3,3	0.43	0	2,2,2	0.30	0
5	PO4	A	827	-	4,4,4	0.93	0	6,6,6	0.52	0
6	EDO	A	841	-	3,3,3	0.41	0	2,2,2	0.19	0
6	EDO	A	834	-	3,3,3	0.40	0	2,2,2	0.56	0
6	EDO	A	828	-	3,3,3	0.45	0	2,2,2	0.44	0
6	EDO	A	840	-	3,3,3	0.40	0	2,2,2	0.43	0
5	PO4	A	823	-	4,4,4	0.91	0	6,6,6	0.60	0
5	PO4	A	816	-	4,4,4	0.94	0	6,6,6	0.59	0
6	EDO	A	844	-	3,3,3	0.40	0	2,2,2	0.47	0
6	EDO	A	837	-	3,3,3	0.42	0	2,2,2	0.41	0
5	PO4	A	818	-	4,4,4	1.07	0	6,6,6	0.49	0
5	PO4	A	826	-	4,4,4	0.98	0	6,6,6	0.46	0
6	EDO	A	833	-	3,3,3	0.36	0	2,2,2	0.57	0
5	PO4	A	825	-	4,4,4	0.91	0	6,6,6	0.57	0
5	PO4	A	821	-	4,4,4	0.94	0	6,6,6	0.54	0
6	EDO	A	835	-	3,3,3	0.41	0	2,2,2	0.33	0
5	PO4	A	819	-	4,4,4	1.02	0	6,6,6	0.52	0
6	EDO	A	843	-	3,3,3	0.42	0	2,2,2	0.43	0
6	EDO	A	830	-	3,3,3	0.41	0	2,2,2	0.44	0
6	EDO	A	839	-	3,3,3	0.39	0	2,2,2	0.46	0
5	PO4	A	822	-	4,4,4	0.89	0	6,6,6	0.46	0
6	EDO	A	832	-	3,3,3	0.50	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	841	-	-	1/1/1/1	-
6	EDO	A	844	-	-	0/1/1/1	-
6	EDO	A	834	-	-	0/1/1/1	-
6	EDO	A	842	-	-	1/1/1/1	-
6	EDO	A	837	-	-	0/1/1/1	-
6	EDO	A	833	-	-	0/1/1/1	-
6	EDO	A	843	-	-	0/1/1/1	-
6	EDO	A	836	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	830	-	-	0/1/1/1	-
6	EDO	A	828	-	-	0/1/1/1	-
6	EDO	A	838	-	-	1/1/1/1	-
6	EDO	A	839	-	-	0/1/1/1	-
6	EDO	A	829	-	-	0/1/1/1	-
6	EDO	A	840	-	-	0/1/1/1	-
6	EDO	A	832	-	-	1/1/1/1	-
6	EDO	A	831	-	-	0/1/1/1	-
6	EDO	A	835	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	842	EDO	O1-C1-C2-O2
6	A	835	EDO	O1-C1-C2-O2
6	A	838	EDO	O1-C1-C2-O2
6	A	841	EDO	O1-C1-C2-O2
6	A	832	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	842	EDO	1	0
6	A	836	EDO	1	0
6	A	835	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	752/783 (96%)	-0.36	11 (1%) 72 72	6, 15, 26, 47	18 (2%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	MET	3.8
1	A	778	GLY	3.3
1	A	775	ILE	3.1
1	A	691	TYR	3.1
1	A	28[A]	HIS	2.9
1	A	49	ILE	2.7
1	A	777	ASN	2.2
1	A	48	GLY	2.2
1	A	697	GLY	2.1
1	A	692	GLY	2.0
1	A	766	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BGC	D	8	11/12	0.59	0.18	61,66,67,67	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BGC	D	10	11/12	0.63	0.21	74,77,77,78	0
4	BGC	D	1	12/12	0.66	0.17	41,52,55,58	0
4	BGC	D	7	11/12	0.72	0.16	44,48,50,56	0
4	BGC	D	9	11/12	0.75	0.20	62,67,71,73	0
4	BGC	D	6	11/12	0.83	0.12	27,30,32,37	0
4	BGC	D	2	11/12	0.86	0.12	23,27,32,34	0
2	BGC	B	3	11/12	0.88	0.11	21,23,25,26	0
4	BGC	D	3	11/12	0.95	0.07	15,17,20,20	0
2	GLC	B	1	12/12	0.95	0.07	14,14,15,15	0
3	BGC	C	1	12/12	0.95	0.07	14,19,21,21	0
4	BGC	D	5	11/12	0.96	0.06	15,18,20,23	0
3	BGC	C	2	11/12	0.97	0.05	12,12,13,14	0
4	BGC	D	4	11/12	0.97	0.05	15,15,16,17	0
2	BGC	B	2	11/12	0.98	0.04	12,13,14,18	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	A	843	4/4	0.68	0.26	56,56,57,58	0
6	EDO	A	835	4/4	0.74	0.21	39,41,41,44	0
5	PO4	A	826	5/5	0.74	0.19	64,65,67,67	0
5	PO4	A	827	5/5	0.79	0.18	75,75,76,77	0
6	EDO	A	834	4/4	0.79	0.20	42,43,43,45	0
6	EDO	A	833	4/4	0.80	0.17	33,34,34,35	0
6	EDO	A	838	4/4	0.84	0.16	34,34,35,35	0
6	EDO	A	842	4/4	0.85	0.21	37,38,38,38	0
6	EDO	A	836	4/4	0.85	0.18	36,37,37,37	0
5	PO4	A	824	5/5	0.86	0.11	48,49,50,51	0
6	EDO	A	841	4/4	0.86	0.14	35,35,36,36	0
6	EDO	A	839	4/4	0.87	0.17	44,44,44,45	0
6	EDO	A	830	4/4	0.87	0.15	41,42,42,44	0
6	EDO	A	844	4/4	0.87	0.18	49,51,52,54	0
6	EDO	A	831	4/4	0.88	0.14	39,40,41,44	0
5	PO4	A	823	5/5	0.89	0.12	44,46,48,48	0
5	PO4	A	821	5/5	0.89	0.11	45,46,47,48	0
5	PO4	A	825	5/5	0.91	0.11	38,40,41,41	0
6	EDO	A	829	4/4	0.91	0.10	17,17,18,19	0
6	EDO	A	837	4/4	0.91	0.13	42,42,43,45	0
6	EDO	A	832	4/4	0.92	0.12	29,29,30,30	0
5	PO4	A	820	5/5	0.92	0.09	45,45,46,46	0
6	EDO	A	840	4/4	0.93	0.12	30,30,31,32	0
5	PO4	A	822	5/5	0.94	0.15	35,36,37,37	0
6	EDO	A	828	4/4	0.96	0.07	19,20,20,20	0
5	PO4	A	818	5/5	0.97	0.06	25,25,26,26	0
5	PO4	A	816	5/5	0.98	0.07	27,27,28,28	0
5	PO4	A	819	5/5	0.98	0.06	23,23,23,24	0
5	PO4	A	817	5/5	0.98	0.05	18,18,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.