



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 08:22 PM UTC

PDB ID : 7T4P / pdb_00007t4p
EMDB ID : EMD-25684
Title : CryoEM structure of Methylococcus capsulatus (Bath) pMMO treated with potassium cyanide and copper in a native lipid nanodisc at 3.62 Angstrom resolution
Authors : Koo, C.W.; Rosenzweig, A.C.
Deposited on : 2021-12-10
Resolution : 3.62 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11773 (3.12 - 4.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	
1	E	414	
1	I	414	
2	C	260	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	260	
2	K	260	
3	B	247	
3	F	247	
3	J	247	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	P1O	C	308	-	-	X	-
8	P1O	G	309	-	-	X	-
8	P1O	K	308	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 24348 atoms, of which 2106 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Particulate methane monooxygenase alpha subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	382	Total	C	N	O	S	0	0
			3017	1938	513	551	15		
1	E	382	Total	C	N	O	S	0	0
			3017	1938	513	551	15		
1	I	382	Total	C	N	O	S	0	0
			3017	1938	513	551	15		

- Molecule 2 is a protein called Ammonia monooxygenase/methane monooxygenase, subunit C family protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	236	Total	C	N	O	S	0	0
			1972	1339	299	329	5		
2	G	236	Total	C	N	O	S	0	0
			1972	1339	299	329	5		
2	K	236	Total	C	N	O	S	0	0
			1972	1339	299	329	5		


- Molecule 3 is a protein called Particulate methane monooxygenase beta subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	B	241	Total	C	N	O	S	0	0
			1976	1329	315	321	11		
3	F	241	Total	C	N	O	S	0	0
			1976	1329	315	321	11		
3	J	241	Total	C	N	O	S	0	0
			1976	1329	315	321	11		

- Molecule 4 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
4	A	2	Total	Cu	0
			2	2	
4	C	1	Total	Cu	0
			1	1	
4	G	1	Total	Cu	0
			1	1	
4	K	1	Total	Cu	0
			1	1	
4	E	2	Total	Cu	0
			2	2	
4	I	2	Total	Cu	0
			2	2	

- Molecule 5 is DECANE (CCD ID: D10) (formula: $C_{10}H_{22}$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
5	A	1	Total	C	H	0
			32	10	22	
5	C	1	Total	C	H	0
			32	10	22	
5	B	1	Total	C	H	0
			32	10	22	
5	B	1	Total	C	H	0
			32	10	22	
5	B	1	Total	C	H	0
			32	10	22	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
5	G	1	Total	C	H	0
			32	10	22	
5	K	1	Total	C	H	0
			32	10	22	
5	F	1	Total	C	H	0
			32	10	22	
5	F	1	Total	C	H	0
			32	10	22	
5	F	1	Total	C	H	0
			32	10	22	
5	J	1	Total	C	H	0
			32	10	22	
5	J	1	Total	C	H	0
			32	10	22	
5	J	1	Total	C	H	0
			32	10	22	
5	E	1	Total	C	H	0
			32	10	22	
5	I	1	Total	C	H	0
			32	10	22	

- Molecule 6 is DIUNDECYL PHOSPHATIDYL CHOLINE (CCD ID: PLC) (formula: C₃₂H₆₅NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
6	C	1	Total	C	H	N	O	P	0
			106	32	64	1	8	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						AltConf
6	C	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	C	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	C	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	B	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	B	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	B	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	G	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	G	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	G	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	G	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	K	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	K	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	K	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	F	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	F	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	J	1	Total 106	C 32	H 64	N 1	O 8	P 1	0
6	J	1	Total 106	C 32	H 64	N 1	O 8	P 1	0

- Molecule 7 is 1,2-dihexanoyl-sn-glycero-3-phosphocholine (CCD ID: HXG) (formula: C₂₀H₄₁NO₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
7	C	1	Total	C	H	N	O	P	0
			70	20	40	1	8	1	
7	G	1	Total	C	H	N	O	P	0
			70	20	40	1	8	1	
7	K	1	Total	C	H	N	O	P	0
			70	20	40	1	8	1	

- Molecule 8 is 1,2-DIDECANOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: P10) (formula: C₂₈H₅₇NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
8	C	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	C	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	B	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	G	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	G	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	K	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	K	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	F	1	Total 94	C 28	H 56	N 1	O 8	P 1	0
8	J	1	Total 94	C 28	H 56	N 1	O 8	P 1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Particulate methane monooxygenase alpha subunit

Chain A:




- Molecule 1: Particulate methane monooxygenase alpha subunit

Chain E:




- Molecule 1: Particulate methane monooxygenase alpha subunit

Chain I: 




• Molecule 2: Ammonia monooxygenase/methane monooxygenase, subunit C family protein

Chain C: 




• Molecule 2: Ammonia monooxygenase/methane monooxygenase, subunit C family protein


Chain G: 




• Molecule 2: Ammonia monooxygenase/methane monooxygenase, subunit C family protein



• Molecule 3: Particulate methane monooxygenase beta subunit




• Molecule 3: Particulate methane monooxygenase beta subunit



• Molecule 3: Particulate methane monooxygenase beta subunit





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	812752	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1.2	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.174	Depositor
Minimum map value	-0.119	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.0272	Depositor
Map size (Å)	204.0, 204.0, 204.0	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.4722222, 0.4722222, 0.4722222	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P1O, D10, PLC, HXG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	1/3099 (0.0%)	0.42	0/4215
1	E	0.27	1/3099 (0.0%)	0.42	0/4215
1	I	0.27	1/3099 (0.0%)	0.42	0/4215
2	C	0.30	1/2051 (0.0%)	0.57	2/2810 (0.1%)
2	G	0.30	1/2051 (0.0%)	0.57	2/2810 (0.1%)
2	K	0.30	1/2051 (0.0%)	0.57	2/2810 (0.1%)
3	B	0.30	0/2052	0.45	0/2808
3	F	0.30	0/2052	0.45	0/2808
3	J	0.30	0/2052	0.45	0/2808
All	All	0.29	6/21606 (0.0%)	0.48	6/29499 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	33	HIS	CE1-NE2	-6.26	1.26	1.32
1	A	33	HIS	CE1-NE2	-6.23	1.26	1.32
1	E	33	HIS	CE1-NE2	-6.21	1.26	1.32
2	K	233	PHE	N-CA	5.20	1.49	1.46
2	G	233	PHE	N-CA	5.19	1.49	1.46

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	233	PHE	CA-C-O	-6.23	114.33	117.94
2	K	233	PHE	CA-C-O	-6.22	114.33	117.94
2	G	233	PHE	CA-C-O	-6.17	114.36	117.94
2	C	233	PHE	O-C-N	5.02	125.23	121.47
2	G	233	PHE	O-C-N	5.01	125.23	121.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3017	0	2980	85	0
1	E	3017	0	2980	84	0
1	I	3017	0	2980	86	0
2	C	1972	0	1904	141	0
2	G	1972	0	1904	138	0
2	K	1972	0	1904	138	0
3	B	1976	0	1936	113	0
3	F	1976	0	1936	106	0
3	J	1976	0	1936	115	0
4	A	2	0	0	0	0
4	C	1	0	0	0	0
4	E	2	0	0	0	0
4	G	1	0	0	0	0
4	I	2	0	0	0	0
4	K	1	0	0	0	0
5	A	10	22	22	0	0
5	B	30	66	66	8	0
5	C	10	22	22	1	0
5	E	10	22	22	0	0
5	F	30	66	66	8	0
5	G	10	22	22	0	0
5	I	10	22	22	0	0
5	J	30	66	66	8	0
5	K	10	22	22	1	0
6	B	126	192	192	15	0
6	C	168	256	256	14	0
6	F	84	128	128	3	0
6	G	168	256	256	17	0
6	J	84	128	128	3	0
6	K	126	192	192	4	0
7	C	30	40	40	14	0
7	G	30	40	40	14	0
7	K	30	40	40	14	0
8	B	38	56	56	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	76	112	112	31	0
8	F	38	56	56	11	0
8	G	76	112	112	31	0
8	J	38	56	56	8	0
8	K	76	112	112	31	0
All	All	22242	2106	22566	863	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 863 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:K:112:TRP:CH2	8:K:308:P1O:H2	1.86	1.11
2:G:112:TRP:CH2	8:G:309:P1O:H2	1.86	1.10
2:K:67:TRP:HD1	7:K:305:HXG:H41	1.15	1.10
2:C:112:TRP:CH2	8:C:308:P1O:H2	1.86	1.09
2:C:67:TRP:HD1	7:C:305:HXG:H41	1.15	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/414 (92%)	355 (93%)	25 (7%)	0	100	100
1	E	380/414 (92%)	355 (93%)	25 (7%)	0	100	100
1	I	380/414 (92%)	355 (93%)	25 (7%)	0	100	100
2	C	234/260 (90%)	215 (92%)	17 (7%)	2 (1%)	14	44
2	G	234/260 (90%)	214 (92%)	18 (8%)	2 (1%)	14	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	K	234/260 (90%)	215 (92%)	17 (7%)	2 (1%)	14	44
3	B	239/247 (97%)	228 (95%)	10 (4%)	1 (0%)	30	59
3	F	239/247 (97%)	228 (95%)	10 (4%)	1 (0%)	30	59
3	J	239/247 (97%)	228 (95%)	10 (4%)	1 (0%)	30	59
All	All	2559/2763 (93%)	2393 (94%)	157 (6%)	9 (0%)	31	59

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	277	SER
3	B	246	SER
2	G	277	SER
2	K	277	SER
3	F	246	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/345 (94%)	322 (100%)	1 (0%)	86	81
1	E	323/345 (94%)	322 (100%)	1 (0%)	86	81
1	I	323/345 (94%)	322 (100%)	1 (0%)	86	81
2	C	200/212 (94%)	198 (99%)	2 (1%)	68	72
2	G	200/212 (94%)	198 (99%)	2 (1%)	68	72
2	K	200/212 (94%)	198 (99%)	2 (1%)	68	72
3	B	206/210 (98%)	203 (98%)	3 (2%)	57	66
3	F	206/210 (98%)	203 (98%)	3 (2%)	57	66
3	J	206/210 (98%)	203 (98%)	3 (2%)	57	66
All	All	2187/2301 (95%)	2169 (99%)	18 (1%)	70	74

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	244	LEU
1	I	414	MET
1	E	414	MET
2	K	90	MET
3	J	241	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	240	ASN
3	F	245	GLN
1	E	182	ASN
3	J	240	ASN
3	J	245	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 54 ligands modelled in this entry, 9 are monoatomic - leaving 45 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PLC	C	309	-	41,41,41	0.81	0	47,49,49	0.59	0
6	PLC	J	302	-	41,41,41	0.87	0	47,49,49	0.75	1 (2%)
6	PLC	B	301	-	41,41,41	0.85	0	47,49,49	0.81	1 (2%)
7	HXG	K	305	-	29,29,29	0.35	0	35,37,37	0.36	0
7	HXG	C	305	-	29,29,29	0.35	0	35,37,37	0.37	0
5	D10	B	303	-	9,9,9	0.21	0	8,8,8	0.55	0
6	PLC	K	304	-	41,41,41	0.89	0	47,49,49	0.70	1 (2%)
5	D10	J	304	-	9,9,9	0.20	0	8,8,8	0.57	0
8	P1O	G	309	-	37,37,37	1.12	2 (5%)	43,45,45	1.14	3 (6%)
8	P1O	K	308	-	37,37,37	1.12	2 (5%)	43,45,45	1.13	3 (6%)
5	D10	B	302	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	G	305	-	41,41,41	0.89	0	47,49,49	0.70	1 (2%)
8	P1O	C	308	-	37,37,37	1.12	2 (5%)	43,45,45	1.13	3 (6%)
6	PLC	C	302	-	41,41,41	1.06	2 (4%)	47,49,49	1.10	3 (6%)
6	PLC	C	306	-	41,41,41	0.91	0	47,49,49	0.79	2 (4%)
8	P1O	K	307	-	37,37,37	0.87	0	43,45,45	0.70	2 (4%)
5	D10	F	304	-	9,9,9	0.21	0	8,8,8	0.55	0
5	D10	C	303	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	F	302	-	41,41,41	0.85	0	47,49,49	0.81	1 (2%)
6	PLC	K	301	-	41,41,41	0.81	0	47,49,49	0.59	0
5	D10	F	305	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	G	303	-	41,41,41	1.06	2 (4%)	47,49,49	1.10	3 (6%)
8	P1O	G	308	-	37,37,37	0.87	0	43,45,45	0.70	2 (4%)
8	P1O	F	306	-	37,37,37	1.12	2 (5%)	43,45,45	1.10	3 (6%)
5	D10	A	503	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	C	304	-	41,41,41	0.89	0	47,49,49	0.70	1 (2%)
5	D10	K	303	-	9,9,9	0.21	0	8,8,8	0.57	0
8	P1O	B	307	-	37,37,37	1.12	2 (5%)	43,45,45	1.11	3 (6%)
6	PLC	B	306	-	41,41,41	0.87	0	47,49,49	0.75	1 (2%)
6	PLC	K	306	-	41,41,41	0.91	0	47,49,49	0.79	2 (4%)
8	P1O	C	307	-	37,37,37	0.87	0	43,45,45	0.70	2 (4%)
5	D10	G	304	-	9,9,9	0.20	0	8,8,8	0.56	0
5	D10	J	306	-	9,9,9	0.20	0	8,8,8	0.56	0
8	P1O	J	301	-	37,37,37	1.12	2 (5%)	43,45,45	1.10	3 (6%)
5	D10	B	304	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	G	307	-	41,41,41	0.90	0	47,49,49	0.79	2 (4%)
6	PLC	B	305	-	41,41,41	1.06	2 (4%)	47,49,49	1.10	3 (6%)
6	PLC	G	301	-	41,41,41	0.81	0	47,49,49	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	HXG	G	306	-	29,29,29	0.35	0	35,37,37	0.36	0
5	D10	J	305	-	9,9,9	0.21	0	8,8,8	0.56	0
5	D10	E	501	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	F	301	-	41,41,41	0.87	0	47,49,49	0.75	1 (2%)
5	D10	F	303	-	9,9,9	0.20	0	8,8,8	0.56	0
6	PLC	J	303	-	41,41,41	0.85	0	47,49,49	0.81	1 (2%)
5	D10	I	501	-	9,9,9	0.21	0	8,8,8	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PLC	C	309	-	-	12/45/45/45	-
6	PLC	J	302	-	-	11/45/45/45	-
6	PLC	B	301	-	-	13/45/45/45	-
7	HXG	K	305	-	-	7/33/33/33	-
7	HXG	C	305	-	-	7/33/33/33	-
5	D10	B	303	-	-	6/7/7/7	-
6	PLC	K	304	-	-	13/45/45/45	-
5	D10	J	304	-	-	0/7/7/7	-
8	P1O	G	309	-	-	26/41/41/41	-
8	P1O	K	308	-	-	26/41/41/41	-
5	D10	B	302	-	-	0/7/7/7	-
6	PLC	G	305	-	-	13/45/45/45	-
8	P1O	C	308	-	-	26/41/41/41	-
6	PLC	C	302	-	-	22/45/45/45	-
6	PLC	C	306	-	-	15/45/45/45	-
8	P1O	K	307	-	-	15/41/41/41	-
5	D10	F	304	-	-	6/7/7/7	-
5	D10	C	303	-	-	0/7/7/7	-
6	PLC	F	302	-	-	13/45/45/45	-
6	PLC	K	301	-	-	12/45/45/45	-
5	D10	F	305	-	-	5/7/7/7	-
6	PLC	G	303	-	-	22/45/45/45	-
8	P1O	G	308	-	-	15/41/41/41	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	P1O	F	306	-	-	20/41/41/41	-
5	D10	A	503	-	-	0/7/7/7	-
6	PLC	C	304	-	-	13/45/45/45	-
5	D10	K	303	-	-	0/7/7/7	-
8	P1O	B	307	-	-	20/41/41/41	-
6	PLC	B	306	-	-	11/45/45/45	-
6	PLC	K	306	-	-	15/45/45/45	-
8	P1O	C	307	-	-	15/41/41/41	-
5	D10	G	304	-	-	0/7/7/7	-
5	D10	J	306	-	-	5/7/7/7	-
8	P1O	J	301	-	-	20/41/41/41	-
5	D10	B	304	-	-	5/7/7/7	-
6	PLC	G	307	-	-	15/45/45/45	-
6	PLC	B	305	-	-	22/45/45/45	-
6	PLC	G	301	-	-	12/45/45/45	-
7	HXG	G	306	-	-	7/33/33/33	-
5	D10	J	305	-	-	6/7/7/7	-
5	D10	E	501	-	-	0/7/7/7	-
6	PLC	F	301	-	-	11/45/45/45	-
5	D10	F	303	-	-	0/7/7/7	-
6	PLC	J	303	-	-	13/45/45/45	-
5	D10	I	501	-	-	0/7/7/7	-

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	F	306	P1O	O5-C9	4.29	1.45	1.33
8	B	307	P1O	O5-C9	4.28	1.45	1.33
6	G	303	PLC	O3-CB	4.28	1.45	1.33
8	J	301	P1O	O5-C9	4.27	1.45	1.33
6	C	302	PLC	O3-CB	4.25	1.45	1.33

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	307	P1O	O7-C19-C20	4.24	120.66	111.48
8	J	301	P1O	O7-C19-C20	4.22	120.62	111.48
8	F	306	P1O	O7-C19-C20	4.21	120.59	111.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	G	309	P1O	O7-C19-C20	4.19	120.54	111.48
8	C	308	P1O	O7-C19-C20	4.17	120.49	111.48

There are no chirality outliers.

5 of 495 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	302	PLC	O4P-C4-C5-N
6	C	302	PLC	C1'-C'-O2-C2
6	C	302	PLC	C1B-CB-O3-C3
6	C	302	PLC	OB-CB-O3-C3
6	C	302	PLC	C1-O3P-P-O2P

There are no ring outliers.

29 monomers are involved in 240 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	309	PLC	2	0
6	B	301	PLC	3	0
7	K	305	HXG	14	0
7	C	305	HXG	14	0
6	K	304	PLC	2	0
5	J	304	D10	6	0
8	G	309	P1O	31	0
8	K	308	P1O	31	0
5	B	302	D10	6	0
6	G	305	PLC	2	0
8	C	308	P1O	31	0
6	C	302	PLC	10	0
5	C	303	D10	1	0
6	F	302	PLC	3	0
6	K	301	PLC	2	0
5	F	305	D10	2	0
6	G	303	PLC	14	0
8	F	306	P1O	11	0
6	C	304	PLC	2	0
5	K	303	D10	1	0
8	B	307	P1O	8	0
5	J	306	D10	2	0
8	J	301	P1O	8	0
5	B	304	D10	2	0


Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	305	PLC	12	0
6	G	301	PLC	1	0
7	G	306	HXG	14	0
5	F	303	D10	6	0
6	J	303	PLC	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























































































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-25684. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 216



Y Index: 216




Z Index: 216


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 179



Y Index: 252




Z Index: 168


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 98 nm³; this corresponds to an approximate mass of 88 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.276 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.276 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.62	-	-
Author-provided FSC curve	3.38	3.71	3.41
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-25684 and PDB model 7T4P. Per-residue inclusion information can be found in section [3](#) on page [10](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0272 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0272).

9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 70% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0272) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7030	0.5090
A	0.7660	0.5330
B	0.7490	0.5230
C	0.5780	0.4630
E	0.7630	0.5310
F	0.7600	0.5270
G	0.5710	0.4610
I	0.7620	0.5310
J	0.7560	0.5260
K	0.5750	0.4630

1.0

0.0

<0.0