



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:57 AM UTC

PDB ID : 3T5I / pdb_00003t5i
Title : Structure of Fully modified farnesylated Rheb Peptide in complex with PDE6D
Authors : Ismail, S.A.; Chen, Y.-X.; Wittinghofer, A.
Deposited on : 2011-07-27
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	84% 14% ..
1	B	152	82% 11% 7%
1	C	152	5% 84% 14% ..
1	D	152	86% 12% ..
2	Q	9	11% 22% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	9	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CMT	R	14	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	142	Total	C	N	O	S	0	2	0
			1168	753	202	208	5			
1	A	149	Total	C	N	O	S	0	5	0
			1232	796	207	223	6			
1	C	149	Total	C	N	O	S	0	3	0
			1222	785	210	221	6			
1	D	149	Total	C	N	O	S	0	0	0
			1205	774	204	221	6			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP O43924
B	0	SER	-	expression tag	UNP O43924
A	-1	GLY	-	expression tag	UNP O43924
A	0	SER	-	expression tag	UNP O43924
C	-1	GLY	-	expression tag	UNP O43924
C	0	SER	-	expression tag	UNP O43924
D	-1	GLY	-	expression tag	UNP O43924
D	0	SER	-	expression tag	UNP O43924

- Molecule 2 is a protein called C-terminal Farnesylated Rheb peptide CSQQGKSS(CMT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Q	3	Total	C	N	O	S	0	0	0
			20	10	3	6	1			
2	R	3	Total	C	N	O	S	0	0	0
			20	10	3	6	1			

- Molecule 3 is FARNESYL (CCD ID: FAR) (formula: C₁₅H₂₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C 15 15	0	0
3	A	1	Total C 15 15	0	0
3	C	1	Total C 15 15	0	0
3	Q	1	Total C 15 15	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	66	Total O 66 66	0	0
4	A	81	Total O 81 81	0	0
4	C	52	Total O 52 52	0	0
4	D	75	Total O 75 75	0	0
4	Q	1	Total O 1 1	0	0
4	R	3	Total O 3 3	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta





- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta




- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta



- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta



- Molecule 2: C-terminal Farnesylated Rheb peptide CSQQGKSS(CMT)



- Molecule 2: C-terminal Farnesylated Rheb peptide CSQQGKSS(CMT)



CYS	SER	GLN	GLY	LYS	S12	S13	C14
-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.75Å 70.05Å 71.23Å 90.00° 81.55° 90.00°	Depositor
Resolution (Å)	29.91 – 2.10 29.91 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.91-2.10) 99.6 (29.91-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.96 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.189 , 0.243 0.190 , 0.245	Depositor DCC
R_{free} test set	1870 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.155	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5205	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAR, CMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1275	0.83	0/1717
1	B	0.65	0/1198	0.78	0/1613
1	C	0.63	0/1257	0.84	0/1695
1	D	0.66	0/1231	0.84	0/1661
2	Q	1.00	0/11	0.79	0/13
2	R	0.02	0/11	0.03	0/13
All	All	0.64	0/4983	0.82	0/6712

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1232	0	1240	17	0
1	B	1168	0	1169	15	0
1	C	1222	0	1216	18	0
1	D	1205	0	1190	16	0
2	Q	20	0	15	1	0
2	R	20	0	16	4	0
3	A	15	0	26	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	15	0	26	5	0
3	C	15	0	26	2	0
3	Q	15	0	24	0	0
4	A	81	0	0	0	0
4	B	66	0	0	0	0
4	C	52	0	0	4	0
4	D	75	0	0	4	0
4	Q	1	0	0	0	0
4	R	3	0	0	0	0
All	All	5205	0	4948	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2010:FAR:H13	2:R:14:CMT:SG	1.71	1.24
3:A:2010:FAR:H11	2:R:14:CMT:SG	1.81	1.10
1:A:33[A]:GLN:HE22	1:D:30:ILE:H	1.00	0.99
1:B:20:MET:HE1	1:B:38:LEU:HD12	1.52	0.92
3:A:2010:FAR:H13	2:R:14:CMT:CB	2.05	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/152 (100%)	149 (98%)	3 (2%)	0	100	100
1	B	140/152 (92%)	138 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	150/152 (99%)	144 (96%)	6 (4%)	0	100	100
1	D	147/152 (97%)	145 (99%)	2 (1%)	0	100	100
2	Q	1/9 (11%)	1 (100%)	0	0	100	100
2	R	1/9 (11%)	1 (100%)	0	0	100	100
All	All	591/626 (94%)	578 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/136 (99%)	133 (98%)	2 (2%)	57	65
1	B	126/136 (93%)	122 (97%)	4 (3%)	34	38
1	C	132/136 (97%)	127 (96%)	5 (4%)	29	32
1	D	130/136 (96%)	125 (96%)	5 (4%)	29	32
2	Q	2/7 (29%)	2 (100%)	0	100	100
2	R	2/7 (29%)	1 (50%)	1 (50%)	0	0
All	All	527/558 (94%)	510 (97%)	17 (3%)	33	38

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	100	ASN
2	R	12	SER
1	C	29	LYS
1	C	83	LYS
1	C	85	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	126	ASN
1	D	33	GLN
1	D	116	GLN
1	D	45	HIS
1	A	45	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CMT	R	14	2,3	7,7,7	0.79	0	6,8,8	1.33	1 (16%)
2	CMT	Q	14	2,3	7,7,7	1.68	1 (14%)	6,8,8	2.99	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CMT	R	14	2,3	-	1/8/8/8	-
2	CMT	Q	14	2,3	-	4/8/8/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Q	14	CMT	OXT-C1	-3.63	1.37	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	14	CMT	C1-OXT-C	-7.04	99.94	115.92
2	R	14	CMT	C1-OXT-C	-2.84	109.47	115.92

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Q	14	CMT	CA-C-OXT-C1
2	Q	14	CMT	O-C-OXT-C1
2	Q	14	CMT	N-CA-CB-SG
2	R	14	CMT	N-CA-CB-SG
2	Q	14	CMT	O-C-CA-N

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	R	14	CMT	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAR	B	3010	-	14,14,14	0.67	0	15,16,16	1.04	0
3	FAR	C	4010	-	14,14,14	0.60	0	15,16,16	1.17	1 (6%)
3	FAR	Q	5010	2	14,14,14	0.45	0	15,16,16	1.74	5 (33%)
3	FAR	A	2010	2	14,14,14	0.41	0	15,16,16	1.58	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAR	B	3010	-	-	4/14/14/14	-
3	FAR	C	4010	-	-	2/14/14/14	-
3	FAR	Q	5010	2	-	2/14/14/14	-
3	FAR	A	2010	2	-	3/14/14/14	-

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Q	5010	FAR	C4-C3-C5	3.11	120.63	115.23
3	A	2010	FAR	C4-C3-C5	2.90	120.26	115.23
3	Q	5010	FAR	C6-C7-C8	-2.75	121.33	127.62
3	A	2010	FAR	C15-C13-C14	2.68	120.75	114.59
3	Q	5010	FAR	C6-C5-C3	-2.64	104.45	113.19

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	3010	FAR	C5-C6-C7-C8
3	B	3010	FAR	C12-C11-C9-C8
3	Q	5010	FAR	C10-C8-C9-C11
3	Q	5010	FAR	C7-C8-C9-C11
3	B	3010	FAR	C2-C3-C5-C6

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	3010	FAR	5	0
3	C	4010	FAR	2	0
3	A	2010	FAR	5	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	149/152 (98%)	-0.17	1 (0%) 84 86	12, 25, 35, 39	5 (3%)
1	B	142/152 (93%)	-0.03	2 (1%) 73 75	16, 26, 37, 53	2 (1%)
1	C	149/152 (98%)	0.22	7 (4%) 36 38	15, 30, 49, 64	3 (2%)
1	D	149/152 (98%)	0.03	2 (1%) 75 77	16, 29, 38, 43	0
2	Q	2/9 (22%)	1.12	0 100 100	38, 38, 38, 39	0
2	R	2/9 (22%)	0.87	0 100 100	33, 33, 33, 35	0
All	All	593/626 (94%)	0.02	12 (2%) 65 67	12, 27, 40, 64	10 (1%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	SER	3.1
1	C	85	GLN	3.0
1	C	115	SER	2.8
1	B	137	ASP	2.8
1	C	112	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CMT	R	14	8/8	0.91	0.10	20,31,32,32	0
2	CMT	Q	14	8/8	0.94	0.09	20,34,34,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FAR	C	4010	15/15	0.77	0.19	42,44,46,47	0
3	FAR	B	3010	15/15	0.80	0.15	41,43,48,48	0
3	FAR	Q	5010	15/15	0.87	0.11	26,32,34,35	0
3	FAR	A	2010	15/15	0.88	0.10	27,29,32,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.