



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 07:46 PM UTC

PDB ID : 6T5A / pdb_00006t5a
Title : Crystal structure of herpes simplex virus 1 pUL7:pUL51 complex
Authors : Butt, B.G.; Graham, S.C.
Deposited on : 2019-10-15
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	
1	B	136	
1	C	136	
1	D	136	
2	E	304	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	304	7% 88% 9%
2	G	304	5% 87% 9%
2	H	304	5% 86% 5%9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tegument protein UL51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	96	Total	C	N	O	S	0	1	0
			738	457	138	136	7			
1	B	96	Total	C	N	O	S	0	1	0
			738	457	138	136	7			
1	C	96	Total	C	N	O	S	0	2	0
			742	461	138	136	7			
1	D	96	Total	C	N	O	S	0	1	0
			738	457	138	136	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	initiating methionine	UNP D3YPL0
A	9	SER	CYS	engineered mutation	UNP D3YPL0
B	7	MET	-	initiating methionine	UNP D3YPL0
B	9	SER	CYS	engineered mutation	UNP D3YPL0
C	7	MET	-	initiating methionine	UNP D3YPL0
C	9	SER	CYS	engineered mutation	UNP D3YPL0
D	7	MET	-	initiating methionine	UNP D3YPL0
D	9	SER	CYS	engineered mutation	UNP D3YPL0

- Molecule 2 is a protein called Cytoplasmic envelopment protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	276	Total	C	N	O	S	0	7	0
			2242	1441	388	396	17			
2	F	276	Total	C	N	O	S	0	7	0
			2242	1441	388	396	17			
2	G	276	Total	C	N	O	S	0	7	0
			2242	1441	388	396	17			
2	H	276	Total	C	N	O	S	0	7	0
			2242	1441	388	396	17			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	297	GLY	-	expression tag	UNP A0A110B4Q7
E	298	SER	-	expression tag	UNP A0A110B4Q7
E	299	LEU	-	expression tag	UNP A0A110B4Q7
E	300	GLU	-	expression tag	UNP A0A110B4Q7
E	301	VAL	-	expression tag	UNP A0A110B4Q7
E	302	LEU	-	expression tag	UNP A0A110B4Q7
E	303	PHE	-	expression tag	UNP A0A110B4Q7
E	304	GLN	-	expression tag	UNP A0A110B4Q7
F	297	GLY	-	expression tag	UNP A0A110B4Q7
F	298	SER	-	expression tag	UNP A0A110B4Q7
F	299	LEU	-	expression tag	UNP A0A110B4Q7
F	300	GLU	-	expression tag	UNP A0A110B4Q7
F	301	VAL	-	expression tag	UNP A0A110B4Q7
F	302	LEU	-	expression tag	UNP A0A110B4Q7
F	303	PHE	-	expression tag	UNP A0A110B4Q7
F	304	GLN	-	expression tag	UNP A0A110B4Q7
G	297	GLY	-	expression tag	UNP A0A110B4Q7
G	298	SER	-	expression tag	UNP A0A110B4Q7
G	299	LEU	-	expression tag	UNP A0A110B4Q7
G	300	GLU	-	expression tag	UNP A0A110B4Q7
G	301	VAL	-	expression tag	UNP A0A110B4Q7
G	302	LEU	-	expression tag	UNP A0A110B4Q7
G	303	PHE	-	expression tag	UNP A0A110B4Q7
G	304	GLN	-	expression tag	UNP A0A110B4Q7
H	297	GLY	-	expression tag	UNP A0A110B4Q7
H	298	SER	-	expression tag	UNP A0A110B4Q7
H	299	LEU	-	expression tag	UNP A0A110B4Q7
H	300	GLU	-	expression tag	UNP A0A110B4Q7
H	301	VAL	-	expression tag	UNP A0A110B4Q7
H	302	LEU	-	expression tag	UNP A0A110B4Q7
H	303	PHE	-	expression tag	UNP A0A110B4Q7
H	304	GLN	-	expression tag	UNP A0A110B4Q7

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total	Cl	0	0
			1	1		
4	F	1	Total	Cl	0	0
			1	1		
4	G	1	Total	Cl	0	0
			1	1		
4	H	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	56	Total	O	0	0
			56	56		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	48	Total 48	O 48	0	0
5	C	49	Total 49	O 49	0	0
5	D	56	Total 56	O 56	0	0
5	E	126	Total 126	O 126	0	0
5	F	111	Total 111	O 111	0	0
5	G	139	Total 139	O 139	0	0
5	H	132	Total 132	O 132	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tegument protein UL51




- Molecule 1: Tegument protein UL51




- Molecule 1: Tegument protein UL51




- Molecule 1: Tegument protein UL51




- Molecule 2: Cytoplasmic envelopment protein 1










- Molecule 2: Cytoplasmic envelopment protein 1



- Molecule 2: Cytoplasmic envelopment protein 1



- Molecule 2: Cytoplasmic envelopment protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.51Å 106.30Å 106.01Å 90.00° 92.05° 90.00°	Depositor
Resolution (Å)	26.57 – 1.83 26.57 – 1.83	Depositor EDS
% Data completeness (in resolution range)	100.0 (26.57-1.83) 99.9 (26.57-1.83)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.83Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.194 , 0.220 0.199 , 0.227	Depositor DCC
R_{free} test set	8093 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.247 for -h,-l,-k 0.032 for -h,l,k 0.086 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12669	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/750	1.17	1/1017 (0.1%)
1	B	0.79	0/750	1.17	1/1017 (0.1%)
1	C	0.83	0/757	1.16	0/1027
1	D	0.80	0/750	1.16	0/1017
2	E	0.77	0/2306	1.12	3/3139 (0.1%)
2	F	0.76	0/2306	1.11	3/3139 (0.1%)
2	G	0.78	0/2306	1.09	2/3139 (0.1%)
2	H	0.78	0/2306	1.09	4/3139 (0.1%)
All	All	0.78	0/12231	1.12	14/16634 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	73	PHE	CA-CB-CG	6.17	119.97	113.80
2	E	179	GLU	CA-C-N	5.46	127.59	120.28
2	E	179	GLU	C-N-CA	5.46	127.59	120.28
2	F	179	GLU	CA-C-N	5.34	127.76	120.54
2	F	179	GLU	C-N-CA	5.34	127.76	120.54
2	H	73	PHE	CA-CB-CG	5.19	118.99	113.80
2	G	74	VAL	N-CA-C	-5.11	100.54	107.99
2	H	74	VAL	N-CA-C	-5.08	100.57	107.99
1	B	54	ASP	CA-CB-CG	5.08	117.68	112.60
1	A	54	ASP	CA-CB-CG	5.06	117.66	112.60
2	G	74	VAL	N-CA-CB	5.06	117.13	111.21
2	H	179	GLU	CA-C-N	5.04	127.04	120.28
2	H	179	GLU	C-N-CA	5.04	127.04	120.28
2	F	74	VAL	N-CA-C	-5.01	100.67	107.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	738	0	756	2	0
1	B	738	0	756	1	0
1	C	742	0	765	1	0
1	D	738	0	756	1	0
2	E	2242	0	2219	8	0
2	F	2242	0	2219	5	0
2	G	2242	0	2219	6	0
2	H	2242	0	2219	8	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
5	A	56	0	0	0	0
5	B	48	0	0	0	0
5	C	49	0	0	0	0
5	D	56	0	0	0	0
5	E	126	0	0	0	0
5	F	111	0	0	1	0
5	G	139	0	0	2	0
5	H	132	0	0	0	0
All	All	12669	0	11941	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:283[B]:THR:HB	2:F:284[B]:PRO:HD3	1.67	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:70:ARG:HH22	2:E:174:ARG:HH12	1.42	0.68
2:G:70:ARG:HH22	2:G:174:ARG:HH12	1.42	0.68
2:H:70:ARG:HH22	2:H:174:ARG:HH12	1.42	0.67
2:F:70:ARG:HH22	2:F:174:ARG:HH12	1.41	0.65
2:H:283[A]:THR:HB	2:H:284[A]:PRO:HD3	1.80	0.62
1:C:37:GLU:HG2	2:G:300:GLU:HG2	1.84	0.60
2:H:125:LEU:HD11	2:H:277:TYR:HD2	1.67	0.59
1:D:37:GLU:HG2	2:H:300:GLU:HG2	1.82	0.59
2:G:125:LEU:HD11	2:G:277:TYR:HD2	1.67	0.59
1:B:37:GLU:HG2	2:F:300:GLU:HG2	1.85	0.58
1:A:37:GLU:HG2	2:E:300:GLU:HG2	1.90	0.54
2:F:265:CYS:HB3	5:F:508:HOH:O	2.09	0.52
2:E:283[B]:THR:HB	2:E:284[B]:PRO:HD3	1.95	0.47
2:H:294:GLN:HG2	2:H:295:PHE:N	2.30	0.46
2:E:294:GLN:HG2	2:E:295:PHE:N	2.31	0.46
2:G:294:GLN:HG2	2:G:295:PHE:N	2.30	0.45
2:F:294:GLN:HG2	2:F:295:PHE:N	2.30	0.45
2:H:282[B]:GLU:H	2:H:282[B]:GLU:CD	2.25	0.44
2:E:125:LEU:HD11	2:E:277:TYR:HD2	1.83	0.43
1:A:65:LEU:HD13	2:E:212:PRO:HB2	2.00	0.42
2:G:62:ARG:HG3	5:G:503:HOH:O	2.19	0.41
2:E:16:LEU:HB2	2:H:302:LEU:HD21	2.02	0.41
2:E:302:LEU:HD21	2:H:16:LEU:HB2	2.03	0.40
2:G:265:CYS:HB3	5:G:527:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	93/136 (68%)	92 (99%)	1 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	93/136 (68%)	92 (99%)	1 (1%)	0	100	100
1	C	94/136 (69%)	93 (99%)	1 (1%)	0	100	100
1	D	93/136 (68%)	92 (99%)	1 (1%)	0	100	100
2	E	279/304 (92%)	272 (98%)	7 (2%)	0	100	100
2	F	279/304 (92%)	270 (97%)	9 (3%)	0	100	100
2	G	279/304 (92%)	274 (98%)	5 (2%)	0	100	100
2	H	279/304 (92%)	276 (99%)	3 (1%)	0	100	100
All	All	1489/1760 (85%)	1461 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/107 (73%)	78 (100%)	0	100	100
1	B	78/107 (73%)	78 (100%)	0	100	100
1	C	79/107 (74%)	79 (100%)	0	100	100
1	D	78/107 (73%)	78 (100%)	0	100	100
2	E	244/258 (95%)	243 (100%)	1 (0%)	84	78
2	F	244/258 (95%)	244 (100%)	0	100	100
2	G	244/258 (95%)	243 (100%)	1 (0%)	84	78
2	H	244/258 (95%)	243 (100%)	1 (0%)	84	78
All	All	1289/1460 (88%)	1286 (100%)	3 (0%)	87	84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	264	SER
2	G	264	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	264	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	115	HIS
2	E	201	HIS
2	E	224	ASN
2	F	96	GLN
2	F	201	HIS
2	F	224	ASN
2	G	115	HIS
2	H	96	GLN
2	H	115	HIS
2	H	201	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	D	201	-	5,5,5	0.06	0	5,5,5	0.12	0
3	GOL	C	201	-	5,5,5	0.07	0	5,5,5	0.13	0
3	GOL	B	201	-	5,5,5	0.06	0	5,5,5	0.12	0
3	GOL	A	201	-	5,5,5	0.09	0	5,5,5	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	201	-	-	2/4/4/4	-
3	GOL	C	201	-	-	1/4/4/4	-
3	GOL	B	201	-	-	0/4/4/4	-
3	GOL	A	201	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	201	GOL	C1-C2-C3-O3
3	D	201	GOL	C1-C2-C3-O3
3	D	201	GOL	O2-C2-C3-O3
3	C	201	GOL	C1-C2-C3-O3
3	A	201	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	96/136 (70%)	0.36	6 (6%)	26	27	23, 37, 65, 108	1 (1%)
1	B	96/136 (70%)	0.37	6 (6%)	26	27	22, 37, 67, 109	1 (1%)
1	C	96/136 (70%)	0.33	6 (6%)	26	27	18, 36, 66, 111	2 (2%)
1	D	96/136 (70%)	0.37	6 (6%)	26	27	21, 37, 68, 107	1 (1%)
2	E	276/304 (90%)	0.51	20 (7%)	21	22	25, 42, 77, 97	7 (2%)
2	F	276/304 (90%)	0.57	21 (7%)	20	20	25, 44, 85, 112	7 (2%)
2	G	276/304 (90%)	0.33	14 (5%)	33	35	25, 38, 71, 89	7 (2%)
2	H	276/304 (90%)	0.35	16 (5%)	29	30	25, 39, 69, 93	7 (2%)
All	All	1488/1760 (84%)	0.42	95 (6%)	25	26	18, 40, 75, 112	33 (2%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	234	CYS	5.5
2	G	234	CYS	4.7
2	E	283[A]	THR	4.5
2	H	234	CYS	4.5
2	G	283[A]	THR	4.4
2	F	283[A]	THR	4.4
1	D	24	ALA	4.1
2	E	234	CYS	4.1
1	C	124	MET	4.1
2	F	284[A]	PRO	4.0
1	A	24	ALA	4.0
2	F	79	PRO	4.0
1	B	24	ALA	3.9
2	E	284[A]	PRO	3.8
2	G	67	PRO	3.8
1	C	125	SER	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	222	ALA	3.7
2	H	283[A]	THR	3.6
2	H	284[A]	PRO	3.6
1	C	24	ALA	3.6
2	G	284[A]	PRO	3.6
1	D	89	HIS	3.5
2	H	66	ALA	3.4
1	B	124	MET	3.3
2	H	282[A]	GLU	3.3
2	E	66	ALA	3.3
2	E	67	PRO	3.2
2	E	79	PRO	3.1
1	B	89	HIS	3.1
1	A	124	MET	3.0
2	E	282[A]	GLU	3.0
1	B	125	SER	3.0
1	D	124	MET	3.0
2	H	50	ASP	3.0
2	H	67	PRO	3.0
1	C	89	HIS	2.9
2	G	282[A]	GLU	2.9
2	H	77	GLY	2.9
2	G	79	PRO	2.8
1	D	125	SER	2.8
2	F	178	PHE	2.8
2	F	282[A]	GLU	2.7
1	B	96	THR	2.7
1	A	125	SER	2.7
2	H	280[A]	LEU	2.7
2	H	265	CYS	2.6
2	F	66	ALA	2.6
2	F	160	THR	2.6
2	G	253	CYS	2.6
1	A	89	HIS	2.6
1	D	88	HIS	2.6
2	E	265	CYS	2.6
2	F	69	CYS	2.6
2	G	265	CYS	2.6
2	E	178	PHE	2.6
2	E	158	PRO	2.5
2	E	71	LEU	2.5
2	E	68	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	67	PRO	2.5
2	H	69	CYS	2.5
2	G	66	ALA	2.5
2	G	50	ASP	2.5
2	G	83	TYR	2.4
2	F	202	VAL	2.4
1	A	88	HIS	2.3
2	E	63	VAL	2.3
2	H	48	VAL	2.3
2	F	83	TYR	2.3
2	H	178	PHE	2.3
2	H	224	ASN	2.2
2	F	49	GLY	2.2
2	F	139	PHE	2.2
2	F	224	ASN	2.2
2	F	48	VAL	2.2
1	A	87	ARG	2.2
1	C	88	HIS	2.1
2	H	70	ARG	2.2
2	E	83	TYR	2.1
2	F	81	ASP	2.1
2	G	63	VAL	2.1
2	F	253	CYS	2.1
2	F	265	CYS	2.1
2	G	71	LEU	2.1
1	C	96	THR	2.1
2	F	68	GLY	2.1
2	F	87	GLU	2.1
2	E	280[A]	LEU	2.1
2	H	281[A]	SER	2.1
2	E	50	ASP	2.1
1	D	96	THR	2.0
2	E	224	ASN	2.0
2	E	69	CYS	2.0
2	G	69	CYS	2.0
2	E	77	GLY	2.0
1	B	87	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	201	6/6	0.89	0.14	66,67,71,71	0
3	GOL	B	201	6/6	0.89	0.13	67,69,72,73	0
3	GOL	D	201	6/6	0.89	0.14	66,67,72,73	0
3	GOL	C	201	6/6	0.90	0.11	63,64,68,69	0
4	CL	E	401	1/1	0.96	0.11	49,49,49,49	0
4	CL	F	401	1/1	0.96	0.12	47,47,47,47	0
4	CL	G	401	1/1	0.96	0.15	47,47,47,47	0
4	CL	H	401	1/1	0.96	0.12	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.