



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:55 PM UTC

PDB ID : 6T5M / pdb_00006t5m
Title : Crystal structure of 2-methylisocitrate lyase (PrpB) from *Pseudomonas aeruginosa* in complex with Mg(II)-pyruvate.
Authors : Wijaya, A.J.; Brear, P.; Dolan, S.K.; Welch, M.
Deposited on : 2019-10-16
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	6% 84% 8% 7%
1	B	298	15% 81% 11% 7%
1	C	298	4% 81% 11% 7%
1	D	298	10% 83% 9% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-methylisocitrate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	277	Total	C	N	O	S	0	4	0
			2120	1334	360	412	14			
1	A	276	Total	C	N	O	S	0	0	0
			2085	1313	355	404	13			
1	D	276	Total	C	N	O	S	0	0	0
			2085	1313	355	404	13			
1	B	276	Total	C	N	O	S	0	1	0
			2093	1319	356	405	13			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PYRUVIC ACID (CCD ID: PYR) (formula: C₃H₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	166	Total	O	0	4
			170	170		
4	A	135	Total	O	0	1
			136	136		
4	D	106	Total	O	0	0
			106	106		
4	B	78	Total	O	0	0
			78	78		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-methylisocitrate lyase




- Molecule 1: 2-methylisocitrate lyase




- Molecule 1: 2-methylisocitrate lyase



- Molecule 1: 2-methylisocitrate lyase





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	153.49Å 59.40Å 147.65Å 90.00° 120.30° 90.00°	Depositor
Resolution (Å)	42.49 – 1.76 42.49 – 1.76	Depositor EDS
% Data completeness (in resolution range)	85.4 (42.49-1.76) 85.4 (42.49-1.76)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.76Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.211 , 0.227 0.209 , 0.222	Depositor DCC
R_{free} test set	4808 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 26.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8901	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PYR, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	4/2114 (0.2%)	1.43	7/2864 (0.2%)
1	B	1.11	3/2122 (0.1%)	1.47	1/2875 (0.0%)
1	C	1.22	7/2149 (0.3%)	1.40	4/2913 (0.1%)
1	D	1.15	3/2114 (0.1%)	1.43	2/2864 (0.1%)
All	All	1.16	17/8499 (0.2%)	1.43	14/11516 (0.1%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	256	ILE	C-O	-7.03	1.16	1.24
1	A	20	HIS	C-O	-7.03	1.18	1.24
1	C	8	PRO	C-O	-6.07	1.16	1.24
1	A	22	LEU	C-O	-5.97	1.16	1.24
1	C	15	ALA	C-O	-5.72	1.17	1.24
1	D	266	ILE	C-O	-5.51	1.17	1.24
1	C	213	ILE	C-O	-5.46	1.17	1.24
1	C	227	LEU	C-O	-5.39	1.17	1.24
1	A	26	GLY	C-O	5.39	1.29	1.23
1	B	250	GLU	C-O	-5.34	1.17	1.24
1	C	16	VAL	C-O	-5.32	1.18	1.24
1	B	79	VAL	N-CA	5.30	1.51	1.46
1	C	228	ALA	C-O	-5.27	1.18	1.24
1	A	32	HIS	CE1-NE2	5.21	1.37	1.32
1	D	267	ASP	C-O	-5.11	1.17	1.24
1	D	20	HIS	C-O	-5.08	1.20	1.24
1	C	18	THR	C-O	-5.03	1.17	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	115	HIS	CA-CB-CG	6.28	120.08	113.80
1	A	118	ASP	CA-CB-CG	6.08	118.68	112.60
1	A	89	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	49	GLY	CA-C-N	5.34	125.76	119.94
1	A	49	GLY	C-N-CA	5.34	125.76	119.94
1	A	168	GLY	CA-C-O	-5.31	118.33	122.52
1	B	191	ALA	N-CA-C	5.29	117.11	110.91
1	D	231	ASP	CA-C-N	5.26	127.63	120.63
1	D	231	ASP	C-N-CA	5.26	127.63	120.63
1	C	85	LEU	CA-C-O	-5.16	114.81	120.69
1	C	280	ASP	CA-C-N	5.15	127.18	120.28
1	C	280	ASP	C-N-CA	5.15	127.18	120.28
1	A	46	LEU	CA-C-N	5.11	128.10	120.95
1	A	46	LEU	C-N-CA	5.11	128.10	120.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2085	0	2084	12	0
1	B	2093	0	2094	22	0
1	C	2120	0	2118	22	0
1	D	2085	0	2084	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
4	A	136	0	0	0	0
4	B	78	0	0	0	0
4	C	170	0	0	4	0
4	D	106	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8901	0	8380	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:246:ASN:HD22	1:B:270:GLN:HE22	1.29	0.76
1:C:5:SER:HA	4:C:449:HOH:O	1.86	0.76
1:C:101:ARG:NH2	1:B:56[B]:LEU:HD11	2.06	0.70
1:C:270:GLN:HE22	1:A:246:ASN:HD22	1.39	0.69
1:C:246:ASN:HD22	1:A:270:GLN:HE22	1.41	0.67
1:B:192:MET:HB2	1:B:198:TYR:CZ	2.31	0.66
1:B:192:MET:HB2	1:B:198:TYR:CE2	2.34	0.62
1:D:246:ASN:HD22	1:B:270:GLN:NE2	1.98	0.60
1:C:164:LEU:HD22	1:C:192:MET:HE2	1.87	0.56
1:B:37:LYS:HA	1:B:82:LEU:HD11	1.90	0.54
1:B:164:LEU:HD22	1:B:192:MET:HE2	1.89	0.54
1:C:250:GLU:O	1:C:254[B]:THR:HG23	2.08	0.54
1:C:119:GLN:NE2	4:C:402[A]:HOH:O	2.38	0.53
1:C:5:SER:CA	4:C:449:HOH:O	2.51	0.52
1:D:103:VAL:HG21	1:D:146:ALA:HA	1.92	0.51
1:C:164:LEU:HD22	1:C:192:MET:CE	2.40	0.50
1:A:192:MET:HG3	1:A:197:MET:HE3	1.93	0.50
1:C:195:LEU:HD11	1:C:226:GLU:HB3	1.94	0.49
1:C:246:ASN:HD22	1:A:270:GLN:NE2	2.07	0.49
1:B:117:GLU:HA	1:B:160:ARG:O	2.12	0.49
1:D:177:CYS:O	1:D:181:GLU:HG2	2.13	0.48
1:C:210:LEU:C	1:C:210:LEU:HD23	2.38	0.48
1:A:45:TYR:CG	1:A:236:LEU:HD11	2.49	0.47
1:D:160:ARG:NH2	1:D:190:GLU:CD	2.71	0.47
1:B:177:CYS:O	1:B:181:GLU:HG2	2.14	0.47
1:D:140:VAL:HG12	1:D:144:LYS:HD2	1.96	0.47
1:D:263:LYS:O	1:D:266:ILE:HG12	2.14	0.47
1:A:136:GLN:O	1:A:140:VAL:HG23	2.14	0.47
1:C:115:HIS:HA	1:C:158:MET:O	2.16	0.46
1:C:270:GLN:NE2	1:A:246:ASN:HD22	2.08	0.46
1:A:291:LEU:HD21	1:B:148:ASP:CG	2.42	0.44
1:C:119:GLN:NE2	4:C:402[B]:HOH:O	2.50	0.44
1:B:115:HIS:HA	1:B:158:MET:O	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:45:TYR:CG	1:C:236:LEU:HD11	2.53	0.44
1:B:192:MET:SD	1:B:197:MET:HG2	2.57	0.44
1:A:115:HIS:HA	1:A:158:MET:O	2.17	0.44
1:D:269:MET:SD	1:B:215:GLU:HG2	2.58	0.44
1:B:19:GLU:HG2	1:B:22:LEU:HA	1.99	0.44
1:C:288:LEU:HD22	1:C:292:PHE:HE2	1.82	0.43
1:B:74:ARG:HB2	1:B:109:PHE:HD2	1.82	0.43
1:B:197:MET:HE2	1:B:197:MET:HB2	1.75	0.43
1:D:117:GLU:HA	1:D:160:ARG:O	2.18	0.43
1:D:45:TYR:CG	1:D:236:LEU:HD11	2.54	0.43
1:B:140:VAL:HG13	1:B:182:ALA:HB2	2.01	0.43
1:D:37:LYS:HA	1:D:82:LEU:HD11	2.00	0.43
1:D:115:HIS:HA	1:D:158:MET:O	2.19	0.43
1:B:86:VAL:O	1:B:114:MET:HA	2.19	0.42
1:B:136:GLN:O	1:B:140:VAL:HG23	2.19	0.42
1:C:219[A]:THR:O	1:C:219[A]:THR:HG23	2.20	0.42
1:B:74:ARG:HB2	1:B:109:PHE:CD2	2.54	0.42
1:D:160:ARG:HH21	1:D:190:GLU:CD	2.28	0.41
1:D:20:HIS:HA	1:D:21:PRO:HA	1.81	0.41
1:C:280:ASP:CG	1:C:283:SER:HB3	2.46	0.41
1:A:291:LEU:HD21	1:B:148:ASP:CB	2.51	0.41
1:C:117:GLU:HA	1:C:160:ARG:O	2.21	0.41
1:C:188:PHE:CD1	1:C:188:PHE:C	2.99	0.41
1:D:271:THR:HA	1:B:216:PHE:CE2	2.55	0.41
1:C:95:SER:HB3	1:C:98:ASN:OD1	2.21	0.40
1:A:280:ASP:OD1	1:A:283:SER:HB3	2.21	0.40
1:A:160:ARG:HA	1:A:188:PHE:O	2.22	0.40
1:D:263:LYS:HB2	1:D:263:LYS:HE3	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/298 (91%)	265 (97%)	7 (3%)	0	100	100
1	B	273/298 (92%)	265 (97%)	8 (3%)	0	100	100
1	C	277/298 (93%)	270 (98%)	7 (2%)	0	100	100
1	D	272/298 (91%)	265 (97%)	7 (3%)	0	100	100
All	All	1094/1192 (92%)	1065 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/235 (93%)	215 (99%)	3 (1%)	59	44
1	B	219/235 (93%)	214 (98%)	5 (2%)	44	24
1	C	223/235 (95%)	220 (99%)	3 (1%)	61	46
1	D	218/235 (93%)	212 (97%)	6 (3%)	38	18
All	All	878/940 (93%)	861 (98%)	17 (2%)	50	31

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	188	PHE
1	C	197	MET
1	C	245	MET
1	A	188	PHE
1	A	245	MET
1	A	263	LYS
1	D	6	LEU
1	D	94	SER
1	D	188	PHE
1	D	245	MET
1	D	267	ASP
1	D	288	LEU
1	B	188	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	190	GLU
1	B	197	MET
1	B	245	MET
1	B	272	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	20	HIS
1	C	136	GLN
1	C	270	GLN
1	C	282	HIS
1	A	270	GLN
1	A	282	HIS
1	D	136	GLN
1	D	270	GLN
1	B	264	ASN
1	B	270	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PYR	B	302	2	5,5,5	2.61	1 (20%)	3,6,6	1.88	1 (33%)
3	PYR	A	302	2	5,5,5	3.25	2 (40%)	3,6,6	1.19	0
3	PYR	C	302	2	5,5,5	2.84	1 (20%)	3,6,6	2.31	1 (33%)
3	PYR	D	302	2	5,5,5	2.81	1 (20%)	3,6,6	1.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYR	B	302	2	-	0/4/4/4	-
3	PYR	A	302	2	-	0/4/4/4	-
3	PYR	C	302	2	-	0/4/4/4	-
3	PYR	D	302	2	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	PYR	CA-C	-6.71	1.32	1.54
3	D	302	PYR	CA-C	-5.94	1.34	1.54
3	C	302	PYR	CA-C	-5.64	1.35	1.54
3	B	302	PYR	CA-C	-5.40	1.36	1.54
3	A	302	PYR	OXT-C	-2.77	1.23	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	302	PYR	OXT-C-CA	3.75	123.98	113.59
3	B	302	PYR	OXT-C-CA	2.50	120.53	113.59

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/298 (92%)	0.55	17 (6%) 26 30	16, 27, 56, 68	0
1	B	276/298 (92%)	1.09	44 (15%) 5 5	11, 35, 63, 97	1 (0%)
1	C	277/298 (92%)	0.29	13 (4%) 36 42	9, 24, 42, 72	4 (1%)
1	D	276/298 (92%)	0.84	29 (10%) 11 13	21, 31, 48, 70	0
All	All	1105/1192 (92%)	0.69	103 (9%) 14 17	9, 29, 56, 97	5 (0%)

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	288	LEU	6.0
1	D	290	ALA	5.7
1	B	292	PHE	5.2
1	A	191	ALA	5.1
1	A	292	PHE	5.0
1	C	120	VAL	5.0
1	C	292	PHE	4.6
1	D	120	VAL	4.5
1	D	292	PHE	4.2
1	B	291	LEU	4.0
1	A	120	VAL	3.9
1	B	166	VAL	3.8
1	C	288	LEU	3.7
1	B	191	ALA	3.7
1	B	133	ILE	3.7
1	C	291	LEU	3.5
1	B	120	VAL	3.5
1	D	285	GLU	3.5
1	D	163	ALA	3.4
1	B	290	ALA	3.3
1	D	288	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	162	ASP	3.2
1	B	6	LEU	3.2
1	B	173	ILE	3.2
1	A	164	LEU	3.2
1	D	109	PHE	3.2
1	C	285	GLU	3.2
1	B	169	LEU	3.1
1	B	109	PHE	3.1
1	A	165	ALA	3.1
1	A	190	GLU	3.1
1	B	221	LEU	3.0
1	C	191	ALA	3.0
1	A	166	VAL	2.9
1	D	289	ASP	2.9
1	D	281	TYR	2.9
1	B	222	PHE	2.9
1	A	290	ALA	2.8
1	B	205	VAL	2.8
1	A	291	LEU	2.8
1	B	287	LYS	2.8
1	C	282	HIS	2.8
1	B	165	ALA	2.7
1	A	94	SER	2.7
1	D	191	ALA	2.7
1	A	288	LEU	2.7
1	C	267	ASP	2.7
1	D	6	LEU	2.7
1	D	291	LEU	2.7
1	D	168	GLY	2.6
1	B	168	GLY	2.6
1	B	171	ALA	2.6
1	A	193	THR	2.6
1	B	265	VAL	2.6
1	C	290	ALA	2.5
1	B	203	ALA	2.5
1	B	103	VAL	2.4
1	B	190	GLU	2.3
1	A	132	GLU	2.3
1	D	138	GLU	2.3
1	C	96	ALA	2.3
1	B	281	TYR	2.3
1	B	132	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	194	GLU	2.3
1	B	285	GLU	2.3
1	B	134	VAL	2.3
1	B	228	ALA	2.3
1	A	169	LEU	2.3
1	C	162	ASP	2.3
1	D	166	VAL	2.3
1	B	282	HIS	2.2
1	B	200	GLU	2.2
1	B	177	CYS	2.2
1	B	202	ALA	2.2
1	B	206	LYS	2.2
1	D	181	GLU	2.2
1	D	219	THR	2.2
1	D	228	ALA	2.2
1	A	286	GLN	2.2
1	D	217	GLY	2.2
1	B	195	LEU	2.2
1	D	276	TYR	2.2
1	A	283	SER	2.2
1	B	280	ASP	2.1
1	C	286	GLN	2.1
1	D	206	LYS	2.1
1	D	284	PHE	2.1
1	B	289	ASP	2.1
1	B	266	ILE	2.1
1	B	207	VAL	2.1
1	C	20	HIS	2.1
1	D	165	ALA	2.1
1	D	202	ALA	2.1
1	B	180	VAL	2.1
1	D	39	ALA	2.0
1	B	211	ALA	2.0
1	B	94	SER	2.0
1	A	282	HIS	2.0
1	D	282	HIS	2.0
1	B	263	LYS	2.0
1	D	273	MET	2.0
1	B	284	PHE	2.0
1	D	171	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PYR	D	302	6/6	0.84	0.12	31,32,37,39	0
3	PYR	B	302	6/6	0.88	0.11	32,36,37,41	0
3	PYR	A	302	6/6	0.91	0.09	29,31,34,34	0
3	PYR	C	302	6/6	0.96	0.08	21,23,26,29	0
2	MG	B	301	1/1	0.96	0.07	44,44,44,44	0
2	MG	D	301	1/1	0.97	0.06	33,33,33,33	0
2	MG	A	301	1/1	0.98	0.04	33,33,33,33	0
2	MG	C	301	1/1	0.99	0.05	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PYR D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PYR B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around PYR A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around PYR C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.