



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:44 PM UTC

PDB ID : 8T5B / pdb_00008t5b
Title : HIV-1 Integrase Catalytic Core Domain and C-Terminal Domain in Complex with Allosteric Integrase Inhibitor EKC-110
Authors : Dinh, T.; Kvaratskhelia, M.
Deposited on : 2023-06-12
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	
1	B	155	
2	C	57	
2	D	57	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6736 atoms, of which 3349 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	150	Total	C	H	N	O	S	25	3	0
			2339	743	1173	204	215	4			
1	B	148	Total	C	H	N	O	S	28	0	0
			2293	727	1154	198	210	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	LYS	PHE	conflict	UNP P12497
B	185	LYS	PHE	conflict	UNP P12497


- Molecule 2 is a protein called Integrase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	56	Total	C	H	N	O	S	0	1	0
			936	292	476	87	80	1			
2	D	57	Total	C	H	N	O	S	0	1	0
			960	297	490	91	81	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	243	GLU	TRP	conflict	UNP P12497
D	243	GLU	TRP	conflict	UNP P12497

- Molecule 3 is (2S)-tert-butoxy{4-(4-chlorophenyl)-2,6-dimethyl-1-[(1-methyl-1H-pyrazol-4-yl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl}acetic acid (CCD ID: QD6) (formula: C₂₆H₂₉ClN₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	H	N	O	0	0
			62	26	1	28	4	3		
3	B	1	Total	C	Cl	H	N	O	0	0
			62	26	1	28	4	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	1	0
			1	1		
4	B	1	Total	Mg	1	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	39	Total	O	0	0
			39	39		
5	B	33	Total	O	0	0
			33	33		
5	C	6	Total	O	0	0
			6	6		
5	D	4	Total	O	0	0
			4	4		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Integrase





- Molecule 1: Integrase



- Molecule 2: Integrase



- Molecule 2: Integrase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.95Å 69.98Å 63.86Å 90.00° 100.73° 90.00°	Depositor
Resolution (Å)	46.72 – 2.08 46.72 – 2.08	Depositor EDS
% Data completeness (in resolution range)	97.2 (46.72-2.08) 97.2 (46.72-2.08)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 2.08Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.240 , 0.265 0.241 , 0.262	Depositor DCC
R_{free} test set	1999 reflections (6.18%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.085 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6736	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QD6, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/1194	0.33	0/1615
1	B	0.18	0/1159	0.34	0/1563
2	C	0.15	0/471	0.39	0/631
2	D	0.29	0/481	0.52	0/643
All	All	0.19	0/3305	0.38	0/4452

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1166	1173	1171	4	0
1	B	1139	1154	1153	12	0
2	C	460	476	476	8	0
2	D	470	490	490	4	0
3	A	34	28	0	0	0
3	B	34	28	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	39	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	33	0	0	0	0
5	C	6	0	0	0	0
5	D	4	0	0	0	0
All	All	3387	3349	3290	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:249:VAL:HG12	2:C:251:ILE:HG23	1.85	0.58
2:D:249:VAL:HG12	2:D:251:ILE:HG23	1.88	0.56
1:A:186:LYS:O	1:A:187:ARG:CB	2.54	0.56
2:D:244:LYS:HA	2:D:249:VAL:HG22	1.89	0.54
2:D:221:GLN:HA	2:D:269[A]:ARG:HE	1.72	0.54
1:B:68:LEU:HD21	1:B:159:LYS:CD	2.41	0.51
1:B:186:LYS:O	1:B:187:ARG:CB	2.59	0.50
2:C:245:GLY:O	2:C:262:ARG:NH1	2.44	0.49
1:B:103:LYS:O	1:B:107:ARG:HG2	2.13	0.48
2:C:244:LYS:HA	2:C:249:VAL:HG22	1.96	0.48
2:C:224[A]:ARG:HD3	2:C:238:PRO:HB3	1.96	0.47
1:B:182:ILE:HG23	2:C:275:MET:HE2	1.96	0.47
1:A:107[A]:ARG:HG3	1:B:83:TYR:OH	2.15	0.46
1:B:58:PRO:HB3	1:B:203:ILE:HG22	1.97	0.45
2:C:249:VAL:HG21	2:C:267:ILE:HD11	2.00	0.44
2:C:269:ARG:HG3	2:C:271:TYR:HB2	1.99	0.44
2:C:250:VAL:HG22	2:C:259:VAL:HG22	1.99	0.44
1:B:62:GLN:OE1	1:B:151:ILE:HB	2.17	0.44
1:B:77:VAL:HG12	1:B:151:ILE:HD13	2.00	0.43
1:B:68:LEU:HD21	1:B:159:LYS:HD3	2.02	0.41
1:B:156:LYS:HD3	1:B:157:GLU:N	2.36	0.41
2:D:241:LEU:CD1	2:D:249:VAL:HG13	2.51	0.40
1:A:107[B]:ARG:HG2	1:A:108:TRP:CD1	2.56	0.40
1:B:68:LEU:HD21	1:B:159:LYS:HD2	2.02	0.40
1:A:107[B]:ARG:HD3	1:B:85:GLU:OE2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/155 (96%)	143 (96%)	4 (3%)	2 (1%)	9	5
1	B	142/155 (92%)	139 (98%)	2 (1%)	1 (1%)	18	14
2	C	55/57 (96%)	51 (93%)	4 (7%)	0	100	100
2	D	56/57 (98%)	52 (93%)	4 (7%)	0	100	100
All	All	402/424 (95%)	385 (96%)	14 (4%)	3 (1%)	16	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	ARG
1	B	187	ARG
1	A	188	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/126 (96%)	117 (97%)	4 (3%)	33	35
1	B	118/126 (94%)	115 (98%)	3 (2%)	42	46
2	C	48/48 (100%)	47 (98%)	1 (2%)	47	52
2	D	49/48 (102%)	47 (96%)	2 (4%)	27	27
All	All	336/348 (97%)	326 (97%)	10 (3%)	39	39

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	SER
1	A	107[A]	ARG
1	A	107[B]	ARG
1	A	160	LYS
1	B	57	SER
1	B	87	GLU
1	B	156	LYS
2	C	255	SER
2	D	273	LYS
2	D	274	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	164	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	QD6	B	301	-	35,37,37	4.25	31 (88%)	40,56,56	3.28	13 (32%)
3	QD6	A	301	-	35,37,37	4.33	31 (88%)	40,56,56	3.21	11 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QD6	B	301	-	-	3/21/21/21	0/4/4/4
3	QD6	A	301	-	-	3/21/21/21	0/4/4/4

All (62) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	QD6	C12-C11	8.13	1.63	1.50
3	B	301	QD6	C18-C17	8.12	1.64	1.49
3	A	301	QD6	C18-C17	8.08	1.64	1.49
3	B	301	QD6	C12-C11	8.04	1.63	1.50
3	A	301	QD6	C24-N23	6.69	1.54	1.46
3	B	301	QD6	C24-N23	6.58	1.54	1.46
3	A	301	QD6	C14-N13	6.56	1.42	1.34
3	B	301	QD6	C14-N13	6.18	1.42	1.34
3	A	301	QD6	C27-C28	6.10	1.60	1.50
3	B	301	QD6	C27-C28	5.96	1.60	1.50
3	A	301	QD6	C20-N19	5.15	1.56	1.46
3	A	301	QD6	C31-CL32	5.05	1.86	1.74
3	B	301	QD6	C27-C10	5.05	1.48	1.40
3	B	301	QD6	C20-N19	4.95	1.56	1.46
3	B	301	QD6	C31-CL32	4.93	1.86	1.74
3	A	301	QD6	C27-C10	4.93	1.48	1.40
3	A	301	QD6	C30-C31	4.40	1.46	1.38
3	A	301	QD6	C11-N13	4.36	1.40	1.34
3	A	301	QD6	C33-C31	4.34	1.46	1.38
3	A	301	QD6	O05-C06	4.33	1.54	1.45
3	B	301	QD6	C33-C31	4.32	1.46	1.38
3	A	301	QD6	C20-C21	4.32	1.63	1.50
3	B	301	QD6	C08-C06	4.27	1.63	1.51
3	A	301	QD6	C08-C06	4.26	1.63	1.51
3	A	301	QD6	C15-C27	4.26	1.48	1.42
3	B	301	QD6	C30-C31	4.26	1.45	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	QD6	C20-C21	4.19	1.63	1.50
3	B	301	QD6	O05-C06	4.17	1.54	1.45
3	B	301	QD6	C15-C27	4.15	1.48	1.42
3	B	301	QD6	C11-N13	4.11	1.40	1.34
3	A	301	QD6	C07-C06	4.10	1.62	1.51
3	B	301	QD6	C07-C06	4.07	1.62	1.51
3	A	301	QD6	C30-C29	3.98	1.45	1.38
3	B	301	QD6	C26-C21	3.94	1.47	1.40
3	A	301	QD6	C26-C21	3.93	1.47	1.40
3	B	301	QD6	C30-C29	3.88	1.45	1.38
3	A	301	QD6	O03-C02	3.82	1.33	1.22
3	A	301	QD6	C34-C33	3.82	1.45	1.38
3	B	301	QD6	O03-C02	3.74	1.33	1.22
3	B	301	QD6	C16-C17	3.72	1.47	1.37
3	A	301	QD6	C16-C17	3.68	1.47	1.37
3	B	301	QD6	C34-C33	3.65	1.44	1.38
3	A	301	QD6	C26-N25	3.62	1.38	1.32
3	B	301	QD6	C26-N25	3.54	1.38	1.32
3	A	301	QD6	C29-C28	3.44	1.46	1.39
3	B	301	QD6	N23-N25	3.43	1.39	1.35
3	A	301	QD6	N23-N25	3.41	1.39	1.35
3	B	301	QD6	C29-C28	3.37	1.46	1.39
3	A	301	QD6	C22-C21	3.30	1.49	1.38
3	A	301	QD6	C09-C06	3.24	1.60	1.51
3	B	301	QD6	C09-C06	3.23	1.60	1.51
3	B	301	QD6	C22-C21	3.21	1.48	1.38
3	A	301	QD6	C34-C28	3.18	1.45	1.39
3	B	301	QD6	C34-C28	3.18	1.45	1.39
3	A	301	QD6	C22-N23	3.08	1.37	1.34
3	B	301	QD6	C22-N23	2.97	1.37	1.34
3	A	301	QD6	O05-C04	-2.84	1.38	1.43
3	B	301	QD6	O05-C04	-2.74	1.39	1.43
3	A	301	QD6	C10-C04	2.24	1.57	1.53
3	A	301	QD6	C15-C14	2.22	1.43	1.40
3	B	301	QD6	C10-C04	2.17	1.57	1.53
3	B	301	QD6	C15-C14	2.08	1.43	1.40

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	QD6	C21-C26-N25	-12.15	107.13	113.14
3	A	301	QD6	C21-C26-N25	-11.74	107.33	113.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	QD6	C26-N25-N23	10.25	110.04	104.41
3	A	301	QD6	C26-N25-N23	10.08	109.94	104.41
3	A	301	QD6	C15-C14-N13	-7.90	121.54	126.56
3	B	301	QD6	C15-C14-N13	-7.53	121.78	126.56
3	B	301	QD6	C24-N23-N25	5.51	124.45	120.03
3	A	301	QD6	C24-N23-N25	5.07	124.10	120.03
3	A	301	QD6	C14-N13-C11	3.58	121.63	117.17
3	B	301	QD6	C14-N13-C11	3.49	121.51	117.17
3	B	301	QD6	C24-N23-C22	-3.19	124.74	127.94
3	A	301	QD6	C18-C17-N19	3.15	126.09	122.75
3	B	301	QD6	C18-C17-N19	2.88	125.81	122.75
3	A	301	QD6	C24-N23-C22	-2.88	125.05	127.94
3	A	301	QD6	N13-C14-N19	2.85	130.46	125.67
3	B	301	QD6	C21-C20-N19	-2.66	109.38	113.32
3	B	301	QD6	N13-C14-N19	2.58	130.01	125.67
3	B	301	QD6	C34-C28-C29	2.54	122.22	117.68
3	A	301	QD6	C34-C28-C29	2.42	122.02	117.68
3	B	301	QD6	C34-C28-C27	-2.40	116.68	120.80
3	B	301	QD6	C22-N23-N25	-2.25	110.81	112.08
3	A	301	QD6	C22-N23-N25	-2.17	110.85	112.08
3	A	301	QD6	C15-C16-C17	-2.04	105.82	108.01
3	B	301	QD6	C30-C29-C28	-2.01	118.55	121.12

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	301	QD6	O01-C02-C04-C10
3	B	301	QD6	O03-C02-C04-C10
3	A	301	QD6	O01-C02-C04-C10
3	A	301	QD6	O03-C02-C04-C10
3	A	301	QD6	C02-C04-O05-C06
3	B	301	QD6	C02-C04-O05-C06


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	150/155 (96%)	0.73	24 (16%)	5 5	19, 39, 72, 92	7 (4%)
1	B	148/155 (95%)	0.79	23 (15%)	5 5	31, 41, 70, 90	5 (3%)
2	C	56/57 (98%)	1.53	15 (26%)	1 1	24, 57, 78, 90	1 (1%)
2	D	57/57 (100%)	1.53	18 (31%)	1 1	30, 57, 86, 106	1 (1%)
All	All	411/424 (96%)	0.97	80 (19%)	3 3	19, 42, 78, 106	14 (3%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	150	VAL	5.8
1	A	191	ILE	5.5
1	A	143	TYR	5.0
2	D	277	GLY	4.9
1	A	149	GLY	4.9
2	C	276	ALA	4.7
1	B	190	GLY	4.3
2	C	222	ASN	4.1
1	A	150	VAL	4.1
1	A	210	THR	4.0
1	B	155	ASN	4.0
1	A	189	GLY	4.0
1	A	152	GLU	3.9
1	B	210	THR	3.8
1	B	149	GLY	3.6
2	D	223	PHE	3.6
1	B	211	LYS	3.6
2	D	256	ASP	3.6
1	A	107[A]	ARG	3.6
1	B	138	GLU	3.5
2	D	269[A]	ARG	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	254	ASN	3.5
1	B	140	GLY	3.4
1	A	144	ASN	3.3
2	D	222	ASN	3.3
2	C	229	ASP	3.3
1	B	152	GLU	3.3
2	D	276	ALA	3.3
2	D	245	GLY	3.2
1	B	151	ILE	3.2
1	A	57	SER	3.1
2	D	275	MET	3.1
2	D	221	GLN	3.1
2	C	252	GLN	3.1
2	C	221	GLN	3.0
1	A	151	ILE	2.9
1	B	58	PRO	2.9
2	C	230	SER	2.9
2	C	224[A]	ARG	2.8
1	B	57	SER	2.8
2	C	254	ASN	2.8
1	A	188	LYS	2.8
2	D	255	SER	2.7
2	D	257	ILE	2.7
1	B	139	PHE	2.6
1	A	209	GLN	2.6
1	B	156	LYS	2.6
1	B	148	GLN	2.5
1	B	186	LYS	2.5
1	A	187	ARG	2.5
1	B	209	GLN	2.5
1	A	156	LYS	2.5
1	B	194	TYR	2.4
2	D	271	TYR	2.4
1	B	153	SER	2.4
1	A	206	THR	2.4
2	C	244	LYS	2.4
1	A	157	GLU	2.3
2	D	274	GLN	2.3
1	B	114	HIS	2.3
1	A	141	ILE	2.3
2	D	249	VAL	2.2
2	D	242	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	231	ARG	2.2
1	A	190	GLY	2.2
2	C	260	VAL	2.1
2	C	242	LEU	2.1
2	C	263	ARG	2.1
2	D	273	LYS	2.1
1	B	154	MET	2.1
1	B	191	ILE	2.1
2	C	249	VAL	2.1
1	A	192	GLY	2.1
1	A	58	PRO	2.0
1	A	194	TYR	2.0
2	C	236	LYS	2.0
2	D	244	LYS	2.0
1	A	87[A]	GLU	2.0
1	A	153	SER	2.0
1	B	70	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	QD6	A	301	34/34	0.92	0.12	29,36,43,56	0
3	QD6	B	301	34/34	0.93	0.11	29,36,47,53	0
4	MG	A	302	1/1	-	-	31,31,31,31	1
4	MG	B	302	1/1	-	-	40,40,40,40	1

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around QD6 A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around QD6 B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.