



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:11 PM UTC

PDB ID : 5T68 / pdb_00005t68
Title : Crystal structure of Syk catalytic domain in complex with a furo[3,2-d]pyrimidine
Authors : Argiriadi, M.A.; Hoemann, M.; Wilson, N.; Banach, D.; Burchat, A.; Calderwood, D.; Clapham, B.; Cox, P.; Duignan, D.B.; Konopacki, D.; Somal, G.; Vasudevan, A.
Deposited on : 2016-09-01
Resolution : 2.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1159 (2.96-2.92)
Clashscore	190562	1184 (2.96-2.92)
Ramachandran outliers	187476	1131 (2.96-2.92)
Sidechain outliers	187428	1131 (2.96-2.92)
RSRZ outliers	180081	1159 (2.96-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	
1	B	291	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase SYK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	0
			2075	1341	343	372	19			
1	B	264	Total	C	N	O	S	0	0	0
			2078	1341	343	375	19			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP P43405
A	354	ALA	-	expression tag	UNP P43405
A	355	LEU	-	expression tag	UNP P43405
A	636	GLU	-	expression tag	UNP P43405
A	637	GLY	-	expression tag	UNP P43405
A	638	HIS	-	expression tag	UNP P43405
A	639	HIS	-	expression tag	UNP P43405
A	640	HIS	-	expression tag	UNP P43405
A	641	HIS	-	expression tag	UNP P43405
A	642	HIS	-	expression tag	UNP P43405
A	643	HIS	-	expression tag	UNP P43405
B	353	MET	-	initiating methionine	UNP P43405
B	354	ALA	-	expression tag	UNP P43405
B	355	LEU	-	expression tag	UNP P43405
B	636	GLU	-	expression tag	UNP P43405
B	637	GLY	-	expression tag	UNP P43405
B	638	HIS	-	expression tag	UNP P43405
B	639	HIS	-	expression tag	UNP P43405
B	640	HIS	-	expression tag	UNP P43405
B	641	HIS	-	expression tag	UNP P43405
B	642	HIS	-	expression tag	UNP P43405
B	643	HIS	-	expression tag	UNP P43405

- Molecule 2 is N 4 -cyclopropyl-N 2 -(3-methyl-1H-indazol-6-yl)furo[3,2-d]pyrimidine-2,4-diamine (CCD ID: 77V) (formula: C₁₇H₁₆N₆O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			24	17	6	1		
2	B	1	Total	C	N	O	0	0
			24	17	6	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		
3	B	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase SYK



• Molecule 1: Tyrosine-protein kinase SYK



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.60Å 40.03Å 84.05Å 90.16° 89.98° 99.80°	Depositor
Resolution (Å)	84.05 – 2.93 84.05 – 2.94	Depositor EDS
% Data completeness (in resolution range)	96.7 (84.05-2.93) 96.7 (84.05-2.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.96Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.183 , 0.281 0.195 , 0.294	Depositor DCC
R_{free} test set	492 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.380 for -h,-k,l 0.036 for -k,-h,-l 0.043 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4224	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 77V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.89	0/2122	1.41	15/2872 (0.5%)
1	B	0.89	0/2125	1.43	9/2875 (0.3%)
All	All	0.89	0/4247	1.42	24/5747 (0.4%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	503	VAL	N-CA-C	-7.87	104.13	111.45
1	B	412	ALA	N-CA-C	-7.15	105.09	113.88
1	B	603	ASP	CA-CB-CG	6.02	118.62	112.60
1	A	494	ASP	CA-CB-CG	5.97	118.57	112.60
1	B	505	GLN	N-CA-C	-5.87	105.83	112.87
1	B	412	ALA	CA-C-N	5.81	128.86	120.38
1	B	412	ALA	C-N-CA	5.81	128.86	120.38
1	A	461	GLN	CA-C-N	5.80	127.99	120.44
1	A	461	GLN	C-N-CA	5.80	127.99	120.44
1	A	366	ASP	CA-CB-CG	5.71	118.31	112.60
1	A	535	PRO	CA-C-N	5.69	132.22	121.97
1	A	535	PRO	C-N-CA	5.69	132.22	121.97
1	A	559	GLY	CA-C-N	5.55	128.07	120.46
1	A	559	GLY	C-N-CA	5.55	128.07	120.46
1	B	619	PHE	CA-CB-CG	-5.50	108.30	113.80
1	A	613	VAL	CB-CA-C	-5.31	104.97	112.14
1	B	621	ALA	CA-C-N	5.27	127.31	120.56
1	B	621	ALA	C-N-CA	5.27	127.31	120.56
1	A	626	LEU	CA-C-N	5.20	127.20	120.44
1	A	626	LEU	C-N-CA	5.20	127.20	120.44
1	A	607	LEU	CA-C-N	5.13	127.16	120.28
1	A	607	LEU	C-N-CA	5.13	127.16	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	554	ASP	CA-CB-CG	5.02	117.62	112.60
1	A	502	LEU	N-CA-C	5.00	117.40	109.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2075	0	1990	21	0
1	B	2078	0	2002	32	0
2	A	24	0	0	4	0
2	B	24	0	0	1	0
3	A	13	0	0	0	0
3	B	10	0	0	0	0
All	All	4224	0	3992	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:LEU:HD13	2:A:701:77V:C3	2.04	0.87
1:B:426:GLN:OE1	1:B:490:PHE:HZ	1.59	0.83
1:B:392:MET:HG2	1:B:393:LYS:H	1.59	0.68
1:A:377:LEU:CD1	2:A:701:77V:N14	2.62	0.62
1:A:404:LEU:HG	1:A:417:LEU:HD13	1.86	0.57
1:A:377:LEU:HD13	2:A:701:77V:N14	2.20	0.57
1:B:392:MET:CG	1:B:393:LYS:H	2.17	0.57
1:B:607:LEU:HD23	1:B:617:PRO:HG3	1.87	0.57
1:B:392:MET:HG2	1:B:393:LYS:N	2.20	0.56
1:A:535:PRO:HD2	1:A:539:TYR:CZ	2.42	0.55
1:B:500:VAL:HG21	1:B:561:LEU:HD11	1.89	0.55
1:B:551:SER:O	1:B:555:VAL:HG23	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:701:77V:C3	2:B:701:77V:N14	2.71	0.53
1:A:624:LEU:HG	1:A:628:ASN:HD21	1.74	0.53
1:B:387:LYS:HE3	1:B:450:MET:HE2	1.90	0.52
1:A:476:HIS:CD2	1:A:480:MET:HE2	2.45	0.52
1:B:439:CYS:HB3	1:B:445:MET:HB2	1.92	0.52
1:B:543:CYS:HB3	1:B:549:PHE:CE1	2.46	0.50
1:A:436:ILE:HB	1:A:447:VAL:HG12	1.94	0.50
1:B:601:MET:HA	1:B:601:MET:HE2	1.95	0.49
1:B:535:PRO:HD2	1:B:539:TYR:CZ	2.47	0.49
1:A:601:MET:HE2	1:A:601:MET:HA	1.95	0.48
1:A:402:LYS:HB3	1:A:446:LEU:HB3	1.95	0.48
1:A:551:SER:O	1:A:555:VAL:HG23	2.13	0.48
1:B:494:ASP:HB2	1:B:515:LEU:HD12	1.96	0.48
1:B:433:VAL:HG21	1:B:511:SER:HB3	1.97	0.47
1:A:607:LEU:HD23	1:A:617:PRO:HG3	1.96	0.46
1:A:562:MET:O	1:A:566:PHE:HD2	1.99	0.46
1:B:500:VAL:CG2	1:B:561:LEU:HD11	2.47	0.45
1:B:490:PHE:CE1	1:B:518:ALA:HB2	2.52	0.44
1:B:528:ALA:HB3	1:B:547:TYR:HB3	1.99	0.44
1:B:461:GLN:HG3	1:B:569:GLY:HA3	1.99	0.43
1:B:482:MET:HE2	1:B:554:ASP:HB3	2.01	0.43
1:B:392:MET:SD	1:B:397:LYS:HB2	2.58	0.43
1:B:504:THR:C	1:B:506:HIS:H	2.25	0.43
1:B:504:THR:C	1:B:506:HIS:N	2.76	0.43
1:A:476:HIS:HE1	1:A:627:ARG:HB2	1.84	0.42
1:A:430:PRO:HG2	1:A:431:TYR:CE2	2.54	0.42
1:A:543:CYS:SG	1:A:553:SER:HB2	2.59	0.42
1:B:576:MET:HE2	1:B:580:GLU:HB3	2.01	0.42
1:B:402:LYS:HE3	1:B:448:MET:HE1	2.02	0.42
1:B:415:ASP:HA	1:B:418:LEU:HB2	2.01	0.42
1:A:476:HIS:CE1	1:A:627:ARG:CA	3.02	0.41
1:B:476:HIS:CE1	1:B:480:MET:HE2	2.56	0.41
1:B:584:MET:HG2	1:B:589:GLU:HB2	2.02	0.41
1:B:562:MET:O	1:B:566:PHE:HD1	2.03	0.41
1:B:590:ARG:HD2	1:B:609:TRP:HB3	2.02	0.41
1:A:537:LYS:HE2	1:A:573:TYR:HB2	2.03	0.40
1:A:377:LEU:HD22	1:A:450:MET:HE1	2.03	0.40
2:A:701:77V:C3	2:A:701:77V:N14	2.85	0.40
1:B:402:LYS:HB3	1:B:446:LEU:HB3	2.04	0.40
1:B:540:ALA:HB2	1:B:556:TRP:CB	2.52	0.40
1:B:563:TRP:CZ3	1:B:572:PRO:HA	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:476:HIS:CE1	1:A:627:ARG:HB2	2.56	0.40
1:A:476:HIS:ND1	1:A:626:LEU:HB3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/291 (89%)	234 (90%)	18 (7%)	8 (3%)	3	8
1	B	258/291 (89%)	240 (93%)	15 (6%)	3 (1%)	10	27
All	All	518/582 (89%)	474 (92%)	33 (6%)	11 (2%)	5	15

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	393	LYS
1	A	536	VAL
1	B	393	LYS
1	B	433	VAL
1	A	568	TYR
1	B	452	GLU
1	A	512	ASP
1	A	636	GLU
1	A	391	GLN
1	A	633	VAL
1	A	395	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/253 (81%)	193 (94%)	13 (6%)	16	36
1	B	210/253 (83%)	195 (93%)	15 (7%)	13	31
All	All	416/506 (82%)	388 (93%)	28 (7%)	15	34

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	392	MET
1	A	398	THR
1	A	401	VAL
1	A	445	MET
1	A	469	LYS
1	A	485	LEU
1	A	491	VAL
1	A	502	LEU
1	A	504	THR
1	A	524	ASN
1	A	534	TRP
1	A	613	VAL
1	A	633	VAL
1	B	413	LEU
1	B	438	ILE
1	B	449	GLU
1	B	458	LYS
1	B	470	ASN
1	B	485	LEU
1	B	489	ASN
1	B	498	ARG
1	B	519	LEU
1	B	534	TRP
1	B	587	LYS
1	B	607	LEU
1	B	622	VAL
1	B	623	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	633	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	422	ASN
1	A	476	HIS
1	A	505	GLN
1	A	506	HIS
1	A	628	ASN
1	B	381	ASN
1	B	425	GLN
1	B	465	HIS
1	B	476	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	77V	A	701	-	27,28,28	1.21	4 (14%)	32,41,41	2.15	13 (40%)
2	77V	B	701	-	27,28,28	1.23	3 (11%)	32,41,41	2.09	13 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	77V	A	701	-	-	0/8/10/10	0/5/5/5
2	77V	B	701	-	-	0/8/10/10	0/5/5/5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	77V	C15-N20	3.85	1.41	1.35
2	A	701	77V	C15-N20	3.62	1.40	1.35
2	B	701	77V	C18-C17	2.52	1.38	1.33
2	A	701	77V	C18-C17	2.43	1.37	1.33
2	A	701	77V	C5-C4	-2.41	1.38	1.41
2	B	701	77V	C5-C4	-2.16	1.38	1.41
2	A	701	77V	C13-N19	2.14	1.41	1.36

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	77V	C15-N20-C21	-6.06	113.95	124.18
2	A	701	77V	C15-N20-C21	-5.37	115.11	124.18
2	B	701	77V	C5-C9-N8	4.15	112.46	110.66
2	A	701	77V	C5-C9-N8	3.98	112.39	110.66
2	A	701	77V	C4-N7-N8	3.75	114.68	111.90
2	A	701	77V	C10-O16-C17	3.56	107.48	105.31
2	B	701	77V	C1-C6-C5	-3.34	115.35	120.86
2	A	701	77V	C10-C15-N20	3.27	124.72	119.88
2	A	701	77V	C1-C6-C5	-2.83	116.20	120.86
2	B	701	77V	C6-C1-C2	2.75	123.47	120.30
2	B	701	77V	N20-C15-N14	2.61	123.57	118.55
2	A	701	77V	O16-C17-C18	-2.60	110.35	112.49
2	A	701	77V	C6-C1-C2	2.59	123.28	120.30
2	B	701	77V	C10-O16-C17	2.56	106.86	105.31
2	B	701	77V	C11-N12-C13	-2.55	113.96	116.69
2	B	701	77V	O16-C17-C18	-2.54	110.40	112.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	77V	C4-N7-N8	2.46	113.72	111.90
2	A	701	77V	C22-C21-N20	2.44	123.16	118.06
2	B	701	77V	C10-C15-N14	-2.40	114.69	121.94
2	A	701	77V	C13-N14-C15	2.40	122.25	116.36
2	A	701	77V	C23-C21-N20	2.39	123.05	118.06
2	A	701	77V	C10-C15-N14	-2.27	115.09	121.94
2	B	701	77V	C13-N14-C15	2.18	121.70	116.36
2	A	701	77V	C5-C4-N7	-2.15	105.05	106.59
2	B	701	77V	C2-C3-C4	-2.04	115.47	119.52
2	B	701	77V	C22-C21-N20	2.01	122.24	118.06

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	77V	4	0
2	B	701	77V	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/291 (91%)	-1.48	0 100 100	35, 57, 88, 109	0
1	B	264/291 (90%)	-1.49	0 100 100	33, 57, 93, 124	0
All	All	530/582 (91%)	-1.49	0 100 100	33, 57, 91, 124	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	77V	A	701	24/24	0.99	0.04	55,60,62,62	0
2	77V	B	701	24/24	0.99	0.04	46,51,60,62	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 77V A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 77V B 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.