



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 01:27 AM UTC

PDB ID : 1T89 / pdb_00001t89
Title : CRYSTAL STRUCTURE OF A HUMAN TYPE III FC GAMMA RECEPTOR IN COMPLEX WITH AN FC FRAGMENT OF IGG1 (HEXAGONAL)
Authors : Radaev, S.; Motyka, S.; Fridman, W.-H.; Sautes-Fridman, C.; Sun, P.D.
Deposited on : 2004-05-11
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	224	28% 50% 13% • 6%
1	B	224	21% 52% 20% • 5%
2	C	176	33% 52% 9% • 5%
3	D	8	38% 12% 50%
4	E	8	12% 50% 38%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	1	X	-	-	-
3	MAN	D	4	-	-	X	-
4	NAG	E	1	X	-	-	-
4	FUC	E	8	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called recombinant IgG1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1677	1067	282	321	7			
1	B	213	Total	C	N	O	S	0	0	0
			1700	1083	285	325	7			

- Molecule 2 is a protein called Low affinity immunoglobulin gamma Fc region receptor III-B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	167	Total	C	N	O	S	101	0	0
			1347	856	231	256	4			

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[beta-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	8	Total	C	N	O	0	0	0
			96	54	3	39			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	8	Total	C	N	O	0	0	0
			99	56	4	39			

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: recombinant IgG1 heavy chain



- Molecule 1: recombinant IgG1 heavy chain




- Molecule 2: Low affinity immunoglobulin gamma Fc region receptor III-B





- Molecule 3: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[beta-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	114.98Å 114.98Å 301.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 3.50	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-3.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.247 , 0.309	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4919	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUL, NDG, MAN, BMA, GAL, FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/1723 (0.1%)	1.13	12/2346 (0.5%)
1	B	0.56	0/1747	1.04	10/2380 (0.4%)
2	C	0.60	0/1385	1.06	8/1884 (0.4%)
All	All	0.64	1/4855 (0.0%)	1.08	30/6610 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	296	TYR	CA-C	7.33	1.56	1.52

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	386	GLN	CA-C-N	9.01	131.10	119.84
1	B	386	GLN	C-N-CA	9.01	131.10	119.84
2	C	54	SER	N-CA-C	-8.91	101.67	112.54
2	C	59	ASP	N-CA-C	8.42	120.08	111.07
2	C	51	SER	N-CA-C	8.23	121.01	109.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1677	0	1644	220	0
1	B	1700	0	1668	232	0
2	C	1347	0	1288	120	0
3	D	96	0	81	9	0
4	E	99	0	84	11	0
All	All	4919	0	4765	561	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 59.

The worst 5 of 561 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:ARG:NE	1:A:293:GLU:H	1.49	1.10
1:A:292:ARG:HE	1:A:292:ARG:CA	1.63	1.10
1:A:292:ARG:HE	1:A:292:ARG:HA	0.98	1.09
1:B:308:VAL:HG12	1:B:309:LEU:H	1.13	1.08
1:A:350:THR:HB	1:A:441:LEU:HD12	1.39	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/224 (93%)	160 (77%)	24 (12%)	24 (12%)	0	5
1	B	211/224 (94%)	147 (70%)	35 (17%)	29 (14%)	0	3
2	C	165/176 (94%)	134 (81%)	20 (12%)	11 (7%)	1	11
All	All	584/624 (94%)	441 (76%)	79 (14%)	64 (11%)	0	5

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	ASP
1	A	287	ALA
1	A	291	PRO
1	A	408	SER
1	A	432	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/207 (94%)	173 (89%)	22 (11%)	5	25
1	B	197/207 (95%)	170 (86%)	27 (14%)	3	19
2	C	152/160 (95%)	142 (93%)	10 (7%)	15	41
All	All	544/574 (95%)	485 (89%)	59 (11%)	6	26

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	285	HIS
2	C	77	THR
1	B	337	SER
2	C	73	THR
2	C	15	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	315	ASN
2	C	115	ASN
1	B	386	GLN
2	C	140	HIS
2	C	74	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	D	1	3,1	14,14,15	0.59	0	17,19,21	1.59	4 (23%)
3	NAG	D	2	3	14,14,15	0.60	0	17,19,21	0.95	1 (5%)
3	BMA	D	3	3	11,11,12	0.69	0	15,15,17	0.61	0
3	MAN	D	4	3	11,11,12	0.78	0	15,15,17	0.90	1 (6%)
3	NDG	D	5	3	14,14,15	0.80	0	17,19,21	1.01	2 (11%)
3	GAL	D	6	3	11,11,12	0.66	0	15,15,17	0.49	0
3	BMA	D	7	3	11,11,12	0.49	0	15,15,17	0.80	0
3	FUL	D	8	3	10,10,11	0.66	0	14,14,16	0.63	0
4	NAG	E	1	4,1	14,14,15	1.18	2 (14%)	17,19,21	1.33	3 (17%)
4	NDG	E	2	4	14,14,15	0.69	0	17,19,21	1.07	1 (5%)
4	BMA	E	3	4	11,11,12	0.53	0	15,15,17	0.47	0
4	MAN	E	4	4	11,11,12	0.66	0	15,15,17	0.65	0
4	NAG	E	5	4	14,14,15	0.55	0	17,19,21	0.81	1 (5%)
4	MAN	E	6	4	11,11,12	0.58	0	15,15,17	0.51	0
4	NAG	E	7	4	14,14,15	0.82	0	17,19,21	1.58	3 (17%)
4	FUC	E	8	4	10,10,11	0.63	0	14,14,16	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	3,1	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	2/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	NDG	D	5	3	-	3/6/23/26	0/1/1/1
3	GAL	D	6	3	-	2/2/19/22	0/1/1/1
3	BMA	D	7	3	-	2/2/19/22	0/1/1/1
3	FUL	D	8	3	-	-	0/1/1/1
4	NAG	E	1	4,1	1/1/5/7	3/6/23/26	0/1/1/1
4	NDG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	0/2/19/22	0/1/1/1
4	MAN	E	4	4	-	2/2/19/22	0/1/1/1
4	NAG	E	5	4	-	2/6/23/26	0/1/1/1
4	MAN	E	6	4	-	2/2/19/22	0/1/1/1
4	NAG	E	7	4	-	4/6/23/26	0/1/1/1
4	FUC	E	8	4	1/1/4/5	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	1	NAG	C1-C2	2.85	1.56	1.52
4	E	1	NAG	O5-C5	2.05	1.47	1.43

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	7	NAG	C4-C3-C2	-4.19	104.88	111.02
3	D	1	NAG	O5-C1-C2	4.03	117.53	111.29
4	E	7	NAG	C2-N2-C7	-3.53	118.17	122.90
4	E	2	NDG	C2-N2-C7	-3.09	118.76	122.90
4	E	1	NAG	C4-C3-C2	-2.85	106.85	111.02

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	1	NAG	C1
4	E	1	NAG	C1
4	E	8	FUC	C1

5 of 30 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
3	D	2	NAG	C8-C7-N2-C2
3	D	2	NAG	O7-C7-N2-C2
3	D	5	NDG	C3-C2-N2-C7


There are no ring outliers.

11 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	6	MAN	3	0
4	E	7	NAG	2	0
3	D	3	BMA	1	0
3	D	4	MAN	6	0
3	D	1	NAG	2	0
4	E	2	NDG	5	0
4	E	8	FUC	2	0
3	D	2	NAG	2	0
3	D	5	NDG	2	0
4	E	3	BMA	2	0
4	E	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.