



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 05:51 PM UTC

PDB ID : 3T8L / pdb_00003t8l
Title : Crystal Structure of adenine deaminase with Mn/Fe
Authors : Bagaria, A.; Kumaran, D.; Burley, S.K.; Swaminathan, S.; New York SGX
Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2011-08-01
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	608	
1	B	608	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	606	-	-	-	X
2	UNX	A	607	-	-	-	X
2	UNX	A	608	-	-	-	X
2	UNX	B	607	-	-	-	X
2	UNX	B	608	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenine deaminase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	587	Total	C	N	O	S	Se	0	0	0
			4337	2723	775	817	6	16			
1	B	587	Total	C	N	O	S	Se	0	0	0
			4335	2721	775	817	6	16			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MSE	-	expression tag	UNP Q7CUX4
A	-1	SER	-	expression tag	UNP Q7CUX4
A	0	LEU	-	expression tag	UNP Q7CUX4
A	598	GLU	-	expression tag	UNP Q7CUX4
A	599	GLY	-	expression tag	UNP Q7CUX4
A	600	HIS	-	expression tag	UNP Q7CUX4
A	601	HIS	-	expression tag	UNP Q7CUX4
A	602	HIS	-	expression tag	UNP Q7CUX4
A	603	HIS	-	expression tag	UNP Q7CUX4
A	604	HIS	-	expression tag	UNP Q7CUX4
A	605	HIS	-	expression tag	UNP Q7CUX4
B	-2	MSE	-	expression tag	UNP Q7CUX4
B	-1	SER	-	expression tag	UNP Q7CUX4
B	0	LEU	-	expression tag	UNP Q7CUX4
B	598	GLU	-	expression tag	UNP Q7CUX4
B	599	GLY	-	expression tag	UNP Q7CUX4
B	600	HIS	-	expression tag	UNP Q7CUX4
B	601	HIS	-	expression tag	UNP Q7CUX4
B	602	HIS	-	expression tag	UNP Q7CUX4
B	603	HIS	-	expression tag	UNP Q7CUX4
B	604	HIS	-	expression tag	UNP Q7CUX4
B	605	HIS	-	expression tag	UNP Q7CUX4

- Molecule 2 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	X 3	0	0
2	B	3	Total 3	X 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total 33	O 33	0	0
3	B	50	Total 50	O 50	0	0

[illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.54Å 131.17Å 69.28Å 90.00° 97.04° 90.00°	Depositor
Resolution (Å)	48.73 – 2.80 48.73 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.73-2.80) 99.7 (48.73-2.80)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.301 0.182 , 0.304	Depositor DCC
R_{free} test set	1362 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	56.2	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8761	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.66	265/4402 (6.0%)	2.19	191/5964 (3.2%)
1	B	2.53	231/4400 (5.2%)	2.17	191/5961 (3.2%)
All	All	2.59	496/8802 (5.6%)	2.18	382/11925 (3.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	6
All	All	0	8

The worst 5 of 496 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	451	MSE	SE-CE	24.10	2.67	1.95
1	A	396	MSE	SE-CE	23.09	2.64	1.95
1	B	189	MSE	SE-CE	22.53	2.63	1.95
1	A	189	MSE	SE-CE	22.34	2.62	1.95
1	B	191	MSE	SE-CE	19.62	2.54	1.95

The worst 5 of 382 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	548	VAL	N-CA-C	-13.68	99.65	113.47
1	B	420	THR	N-CA-C	13.13	130.73	109.72
1	A	210	ALA	N-CA-C	-13.10	97.08	111.36
1	A	502	THR	N-CA-C	-12.08	98.52	113.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	552	GLN	CA-C-N	11.04	131.75	120.38

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	396	MSE	Peptide
1	A	430	LYS	Peptide
1	B	152	PRO	Peptide
1	B	412	ALA	Peptide
1	B	417	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4337	0	4344	237	0
1	B	4335	0	4339	315	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	33	0	0	5	0
3	B	50	0	0	16	0
All	All	8761	0	8683	539	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 539 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ($\text{\AA}$)	Clash overlap ($\text{\AA}$)
1:A:385:VAL:CB	1:A:385:VAL:CG1	1.76	1.58
1:B:231:MSE:SE	1:B:231:MSE:CG	2.14	1.44
1:A:492:MSE:SE	1:A:492:MSE:CE	2.14	1.44
1:A:393:PRO:CB	1:A:393:PRO:CG	1.79	1.43
1:A:231:MSE:SE	1:A:231:MSE:CE	2.17	1.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/608 (96%)	520 (89%)	49 (8%)	16 (3%)	4	15
1	B	585/608 (96%)	483 (83%)	69 (12%)	33 (6%)	1	4
All	All	1170/1216 (96%)	1003 (86%)	118 (10%)	49 (4%)	2	7

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	ASN
1	A	406	GLY
1	B	72	ASP
1	B	224	ASN
1	B	418	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/444 (100%)	371 (83%)	74 (17%)	2	8
1	B	444/444 (100%)	370 (83%)	74 (17%)	2	8
All	All	889/888 (100%)	741 (83%)	148 (17%)	2	8

5 of 148 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	409	VAL
1	B	570	ILE
1	B	420	THR
1	B	474	THR
1	A	435	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	65	HIS
1	B	228	ASN
1	B	552	GLN
1	B	128	HIS
1	B	265	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	571/608 (93%)	-0.64	0 100 100	28, 44, 66, 80	0
1	B	571/608 (93%)	-0.27	26 (4%) 37 29	23, 47, 83, 98	27 (4%)
All	All	1142/1216 (93%)	-0.46	26 (2%) 61 51	23, 46, 75, 98	27 (2%)

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	419	PHE	7.9
1	B	418	ARG	6.1
1	B	427	ALA	5.7
1	B	417	PRO	5.4
1	B	593	ILE	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UNX	A	606	1/1	0.39	0.55	53,53,53,53	0
2	UNX	B	607	1/1	0.47	0.41	60,60,60,60	0
2	UNX	A	608	1/1	0.52	0.52	43,43,43,43	0
2	UNX	A	607	1/1	0.54	0.49	43,43,43,43	0
2	UNX	B	608	1/1	0.80	0.42	50,50,50,50	0
2	UNX	B	606	1/1	0.84	0.42	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.