



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:53 PM UTC

PDB ID : 5T8I / pdb_00005t8i
Title : PI3Kdelta in complex with the inhibitor GS-9901
Authors : Somoza, J.R.; Villasenor, A.
Deposited on : 2016-09-07
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	939	5% 78% 7% 14%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 13091 atoms, of which 6529 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	808	13040	4183	6512	1108	1182	55	0	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	GLN	-	insertion	UNP O35904

- Molecule 2 is 2,4-diamino-6-[[[(S)-[5-chloro-8-fluoro-4-oxo-3-(pyridin-3-yl)-3,4-dihydroquinazolin-2-yl](cyclopropyl)methyl]amino}pyrimidine-5-carbonitrile (CCD ID: 77C) (formula: C₂₂H₁₇ClFN₉O).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	Cl	F	H	N	O		
2	A	1	51	22	1	1	17	9	1	0	0

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.05Å 142.20Å 222.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.88 – 2.60 45.88 – 2.60	Depositor EDS
% Data completeness (in resolution range)	91.2 (45.88-2.60) 91.2 (45.88-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.252 , 0.312 0.258 , 0.316	Depositor DCC
R_{free} test set	1999 reflections (6.33%)	wwPDB-VP
Wilson B-factor (Å ²)	70.0	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13091	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 77C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/6693	0.65	1/9029 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	714	MET	CG-SD-CE	8.46	119.51	100.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6528	6512	6492	32	1
2	A	34	17	0	0	0
All	All	6562	6529	6492	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ASP:O	1:A:132:ARG:NH1	2.25	0.70
1:A:367:SER:HA	1:A:368:GLU:HB2	1.83	0.60
1:A:1006:LYS:NZ	1:A:1007:THR:O	2.37	0.58
1:A:897:ASP:N	1:A:897:ASP:OD1	2.37	0.57
1:A:1016:ARG:O	1:A:1020:ASN:ND2	2.38	0.57
1:A:982:ARG:NH1	1:A:991:CYS:O	2.36	0.57
1:A:246:ARG:NH1	1:A:248:GLU:OE1	2.40	0.55
1:A:512:ARG:O	1:A:514:ILE:N	2.41	0.54
1:A:532:TRP:C	1:A:533:LYS:HE2	2.34	0.53
1:A:383:ASP:OD1	1:A:558:HIS:ND1	2.42	0.53
1:A:533:LYS:HZ2	1:A:538:VAL:CG2	2.23	0.50
1:A:750:THR:OG1	1:A:751:PHE:N	2.45	0.50
1:A:533:LYS:O	1:A:535:ARG:N	2.46	0.49
1:A:982:ARG:NH2	1:A:990:SER:O	2.46	0.49
1:A:852:LYS:NZ	1:A:853:ASP:OD2	2.45	0.49
1:A:511:LEU:O	1:A:513:GLU:N	2.47	0.47
1:A:844:MET:O	1:A:847:THR:N	2.48	0.47
1:A:528:LYS:HA	1:A:531:VAL:HG12	1.96	0.46
1:A:533:LYS:HZ2	1:A:538:VAL:HG23	1.80	0.46
1:A:533:LYS:HD2	1:A:534:MET:N	2.31	0.46
1:A:944:GLN:NE2	1:A:949:ASN:OD1	2.48	0.46
1:A:367:SER:CA	1:A:368:GLU:HB2	2.47	0.45
1:A:1021:GLU:O	1:A:1024:ARG:NH1	2.49	0.44
1:A:531:VAL:HG13	1:A:552:VAL:HG11	1.98	0.44
1:A:533:LYS:HE3	1:A:534:MET:C	2.43	0.44
1:A:511:LEU:HG	1:A:512:ARG:N	2.33	0.43
1:A:343:VAL:HG22	1:A:394:LEU:HD12	2.01	0.42
1:A:765:SER:HB3	1:A:772:GLY:HA3	2.02	0.42
1:A:528:LYS:HA	1:A:531:VAL:CG1	2.49	0.41
1:A:531:VAL:C	1:A:533:LYS:HE2	2.45	0.41
1:A:996:GLN:NE2	1:A:1000:ASP:OD1	2.49	0.41
1:A:157:LEU:O	1:A:286:ARG:NH1	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:714:MET:SD	1:A:714:MET:CE[4_566]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	792/939 (84%)	734 (93%)	46 (6%)	12 (2%)	8	18

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	512	ARG
1	A	513	GLU
1	A	533	LYS
1	A	534	MET
1	A	319	LEU
1	A	767	GLU
1	A	865	GLY
1	A	229	ARG
1	A	727	ALA
1	A	227	VAL
1	A	328	ILE
1	A	766	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	721/827 (87%)	697 (97%)	24 (3%)	33	61

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	149	GLU
1	A	187	ASN
1	A	247	HIS
1	A	319	LEU
1	A	331	ARG
1	A	334	ASN
1	A	433	LYS
1	A	473	ILE
1	A	492	LEU
1	A	494	LEU
1	A	511	LEU
1	A	533	LYS
1	A	594	SER
1	A	706	THR
1	A	743	GLU
1	A	770	SER
1	A	792	GLN
1	A	852	LYS
1	A	897	ASP
1	A	901	ILE
1	A	915	PHE
1	A	951	GLU
1	A	1010	GLU
1	A	1013	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	170	GLN
1	A	334	ASN
1	A	840	ASN
1	A	906	GLN
1	A	976	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	77C	A	1101	-	38,38,38	1.10	5 (13%)	43,56,56	1.29	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	77C	A	1101	-	-	2/12/20/20	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	77C	C21-N14	3.24	1.40	1.35
2	A	1101	77C	C25-N27	2.30	1.38	1.33
2	A	1101	77C	C22-C23	-2.18	1.39	1.42
2	A	1101	77C	C23-N28	2.09	1.39	1.34
2	A	1101	77C	C13-C08	2.06	1.54	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	77C	C05-C06-CL3	4.12	124.93	119.62
2	A	1101	77C	N09-C08-N07	-2.85	121.75	124.21
2	A	1101	77C	C15-C13-C08	2.82	116.40	111.61
2	A	1101	77C	C21-C22-C29	2.08	122.01	119.21

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	77C	C08-C13-C15-C33
2	A	1101	77C	C08-C13-C15-C34

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	808/939 (86%)	0.72	50 (6%) 26 21	43, 97, 115, 124	3 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	511	LEU	5.7
1	A	335	ALA	4.8
1	A	109	VAL	3.9
1	A	792	GLN	3.7
1	A	452	LEU	3.6
1	A	530	LEU	3.5
1	A	332	LYS	3.4
1	A	316	LEU	3.3
1	A	173	PRO	3.1
1	A	474	TYR	3.0
1	A	365	VAL	2.9
1	A	479	ALA	2.8
1	A	377	PHE	2.8
1	A	842	SER	2.7
1	A	482	PRO	2.7
1	A	568	LEU	2.6
1	A	936	TYR	2.6
1	A	839	LEU	2.6
1	A	445	VAL	2.6
1	A	904	SER	2.6
1	A	330	GLY	2.5
1	A	226	THR	2.5
1	A	492	LEU	2.5
1	A	366	CYS	2.5
1	A	240	ALA	2.5
1	A	334	ASN	2.4
1	A	136	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	437	ARG	2.4
1	A	512	ARG	2.4
1	A	637	ALA	2.4
1	A	494	LEU	2.4
1	A	677	HIS	2.3
1	A	516	GLU	2.3
1	A	861	SER	2.3
1	A	150	ALA	2.3
1	A	1007	THR	2.3
1	A	484	TYR	2.3
1	A	388	ALA	2.2
1	A	835	ALA	2.2
1	A	847	THR	2.2
1	A	846	ALA	2.2
1	A	398	VAL	2.2
1	A	347	LEU	2.1
1	A	766	GLU	2.1
1	A	358	VAL	2.1
1	A	352	GLU	2.1
1	A	397	VAL	2.1
1	A	196	PHE	2.0
1	A	434	THR	2.0
1	A	1012	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	77C	A	1101	34/34	0.93	0.10	73,79,96,99	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.