



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:05 AM UTC

PDB ID : 7T87 / pdb_00007t87
Title : CRYSTAL STRUCTURE OF LEUKOCIDIN AB/CENTYRIN S17/FAB
214F COMPLEX
Authors : Luo, J.; Malia, T.J.; Buckley, P.T.
Deposited on : 2021-12-15
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 77% 16% • 6%
2	B	332	 68% 17% • 15%
3	L	213	 92% 7%
4	H	233	 84% 10% 6%
5	C	98	 82% 14% •

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leukocidin B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2346	1470	411	460	5			

- Molecule 2 is a protein called Leukocidin A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	283	Total	C	N	O	0	0	0
			2291	1447	398	446			

- Molecule 3 is a protein called Antibody Fab B214 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1624	1018	272	327	7			

- Molecule 4 is a protein called Antibody Fab B214 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	220	Total	C	N	O	S	0	0	0
			1654	1049	268	330	7			

- Molecule 5 is a protein called Centyrin S17.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	C	94	Total	C	N	O	S	0	0	0
			726	471	119	135	1			

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Leukocidin B

Chain A: 




• Molecule 2: Leukocidin A

Chain B: 



• Molecule 3: Antibody Fab B214 Light Chain

Chain L: 





• Molecule 4: Antibody Fab B214 Heavy Chain

Chain H: 



HIS
HIS
HIS
HIS
HIS
HIS

● Molecule 5: Centyrin S17



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.98Å 173.74Å 174.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.16 – 3.00 49.16 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.0 (49.16-3.00) 95.0 (49.16-3.00)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.196 , 0.247 0.197 , 0.248	Depositor DCC
R_{free} test set	1361 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	63.8	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8641	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/2404	0.32	0/3253
2	B	0.14	0/2343	0.32	0/3172
3	L	0.13	0/1660	0.30	0/2254
4	H	0.13	0/1697	0.31	0/2308
5	C	0.15	0/749	0.36	0/1023
All	All	0.14	0/8853	0.32	0/12010

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2346	0	2219	22	0
2	B	2291	0	2184	28	0
3	L	1624	0	1583	6	0
4	H	1654	0	1590	13	0
5	C	726	0	701	9	0
All	All	8641	0	8277	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:302:ASP:OD2	2:B:305:ASN:ND2	2.19	0.76
5:C:4:ALA:HB2	5:C:82:TRP:HB2	1.68	0.74
2:B:181:SER:O	2:B:184:ASN:ND2	2.20	0.74
1:A:38:LEU:HD21	1:A:249:VAL:HG21	1.73	0.70
1:A:85:SER:HB2	1:A:252:LYS:HG3	1.75	0.68
4:H:12:VAL:HG11	4:H:18:VAL:HB	1.75	0.68
1:A:264:ARG:NH2	5:C:37:GLU:OE2	2.26	0.66
2:B:70:ASP:OD2	2:B:306:LYS:NZ	2.20	0.66
1:A:26:THR:HG22	1:A:37:SER:HA	1.77	0.65
2:B:122:LYS:HD3	2:B:257:ASN:HB2	1.79	0.65
1:A:264:ARG:HG2	5:C:35:THR:HG21	1.77	0.64
1:A:89:GLN:HG3	1:A:165:GLY:HA3	1.80	0.64
2:B:112:HIS:HB2	2:B:267:THR:HB	1.81	0.63
5:C:72:ILE:HG23	5:C:84:LEU:HB3	1.80	0.62
1:A:45:GLU:OE1	2:B:48:LYS:NZ	2.33	0.61
1:A:183:THR:HG23	1:A:185:GLN:H	1.64	0.61
3:L:146:LYS:HB3	3:L:198:THR:HB	1.84	0.58
1:A:41:ASN:HB2	1:A:54:PHE:HB2	1.86	0.58
2:B:110:GLU:HB2	2:B:269:ASN:HB2	1.85	0.57
5:C:29:PHE:HA	5:C:76:LYS:HB2	1.85	0.57
1:A:178:MET:HE1	1:A:203:ARG:HB2	1.87	0.57
1:A:290:VAL:HG22	1:A:297:VAL:HG22	1.87	0.56
2:B:232:LYS:HD2	2:B:240:ARG:NH1	2.20	0.56
1:A:18:THR:HG22	1:A:46:PRO:HD3	1.88	0.56
2:B:294:ARG:HH21	2:B:319:LYS:HG3	1.72	0.55
3:L:121:PRO:HD3	3:L:133:VAL:HG22	1.89	0.55
4:H:179:ALA:HB2	4:H:189:LEU:HD23	1.89	0.54
4:H:134:PRO:HD3	4:H:220:LYS:HE2	1.90	0.54
5:C:34:ILE:HG12	5:C:72:ILE:HD13	1.90	0.54
2:B:82:LYS:HD2	2:B:84:GLN:HE21	1.72	0.54
3:L:6:GLN:HE22	3:L:88:PHE:HA	1.74	0.53
2:B:274:LYS:HB2	2:B:285:ILE:HB	1.89	0.53
2:B:106:LYS:HG2	2:B:272:ILE:HG12	1.91	0.52
2:B:222:VAL:HG22	2:B:277:PRO:HG2	1.92	0.52
1:A:132:LEU:HD12	2:B:164:SER:HA	1.92	0.52
1:A:163:HIS:CD2	1:A:164:LYS:HG3	2.45	0.51
2:B:291:ASP:OD1	2:B:292:GLY:N	2.41	0.50
3:L:194:ALA:HB2	3:L:209:SER:HB3	1.94	0.49
2:B:114:ASP:HB3	2:B:265:THR:HB	1.94	0.48
1:A:281:GLU:HB2	4:H:107:TYR:CE1	2.49	0.48
1:A:63:SER:HB2	1:A:81:PRO:HG3	1.96	0.47
5:C:25:PRO:HB2	5:C:28:ALA:HB2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:126:LEU:HD22	2:B:252:SER:HB3	1.97	0.47
2:B:218:ARG:HG2	2:B:279:ILE:HA	1.96	0.47
2:B:79:LEU:HD23	2:B:251:LEU:HD12	1.95	0.47
2:B:88:HIS:O	2:B:268:ARG:NH1	2.46	0.47
4:H:67:LYS:NZ	4:H:85:SER:O	2.47	0.47
1:A:280:LYS:NZ	4:H:55:ASP:OD2	2.45	0.46
4:H:46:GLU:OE2	4:H:63:LYS:NZ	2.46	0.46
2:B:111:TYR:HB2	2:B:193:VAL:HB	1.97	0.46
4:H:154:LYS:NZ	4:H:182:GLN:OE1	2.49	0.46
1:A:238:ASP:CG	1:A:240:LYS:HG2	2.41	0.45
2:B:47:THR:HB	2:B:68:PHE:HB2	1.98	0.45
1:A:68:LEU:HD21	1:A:79:ARG:HB2	1.99	0.45
1:A:28:SER:HA	1:A:35:THR:HA	1.98	0.44
1:A:128:ASN:OD1	1:A:130:GLY:N	2.31	0.44
2:B:195:ALA:HB1	2:B:198:LEU:HD21	1.99	0.44
4:H:130:PRO:HB3	4:H:156:TYR:HB3	1.98	0.44
2:B:294:ARG:NH1	2:B:296:ILE:HD11	2.33	0.44
4:H:12:VAL:HG21	4:H:86:LEU:HD13	2.00	0.43
2:B:110:GLU:HG2	2:B:194:ILE:HG23	2.01	0.43
4:H:179:ALA:HA	4:H:189:LEU:HB3	1.99	0.43
2:B:142:THR:HG22	2:B:168:THR:HG22	2.00	0.43
5:C:32:PHE:HB2	5:C:50:VAL:HG22	1.99	0.43
5:C:32:PHE:HB3	5:C:72:ILE:HD11	2.00	0.43
4:H:36:TRP:CD1	4:H:70:LEU:HD22	2.53	0.42
2:B:296:ILE:HB	2:B:314:TYR:HB3	2.01	0.42
2:B:66:PHE:CE1	2:B:308:VAL:HG11	2.54	0.42
3:L:109:ARG:HG2	3:L:172:SER:HB2	2.01	0.42
4:H:10:GLU:HB2	4:H:120:VAL:HG22	2.02	0.42
3:L:141:TYR:CD1	3:L:142:PRO:HA	2.54	0.42
1:A:20:MET:HE2	1:A:43:LEU:HD13	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/309 (93%)	270 (94%)	18 (6%)	0	100	100
2	B	281/332 (85%)	262 (93%)	19 (7%)	0	100	100
3	L	211/213 (99%)	196 (93%)	14 (7%)	1 (0%)	24	60
4	H	216/233 (93%)	210 (97%)	6 (3%)	0	100	100
5	C	92/98 (94%)	86 (94%)	5 (5%)	1 (1%)	11	43
All	All	1088/1185 (92%)	1024 (94%)	62 (6%)	2 (0%)	43	76

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	C	3	PRO
3	L	139	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/281 (92%)	242 (94%)	16 (6%)	16	49
2	B	248/308 (80%)	238 (96%)	10 (4%)	28	62
3	L	185/185 (100%)	181 (98%)	4 (2%)	45	74
4	H	181/195 (93%)	178 (98%)	3 (2%)	53	78
5	C	75/81 (93%)	75 (100%)	0	100	100
All	All	947/1050 (90%)	914 (96%)	33 (4%)	32	65

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	THR
1	A	66	ARG
1	A	69	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	109	GLU
1	A	116	THR
1	A	136	ILE
1	A	149	GLN
1	A	158	ASP
1	A	204	ASN
1	A	207	LEU
1	A	210	LYS
1	A	216	LYS
1	A	225	GLU
1	A	264	ARG
1	A	279	LYS
1	A	305	ASP
2	B	48	LYS
2	B	98	GLU
2	B	99	GLU
2	B	117	VAL
2	B	139	VAL
2	B	169	ILE
2	B	279	ILE
2	B	301	VAL
2	B	304	LYS
2	B	319	LYS
3	L	10	THR
3	L	19	ILE
3	L	51	ARG
3	L	109	ARG
4	H	43	LYS
4	H	98	ARG
4	H	111	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	ASN
1	A	96	ASN
1	A	107	GLN
1	A	192	ASN
1	A	276	HIS
2	B	77	ASN
2	B	120	ASN
2	B	174	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	216	ASN
2	B	253	ASN
2	B	257	ASN
2	B	318	ASN
3	L	38	GLN
3	L	125	GLN
3	L	139	ASN
4	H	5	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	290/309 (93%)	-0.42	0	100	100	24, 47, 81, 101	0
2	B	283/332 (85%)	-0.12	0	100	100	32, 59, 91, 106	0
3	L	213/213 (100%)	0.34	1 (0%)	87	72	46, 81, 122, 129	0
4	H	220/233 (94%)	-0.17	2 (0%)	81	61	33, 64, 98, 135	0
5	C	94/98 (95%)	0.17	0	100	100	39, 75, 94, 111	0
All	All	1100/1185 (92%)	-0.09	3 (0%)	90	79	24, 62, 106, 135	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	143	SER	4.0
3	L	197	VAL	2.1
4	H	138	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.