



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2026 – 11:33 AM UTC

PDB ID : 1T9G / pdb_00001t9g
Title : Structure of the human MCAD:ETF complex
Authors : Toogood, H.S.; van Thiel, A.; Basran, J.; Sutcliffe, M.J.; Scrutton, N.S.; Leys, D.
Deposited on : 2004-05-17
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 63% 28% 7% .
1	B	396	 58% 33% 6% ..
1	C	396	 61% 29% 7% ..
1	D	396	 59% 32% 7% ..
2	R	333	 35% 17% . 45%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	S	255	 A horizontal bar chart showing the quality of chain S. The bar is divided into four segments: green (50%), yellow (33%), orange (5%), and grey (11%). The percentages are labeled below the bar.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA dehydrogenase, medium-chain specific, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			2929	1859	495	557	18			
1	B	387	Total	C	N	O	S	0	0	0
			2930	1857	496	559	18			
1	C	388	Total	C	N	O	S	0	0	0
			2951	1870	507	556	18			
1	D	387	Total	C	N	O	S	0	0	0
			2957	1872	507	560	18			


- Molecule 2 is a protein called Electron transfer flavoprotein alpha-subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	R	184	Total	C	N	O	S	0	0	0
			1323	835	221	261	6			

- Molecule 3 is a protein called Electron transfer flavoprotein beta-subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	S	228	Total	C	N	O	S	0	0	0
			1673	1062	282	321	8			

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
4	B	1	Total 53	C 27	N 9	O 15	P 2	0	0
4	C	1	Total 53	C 27	N 9	O 15	P 2	0	0
4	D	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 5 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: $\text{C}_{10}\text{H}_{14}\text{N}_5\text{O}_7\text{P}$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	S	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acyl-CoA dehydrogenase, medium-chain specific, mitochondrial




- Molecule 1: Acyl-CoA dehydrogenase, medium-chain specific, mitochondrial




- Molecule 1: Acyl-CoA dehydrogenase, medium-chain specific, mitochondrial



- Molecule 1: Acyl-CoA dehydrogenase, medium-chain specific, mitochondrial




- Molecule 2: Electron transfer flavoprotein alpha-subunit, mitochondrial



ALA
ASP
LEU
PHE
LYS
VAL
VAL
PRO
GLU
MET
THR
GLU
ILE
LEU
LYS
LYS

● Molecule 3: Electron transfer flavoprotein beta-subunit

Chain S: 50% 33% 5% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.33Å 101.32Å 244.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.90 20.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.90) 97.9 (20.00-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 2.88Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.191 , 0.263 0.191 , 0.264	Depositor DCC
R_{free} test set	2614 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	52.6	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 29.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14998	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	6/2984 (0.2%)	1.30	14/4036 (0.3%)
1	B	1.17	3/2985 (0.1%)	1.30	8/4039 (0.2%)
1	C	1.19	3/3006 (0.1%)	1.37	20/4062 (0.5%)
1	D	1.20	5/3012 (0.2%)	1.34	15/4067 (0.4%)
2	R	1.02	1/1342 (0.1%)	1.18	4/1832 (0.2%)
3	S	1.07	1/1694 (0.1%)	1.31	10/2306 (0.4%)
All	All	1.17	19/15023 (0.1%)	1.31	71/20342 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
3	S	0	3
All	All	0	4

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	36	ILE	CA-CB	-7.62	1.50	1.54
1	A	36	ILE	CA-CB	7.49	1.57	1.54
1	A	110	ILE	CA-CB	6.90	1.63	1.54
1	A	226	VAL	CA-CB	6.50	1.63	1.54
1	D	381	ILE	CA-CB	6.35	1.63	1.54

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	GLY	N-CA-C	-8.71	101.17	111.36
1	D	193	ALA	N-CA-C	7.76	122.01	112.54
3	S	57	LEU	N-CA-C	-7.68	102.97	114.64
1	A	238	LEU	CA-C-N	-7.51	114.89	122.77
1	A	238	LEU	C-N-CA	-7.51	114.89	122.77

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	324	ARG	Peptide
3	S	176	LYS	Peptide
3	S	177	LEU	Peptide
3	S	230	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2929	0	2844	100	0
1	B	2930	0	2826	110	0
1	C	2951	0	2888	103	0
1	D	2957	0	2895	110	0
2	R	1323	0	1320	53	0
3	S	1673	0	1694	95	0
4	A	53	0	31	10	0
4	B	53	0	31	6	0
4	C	53	0	31	2	0
4	D	53	0	31	7	0
5	S	23	0	12	4	0
All	All	14998	0	14603	531	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:399:FAD:N10	4:A:399:FAD:C1'	1.78	1.47
1:A:124:MET:HE3	1:A:175:TRP:HE1	1.07	1.11
1:A:124:MET:CE	1:A:175:TRP:HE1	1.63	1.11
1:D:73:LEU:HG	3:S:199:MET:HE1	1.11	1.10
1:A:363:GLU:HG3	1:B:215:MET:HB2	1.24	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/396 (97%)	352 (92%)	27 (7%)	5 (1%)	9	32
1	B	385/396 (97%)	351 (91%)	30 (8%)	4 (1%)	12	39
1	C	386/396 (98%)	362 (94%)	20 (5%)	4 (1%)	12	39
1	D	385/396 (97%)	361 (94%)	19 (5%)	5 (1%)	9	32
2	R	182/333 (55%)	165 (91%)	11 (6%)	6 (3%)	3	13
3	S	226/255 (89%)	197 (87%)	25 (11%)	4 (2%)	6	25
All	All	1948/2172 (90%)	1788 (92%)	132 (7%)	28 (1%)	9	30

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	234	LYS
1	D	156	ASP
1	D	234	LYS
2	R	112	ALA
3	S	25	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/311 (92%)	244 (85%)	42 (15%)	3	10
1	B	284/311 (91%)	240 (84%)	44 (16%)	2	9
1	C	290/311 (93%)	245 (84%)	45 (16%)	2	9
1	D	292/311 (94%)	241 (82%)	51 (18%)	2	6
2	R	139/262 (53%)	124 (89%)	15 (11%)	6	21
3	S	174/214 (81%)	150 (86%)	24 (14%)	3	12
All	All	1465/1720 (85%)	1244 (85%)	221 (15%)	3	10

5 of 221 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	238	LEU
1	D	98	ILE
3	S	222	LEU
3	S	14	ILE
1	C	259	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	373	GLN
1	D	342	GLN
1	C	382	GLN
1	D	174	ASN
1	D	382	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FAD	B	399	-	58,58,58	1.21	5 (8%)	85,89,89	2.09	19 (22%)
4	FAD	A	399	-	58,58,58	2.64	13 (22%)	85,89,89	2.46	31 (36%)
4	FAD	D	399	-	58,58,58	1.06	4 (6%)	85,89,89	1.74	22 (25%)
5	AMP	S	600	-	25,25,25	1.79	4 (16%)	37,38,38	1.78	9 (24%)
4	FAD	C	399	-	58,58,58	1.60	11 (18%)	85,89,89	2.23	30 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	B	399	-	-	1/34/50/50	0/6/6/6
4	FAD	A	399	-	-	6/34/50/50	0/6/6/6
4	FAD	D	399	-	-	0/34/50/50	0/6/6/6
5	AMP	S	600	-	-	5/10/26/26	0/3/3/3
4	FAD	C	399	-	-	4/34/50/50	0/6/6/6

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	399	FAD	C1'-N10	12.60	1.78	1.47
4	A	399	FAD	C1'-C2'	7.27	1.62	1.52
4	A	399	FAD	C8M-C8	-7.22	1.37	1.51
5	S	600	AMP	C5-C4	6.05	1.49	1.39
4	A	399	FAD	C4X-N5	4.79	1.41	1.30

The worst 5 of 111 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	399	FAD	O4-C4-C4X	-7.98	105.47	126.53
4	A	399	FAD	O4-C4-N3	7.59	134.39	120.11
4	C	399	FAD	C8M-C8-C9	-7.00	107.24	119.57
4	C	399	FAD	C5A-C4A-N3A	-6.21	118.17	126.72
4	C	399	FAD	O5'-C5'-C4'	-6.11	93.04	109.36

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	399	FAD	C2'-C1'-N10-C9A
4	A	399	FAD	C2'-C1'-N10-C10
4	A	399	FAD	N10-C1'-C2'-O2'
4	A	399	FAD	N10-C1'-C2'-C3'
4	A	399	FAD	C1'-C2'-C3'-C4'


There are no ring outliers.


5 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	399	FAD	6	0
4	A	399	FAD	10	0
4	D	399	FAD	7	0
5	S	600	AMP	4	0
4	C	399	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand FAD D 399



Ligand AMP S 600





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/396 (97%)	-0.73	0 100 100	21, 37, 53, 71	0
1	B	387/396 (97%)	-0.59	0 100 100	23, 43, 63, 81	0
1	C	388/396 (97%)	-0.71	0 100 100	20, 36, 53, 70	0
1	D	387/396 (97%)	-0.68	0 100 100	20, 36, 56, 68	0
2	R	184/333 (55%)	-0.06	0 100 100	43, 62, 74, 79	0
3	S	228/255 (89%)	-0.14	0 100 100	36, 57, 85, 90	0
All	All	1960/2172 (90%)	-0.56	0 100 100	20, 41, 71, 90	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	AMP	S	600	23/23	0.95	0.07	37,42,47,51	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FAD	C	399	53/53	0.96	0.08	24,29,37,39	0
4	FAD	A	399	53/53	0.96	0.07	19,28,34,36	0
4	FAD	D	399	53/53	0.97	0.06	24,31,34,35	0
4	FAD	B	399	53/53	0.97	0.05	25,32,37,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AMP S 600:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 399:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 399:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD D 399:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 399:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.