



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 07:08 PM UTC

PDB ID : 6T9K / pdb_00006t9k
EMDB ID : EMD-10414
Title : SAGA Core module
Authors : Wang, H.; Cheung, A.; Cramer, P.
Deposited on : 2019-10-28
Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15087 (2.80 - 3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	604	
2	C	337	
3	D	798	
4	E	516	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	157	
6	G	206	
7	H	488	
8	I	539	
9	K	1332	
10	Q	657	
11	U	155	

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 19241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription factor SPT20.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	181	Total	C	N	O	S	0	0
			1526	975	254	291	6		

- Molecule 2 is a protein called Protein SPT3.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	214	Total	C	N	O	S	0	0
			1774	1130	319	318	7		

- Molecule 3 is a protein called Transcription initiation factor TFIID subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	566	Total	C	N	O	S	0	0
			4474	2830	772	854	18		

- Molecule 4 is a protein called Transcription initiation factor TFIID subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	433	Total	C	N	O	S	0	0
			3441	2207	572	657	5		

- Molecule 5 is a protein called Transcription initiation factor TFIID subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	F	111	Total	C	N	O	S	0	0
			902	581	158	159	4		

- Molecule 6 is a protein called Transcription initiation factor TFIID subunit 10.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	G	140	Total	C	N	O	S	0	0
			1108	689	199	218	2		

- Molecule 7 is a protein called Transcriptional coactivator HFI1/ADA1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	H	227	Total	C	N	O	S	0	0
			1803	1121	314	359	9		

- Molecule 8 is a protein called Transcription initiation factor TFIID subunit 12.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	I	108	Total	C	N	O	S	0	0
			865	533	159	172	1		

- Molecule 9 is a protein called Transcriptional activator SPT7.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	K	279	Total	C	N	O	S	0	0
			2258	1427	392	428	11		

- Molecule 10 is a protein called SAGA-associated factor 73.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	Q	84	Total	C	N	O	S	0	0
			675	426	119	127	3		


- Molecule 11 is a protein called unassigned sequence.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	U	83	Total	C	N	O	0	0
			415	249	83	83		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcription factor SPT20




• Molecule 2: Protein SPT3









- Molecule 5: Transcription initiation factor TFIID subunit 9



- Molecule 6: Transcription initiation factor TFIID subunit 10



- Molecule 7: Transcriptional coactivator HFI1/ADA1



NET ARG SER SER ASP GLY ALA GLU ILE LYS GLY ILE LYS PRO PRO LYS VAL ILE GLU GLU TYR SER SER SER SER GLY SER GLY PRO PRO SER SER ASN ASP SER SER THR THR PRO PRO GLN LEU LEU TYR TYR ASP ASP THR THR LEU LEU LYS LYS TYR PHE ASN ASN

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	27602	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	42.45	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.537	Depositor
Minimum map value	-0.359	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.07	Depositor
Map size (Å)	419.99997, 419.99997, 419.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	B	0.19	0/1562	0.51	0/2113
2	C	0.20	0/1810	0.51	0/2435
3	D	0.24	0/4582	0.58	5/6211 (0.1%)
4	E	0.22	0/3504	0.54	0/4759
5	F	0.24	0/922	0.57	0/1249
6	G	0.21	0/1123	0.50	0/1518
7	H	0.24	0/1826	0.61	2/2456 (0.1%)
8	I	0.22	0/876	0.53	0/1183
9	K	0.20	0/2288	0.52	0/3080
10	Q	0.21	0/692	0.52	0/937
11	U	0.13	0/412	0.42	0/572
All	All	0.22	0/19597	0.55	7/26513 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	2

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	H	239	ASN	CA-C-N	5.82	131.69	122.61
7	H	239	ASN	C-N-CA	5.82	131.69	122.61
3	D	504	ASN	CA-C-N	5.67	132.17	121.97
3	D	504	ASN	C-N-CA	5.67	132.17	121.97
3	D	775	LYS	CA-CB-CG	5.50	125.11	114.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	476	ASP	Peptide
3	D	504	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1526	0	1469	23	0
2	C	1774	0	1790	25	0
3	D	4474	0	4377	59	0
4	E	3441	0	3493	52	0
5	F	902	0	919	24	0
6	G	1108	0	1110	20	0
7	H	1803	0	1800	38	0
8	I	865	0	855	23	0
9	K	2258	0	2308	28	0
10	Q	675	0	666	17	0
11	U	415	0	415	3	0
All	All	19241	0	19202	229	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:H:293:CYS:HB2	8:I:478:ILE:HD12	1.80	0.62
2:C:15:MET:HG2	2:C:246:THR:HG21	1.82	0.61
2:C:141:LEU:HB2	2:C:144:GLU:HG2	1.82	0.60
3:D:568:HIS:HB3	3:D:587:HIS:HB3	1.83	0.60
1:B:200:ARG:NH2	1:B:298:SER:OG	2.34	0.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	175/604 (29%)	163 (93%)	12 (7%)	0	100	100
2	C	206/337 (61%)	200 (97%)	6 (3%)	0	100	100
3	D	558/798 (70%)	510 (91%)	47 (8%)	1 (0%)	43	71
4	E	425/516 (82%)	393 (92%)	31 (7%)	1 (0%)	43	71
5	F	107/157 (68%)	104 (97%)	3 (3%)	0	100	100
6	G	138/206 (67%)	126 (91%)	12 (9%)	0	100	100
7	H	221/488 (45%)	203 (92%)	18 (8%)	0	100	100
8	I	106/539 (20%)	96 (91%)	10 (9%)	0	100	100
9	K	271/1332 (20%)	253 (93%)	18 (7%)	0	100	100
10	Q	82/657 (12%)	75 (92%)	7 (8%)	0	100	100
11	U	77/155 (50%)	73 (95%)	4 (5%)	0	100	100
All	All	2366/5789 (41%)	2196 (93%)	168 (7%)	2 (0%)	49	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	505	ILE
4	E	13	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	170/536 (32%)	170 (100%)	0	100	100
2	C	198/302 (66%)	198 (100%)	0	100	100
3	D	501/701 (72%)	498 (99%)	3 (1%)	78	81
4	E	391/464 (84%)	389 (100%)	2 (0%)	81	83
5	F	95/131 (72%)	95 (100%)	0	100	100
6	G	122/177 (69%)	120 (98%)	2 (2%)	55	72
7	H	206/434 (48%)	205 (100%)	1 (0%)	81	83
8	I	95/483 (20%)	94 (99%)	1 (1%)	65	76
9	K	257/1225 (21%)	257 (100%)	0	100	100
10	Q	77/579 (13%)	77 (100%)	0	100	100
All	All	2112/5032 (42%)	2103 (100%)	9 (0%)	81	84

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	H	353	LEU
8	I	437	VAL
4	E	128	VAL
4	E	148	ILE
6	G	130	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
6	G	144	ASN
7	H	352	GLN
6	G	157	GLN
7	H	219	ASN
8	I	485	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
11	U	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	U	29:ALA	C	101:ALA	N	57.64
1	U	9:ALA	C	11:ALA	N	3.53


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10414. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 185




Y Index: 232




Z Index: 136


6.3.2 Raw map



X Index: 184



Y Index: 233




Z Index: 145


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.07. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 133 nm³; this corresponds to an approximate mass of 120 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.303 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.303 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.41	4.18	3.57
Unmasked-calculated*	4.33	8.79	6.04

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.33 differs from the reported value 3.3 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-10414 and PDB model 6T9K. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.07).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.07) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7600	0.4730
B	0.8210	0.5100
C	0.7310	0.4520
D	0.7590	0.4810
E	0.7390	0.4590
F	0.8640	0.5170
G	0.6750	0.4530
H	0.7730	0.4820
I	0.8140	0.5090
K	0.7360	0.4530
Q	0.7330	0.4680
U	0.8550	0.4320

1.0

0.0

<0.0