



wwPDB EM Validation Summary Report ⓘ

Mar 28, 2026 – 03:18 AM UTC

PDB ID : 6TA6 / pdb_00006ta6
EMDB ID : EMD-10395
Title : MexAB assembly of the Pseudomonas MexAB-OprM efflux pump reconstituted in nanodiscs
Authors : Glavier, M.; Schoehn, G.; Taveau, J.C.; Phan, G.; Daury, L.; Lambert, O.; Broutin, I.
Deposited on : 2019-10-29
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	
1	B	474	
1	C	474	
2	D	366	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	366	 78%15%6%
2	F	366	 82%12%6%
2	G	366	 79%14%6%
2	H	366	 84%10%6%
2	I	366	 81%13%6%
3	J	1052	 80%17%.
3	K	1052	 78%19%..
3	L	1052	 79%18%..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 49550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane protein OprM.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	455	Total	C	N	O	S	0	0
			3487	2182	621	681	3		
1	B	455	Total	C	N	O	S	0	0
			3487	2182	621	681	3		
1	C	455	Total	C	N	O	S	0	0
			3487	2182	621	681	3		

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	469	HIS	-	expression tag	UNP Q51487
A	470	HIS	-	expression tag	UNP Q51487
A	471	HIS	-	expression tag	UNP Q51487
A	472	HIS	-	expression tag	UNP Q51487
A	473	HIS	-	expression tag	UNP Q51487
A	474	HIS	-	expression tag	UNP Q51487
B	469	HIS	-	expression tag	UNP Q51487
B	470	HIS	-	expression tag	UNP Q51487
B	471	HIS	-	expression tag	UNP Q51487
B	472	HIS	-	expression tag	UNP Q51487
B	473	HIS	-	expression tag	UNP Q51487
B	474	HIS	-	expression tag	UNP Q51487
C	469	HIS	-	expression tag	UNP Q51487
C	470	HIS	-	expression tag	UNP Q51487
C	471	HIS	-	expression tag	UNP Q51487
C	472	HIS	-	expression tag	UNP Q51487
C	473	HIS	-	expression tag	UNP Q51487
C	474	HIS	-	expression tag	UNP Q51487

- Molecule 2 is a protein called MexA family multidrug efflux RND transporter periplasmic adaptor subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	345	Total	C	N	O	S	0	0
			2615	1631	468	513	3		
2	E	343	Total	C	N	O	S	0	0
			2603	1625	466	509	3		
2	F	345	Total	C	N	O	S	0	0
			2615	1631	468	513	3		
2	G	343	Total	C	N	O	S	0	0
			2603	1625	466	509	3		
2	H	345	Total	C	N	O	S	0	0
			2614	1631	468	512	3		
2	I	343	Total	C	N	O	S	0	0
			2603	1625	466	509	3		

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	361	HIS	-	expression tag	UNP A0A2V3GTR8
D	362	HIS	-	expression tag	UNP A0A2V3GTR8
D	363	HIS	-	expression tag	UNP A0A2V3GTR8
D	364	HIS	-	expression tag	UNP A0A2V3GTR8
D	365	HIS	-	expression tag	UNP A0A2V3GTR8
D	366	HIS	-	expression tag	UNP A0A2V3GTR8
E	361	HIS	-	expression tag	UNP A0A2V3GTR8
E	362	HIS	-	expression tag	UNP A0A2V3GTR8
E	363	HIS	-	expression tag	UNP A0A2V3GTR8
E	364	HIS	-	expression tag	UNP A0A2V3GTR8
E	365	HIS	-	expression tag	UNP A0A2V3GTR8
E	366	HIS	-	expression tag	UNP A0A2V3GTR8
F	361	HIS	-	expression tag	UNP A0A2V3GTR8
F	362	HIS	-	expression tag	UNP A0A2V3GTR8
F	363	HIS	-	expression tag	UNP A0A2V3GTR8
F	364	HIS	-	expression tag	UNP A0A2V3GTR8
F	365	HIS	-	expression tag	UNP A0A2V3GTR8
F	366	HIS	-	expression tag	UNP A0A2V3GTR8
G	361	HIS	-	expression tag	UNP A0A2V3GTR8
G	362	HIS	-	expression tag	UNP A0A2V3GTR8
G	363	HIS	-	expression tag	UNP A0A2V3GTR8
G	364	HIS	-	expression tag	UNP A0A2V3GTR8
G	365	HIS	-	expression tag	UNP A0A2V3GTR8
G	366	HIS	-	expression tag	UNP A0A2V3GTR8
H	361	HIS	-	expression tag	UNP A0A2V3GTR8
H	362	HIS	-	expression tag	UNP A0A2V3GTR8
H	363	HIS	-	expression tag	UNP A0A2V3GTR8
H	364	HIS	-	expression tag	UNP A0A2V3GTR8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
H	365	HIS	-	expression tag	UNP A0A2V3GTR8
H	366	HIS	-	expression tag	UNP A0A2V3GTR8
I	361	HIS	-	expression tag	UNP A0A2V3GTR8
I	362	HIS	-	expression tag	UNP A0A2V3GTR8
I	363	HIS	-	expression tag	UNP A0A2V3GTR8
I	364	HIS	-	expression tag	UNP A0A2V3GTR8
I	365	HIS	-	expression tag	UNP A0A2V3GTR8
I	366	HIS	-	expression tag	UNP A0A2V3GTR8

- Molecule 3 is a protein called Efflux pump membrane transporter.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	J	1030	Total	C	N	O	S	0	0
			7812	5027	1298	1447	40		
3	K	1030	Total	C	N	O	S	0	0
			7812	5027	1298	1447	40		
3	L	1030	Total	C	N	O	S	0	0
			7812	5027	1298	1447	40		

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
J	1047	HIS	-	expression tag	UNP A0A069Q9M6
J	1048	HIS	-	expression tag	UNP A0A069Q9M6
J	1049	HIS	-	expression tag	UNP A0A069Q9M6
J	1050	HIS	-	expression tag	UNP A0A069Q9M6
J	1051	HIS	-	expression tag	UNP A0A069Q9M6
J	1052	HIS	-	expression tag	UNP A0A069Q9M6
K	1047	HIS	-	expression tag	UNP A0A069Q9M6
K	1048	HIS	-	expression tag	UNP A0A069Q9M6
K	1049	HIS	-	expression tag	UNP A0A069Q9M6
K	1050	HIS	-	expression tag	UNP A0A069Q9M6
K	1051	HIS	-	expression tag	UNP A0A069Q9M6
K	1052	HIS	-	expression tag	UNP A0A069Q9M6
L	1047	HIS	-	expression tag	UNP A0A069Q9M6
L	1048	HIS	-	expression tag	UNP A0A069Q9M6
L	1049	HIS	-	expression tag	UNP A0A069Q9M6
L	1050	HIS	-	expression tag	UNP A0A069Q9M6
L	1051	HIS	-	expression tag	UNP A0A069Q9M6
L	1052	HIS	-	expression tag	UNP A0A069Q9M6




- Molecule 2: MexA family multidrug efflux RND transporter periplasmic adaptor subunit




- Molecule 2: MexA family multidrug efflux RND transporter periplasmic adaptor subunit




- Molecule 2: MexA family multidrug efflux RND transporter periplasmic adaptor subunit




- Molecule 2: MexA family multidrug efflux RND transporter periplasmic adaptor subunit






• Molecule 3: Efflux pump membrane transporter


Chain K: 78% 19%



• Molecule 3: Efflux pump membrane transporter

Chain L: 79% 18%





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	82432	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	42	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.077	Depositor
Minimum map value	-0.024	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0125	Depositor
Map size (Å)	696.32, 696.32, 696.32	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.36, 1.36, 1.36	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.19	0/3543	0.47	2/4819 (0.0%)
1	B	0.19	0/3543	0.45	0/4819
1	C	0.20	0/3543	0.46	2/4819 (0.0%)
2	D	0.25	0/2649	0.57	1/3597 (0.0%)
2	E	0.22	0/2637	0.51	0/3580
2	F	0.23	0/2649	0.52	0/3597
2	G	0.23	0/2637	0.51	0/3580
2	H	0.23	0/2648	0.49	0/3595
2	I	0.22	0/2637	0.52	0/3580
3	J	0.26	0/7971	0.61	2/10833 (0.0%)
3	K	0.27	1/7971 (0.0%)	0.60	2/10833 (0.0%)
3	L	0.25	0/7971	0.61	2/10833 (0.0%)
All	All	0.24	1/50399 (0.0%)	0.55	11/68485 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	3
3	J	0	1
3	L	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	115	THR	C-N	5.54	1.38	1.33

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	60	THR	CA-C-N	-5.90	116.58	122.77
3	L	60	THR	C-N-CA	-5.90	116.58	122.77
2	D	196	LEU	N-CA-C	5.70	119.54	112.47
1	A	430	THR	CA-C-N	5.53	132.10	121.54
1	A	430	THR	C-N-CA	5.53	132.10	121.54

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	195	GLN	Peptide
2	D	265	GLN	Peptide
2	D	268	ILE	Peptide
3	J	469	GLN	Peptide
3	L	143	VAL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3487	0	3466	41	0
1	B	3487	0	3466	31	0
1	C	3487	0	3466	28	0
2	D	2615	0	2666	27	0
2	E	2603	0	2656	34	0
2	F	2615	0	2666	29	0
2	G	2603	0	2656	31	0
2	H	2614	0	2666	23	0
2	I	2603	0	2656	31	0
3	J	7812	0	7944	114	0
3	K	7812	0	7944	131	0
3	L	7812	0	7944	128	0
All	All	49550	0	50196	599	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 599 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:J:652:GLN:O	3:J:656:PHE:HB2	1.89	0.72
3:K:782:PRO:O	3:K:785:LEU:HB2	1.90	0.69
3:L:175:PHE:HB2	3:L:289:ILE:HD11	1.75	0.68
3:J:377:LEU:O	3:J:380:PHE:HB2	1.94	0.67
3:K:415:ASN:HD21	3:K:437:GLN:HE21	1.44	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/474 (96%)	447 (99%)	6 (1%)	0	100	100
1	B	453/474 (96%)	448 (99%)	5 (1%)	0	100	100
1	C	453/474 (96%)	448 (99%)	5 (1%)	0	100	100
2	D	343/366 (94%)	316 (92%)	26 (8%)	1 (0%)	36	68
2	E	341/366 (93%)	318 (93%)	22 (6%)	1 (0%)	36	68
2	F	343/366 (94%)	323 (94%)	18 (5%)	2 (1%)	21	56
2	G	341/366 (93%)	309 (91%)	31 (9%)	1 (0%)	36	68
2	H	343/366 (94%)	323 (94%)	19 (6%)	1 (0%)	36	68
2	I	341/366 (93%)	315 (92%)	25 (7%)	1 (0%)	36	68
3	J	1028/1052 (98%)	988 (96%)	40 (4%)	0	100	100
3	K	1028/1052 (98%)	979 (95%)	49 (5%)	0	100	100
3	L	1028/1052 (98%)	975 (95%)	53 (5%)	0	100	100
All	All	6495/6774 (96%)	6189 (95%)	299 (5%)	7 (0%)	49	79

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	73	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	74	ALA
2	D	73	PRO
2	E	73	PRO
2	I	73	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/375 (96%)	358 (100%)	0	100	100
1	B	358/375 (96%)	357 (100%)	1 (0%)	86	88
1	C	358/375 (96%)	358 (100%)	0	100	100
2	D	278/293 (95%)	278 (100%)	0	100	100
2	E	277/293 (94%)	276 (100%)	1 (0%)	84	86
2	F	278/293 (95%)	276 (99%)	2 (1%)	76	83
2	G	277/293 (94%)	276 (100%)	1 (0%)	84	86
2	H	278/293 (95%)	277 (100%)	1 (0%)	84	86
2	I	277/293 (94%)	276 (100%)	1 (0%)	84	86
3	J	841/860 (98%)	832 (99%)	9 (1%)	65	79
3	K	841/860 (98%)	834 (99%)	7 (1%)	73	82
3	L	841/860 (98%)	831 (99%)	10 (1%)	63	79
All	All	5262/5463 (96%)	5229 (99%)	33 (1%)	76	84

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	770	VAL
3	L	904	VAL
3	L	1015	LEU
3	J	854	VAL
3	J	690	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 134 such sidechains are listed below:

Mol	Chain	Res	Type
3	L	161	ASN
3	L	248	ASN
3	L	927	GLN
2	G	110	GLN
2	F	289	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10395. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 268



Y Index: 269




Z Index: 250


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0125. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 731 nm^3 ; this corresponds to an approximate mass of 661 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.12	3.63	3.15
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10395 and PDB model 6TA6. Per-residue inclusion information can be found in section [3](#) on page [7](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0125 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0125).

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0125) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8810	0.4180
A	0.8730	0.3320
B	0.8580	0.3460
C	0.8700	0.3320
D	0.9190	0.4680
E	0.9020	0.4490
F	0.9200	0.4700
G	0.8890	0.4400
H	0.9220	0.4690
I	0.9030	0.4510
J	0.8730	0.4340
K	0.8740	0.4330
L	0.8600	0.4160

1.0

0.0

<0.0