



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 01:03 PM UTC

PDB ID : 7TAA / pdb_00007taa
Title : FAMILY 13 ALPHA AMYLASE IN COMPLEX WITH ACARBOSE
Authors : Davies, G.J.; Brzozowski, A.M.
Deposited on : 1997-10-06
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4218 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TAKA AMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	476	3688	2336	595	739	18	0	2	0

- Molecule 2 is MODIFIED ACARBOSE HEXASACCHARIDE (CCD ID: ABC) (formula: $C_{37}H_{63}NO_{26}$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	465	Total 465	O 465	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TAKA AMYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.04Å 67.18Å 133.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.98 20.00 – 1.99	Depositor EDS
% Data completeness (in resolution range)	97.0 (20.00-1.98) 95.6 (20.00-1.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.85 (at 1.99Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.159 , 0.236 0.155 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.2	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 73.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4218	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ABC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	1/3793 (0.0%)	1.71	42/5180 (0.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	GLY	CA-C	5.99	1.57	1.51

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	PHE	CA-CB-CG	9.12	122.92	113.80
1	A	323	GLY	CA-C-N	-8.54	115.19	123.04
1	A	323	GLY	C-N-CA	-8.54	115.19	123.04
1	A	421	LEU	CA-C-O	-8.09	111.62	120.43
1	A	253	PRO	O-C-N	-7.64	113.93	122.18
1	A	182	VAL	CB-CA-C	-7.40	102.01	112.22
1	A	322	ASP	CA-CB-CG	7.13	119.73	112.60
1	A	63	THR	CA-C-N	6.78	126.74	119.76
1	A	63	THR	C-N-CA	6.78	126.74	119.76
1	A	386	PRO	N-CA-CB	6.15	107.35	103.23
1	A	64	PRO	N-CA-CB	6.12	108.75	103.31
1	A	117	ASP	CA-CB-CG	6.08	118.68	112.60
1	A	208	VAL	CA-C-O	-6.07	114.64	120.95
1	A	390	ASP	CA-CB-CG	6.05	118.65	112.60
1	A	71	GLN	N-CA-C	6.02	118.47	110.53
1	A	216	TRP	N-CA-C	5.98	121.65	113.16
1	A	123	MET	CA-C-O	-5.91	113.61	121.46
1	A	235	ASP	CA-C-O	5.89	123.68	119.32
1	A	359	TYR	N-CA-C	-5.79	104.87	111.07
1	A	225	VAL	CA-C-O	-5.77	115.98	121.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	98	ASP	CA-CB-CG	-5.76	106.84	112.60
1	A	239	THR	N-CA-C	5.71	118.36	111.40
1	A	251	ASN	OD1-CG-ND2	-5.69	116.91	122.60
1	A	11	ILE	N-CA-C	5.63	116.69	108.53
1	A	268	SER	N-CA-C	5.60	117.42	108.41
1	A	69	LEU	CA-C-N	5.59	125.29	119.64
1	A	69	LEU	C-N-CA	5.59	125.29	119.64
1	A	5	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	298	ASN	CA-CB-CG	-5.49	107.11	112.60
1	A	336	ALA	N-CA-C	5.44	119.69	112.30
1	A	420	THR	CA-C-N	-5.30	115.41	122.72
1	A	420	THR	C-N-CA	-5.30	115.41	122.72
1	A	89	SER	CA-CB-OG	5.24	121.58	111.10
1	A	193	SER	CA-CB-OG	-5.22	100.65	111.10
1	A	15	LEU	N-CA-CB	5.18	118.87	110.42
1	A	35	GLN	CG-CD-NE2	5.14	124.11	116.40
1	A	437	VAL	N-CA-C	5.12	120.04	113.07
1	A	168	ASP	CA-C-N	5.12	129.27	120.72
1	A	168	ASP	C-N-CA	5.12	129.27	120.72
1	A	390	ASP	CA-C-O	-5.10	115.30	121.32
1	A	143	GLN	CA-CB-CG	5.09	124.27	114.10
1	A	465	PRO	N-CA-CB	5.00	107.62	103.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3688	0	3481	40	0
2	A	64	0	63	0	0
3	A	1	0	0	0	0
4	A	465	0	0	16	1
All	All	4218	0	3544	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:LYS:HE2	1:A:383:LYS:HB3	1.56	0.85
1:A:441:THR:HB	4:A:881:HOH:O	1.77	0.83
1:A:465:PRO:HB2	1:A:468:LYS:HG3	1.66	0.77
1:A:441:THR:HG22	4:A:539:HOH:O	1.93	0.67
1:A:279:VAL:HA	4:A:926:HOH:O	1.95	0.66
1:A:371:TYR:O	1:A:374[A]:SER:HB2	2.02	0.59
1:A:240:CYS:SG	4:A:926:HOH:O	2.57	0.59
1:A:415:SER:HB2	4:A:523:HOH:O	2.03	0.59
1:A:36:LYS:HG2	1:A:37:TYR:N	2.17	0.58
1:A:467:GLU:HG2	1:A:468:LYS:N	2.20	0.57
1:A:31:ASN:HB2	4:A:846:HOH:O	2.12	0.50
1:A:448:ASP:HB3	4:A:866:HOH:O	2.11	0.50
1:A:280:LYS:HG3	1:A:383:LYS:HG2	1.94	0.49
1:A:36:LYS:HG2	1:A:37:TYR:H	1.77	0.49
1:A:2:THR:HB	1:A:3:PRO:HD2	1.94	0.49
1:A:15:LEU:HD22	1:A:63:THR:HG21	1.94	0.48
1:A:178:THR:O	1:A:184:LYS:HD2	2.13	0.48
1:A:87:ILE:HD12	1:A:190:TRP:CD1	2.49	0.48
1:A:169:ASN:HB3	4:A:568:HOH:O	2.13	0.47
1:A:352:TYR:N	1:A:353:PRO:CD	2.77	0.47
1:A:308:ILE:HD11	1:A:358:LEU:CD2	2.46	0.46
1:A:252:TYR:N	1:A:253:PRO:CD	2.79	0.46
1:A:184:LYS:NZ	4:A:645:HOH:O	2.40	0.44
1:A:373:ILE:HG22	1:A:379:PHE:CD2	2.52	0.44
1:A:352:TYR:N	1:A:353:PRO:HD3	2.32	0.44
1:A:371:TYR:O	1:A:374[B]:SER:HB3	2.18	0.44
1:A:100:LEU:HD13	1:A:198:TYR:CD1	2.53	0.44
1:A:273:TYR:CG	1:A:389:LYS:HE3	2.53	0.44
1:A:240:CYS:N	1:A:241:PRO:CD	2.82	0.43
1:A:467:GLU:HG2	1:A:468:LYS:HG2	2.00	0.43
1:A:197:ASN:ND2	4:A:602:HOH:O	2.39	0.43
1:A:169:ASN:HA	4:A:568:HOH:O	2.18	0.42
1:A:134:TYR:CD2	1:A:143:GLN:HB3	2.54	0.42
1:A:36:LYS:HE3	4:A:687:HOH:O	2.20	0.42
1:A:36:LYS:HG3	4:A:761:HOH:O	2.20	0.41
1:A:151:PHE:HE1	4:A:819:HOH:O	2.03	0.41
1:A:138:LYS:HA	1:A:139:PRO:HA	1.69	0.41
1:A:369:ARG:HD3	4:A:532:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:LYS:HD2	4:A:786:HOH:O	2.21	0.41
1:A:2:THR:HB	1:A:3:PRO:CD	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:570:HOH:O	4:A:762:HOH:O[3_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/478 (100%)	466 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/399 (100%)	373 (94%)	26 (6%)	15	6

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	61	TRP
1	A	100	LEU
1	A	131	SER
1	A	141	SER
1	A	153	GLN
1	A	160	GLN
1	A	169	ASN
1	A	182	VAL
1	A	184	LYS
1	A	187	TRP
1	A	207	THR
1	A	283	CYS
1	A	296	HIS
1	A	356	SER
1	A	364[A]	SER
1	A	364[B]	SER
1	A	373	ILE
1	A	374[A]	SER
1	A	374[B]	SER
1	A	396	MET
1	A	443	VAL
1	A	459	LEU
1	A	467	GLU
1	A	468	LYS
1	A	476	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	35	GLN
1	A	91	ASN
1	A	93	ASN
1	A	251	ASN
1	A	433	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ABC	A	479	-	68,69,69	1.10	3 (4%)	92,104,104	1.51	18 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ABC	A	479	-	-	6/26/146/146	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	479	ABC	C3I-C4I	3.96	1.59	1.53
2	A	479	ABC	C1I-C7I	3.13	1.54	1.50
2	A	479	ABC	C3H-C4H	2.07	1.57	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	479	ABC	O2K-C2K-C3K	-4.31	100.22	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	479	ABC	O2L-C2L-C1L	3.97	118.40	109.25
2	A	479	ABC	O2G-C2G-C1G	3.87	119.29	110.08
2	A	479	ABC	O4L-C4L-C3L	3.08	115.06	107.23
2	A	479	ABC	C1H-O4I-C4I	3.06	119.13	115.23
2	A	479	ABC	C1J-O5J-C5J	3.00	118.75	113.63
2	A	479	ABC	O2I-C2I-C1I	2.56	114.11	109.08
2	A	479	ABC	O5L-C5L-C4L	-2.53	104.51	109.19
2	A	479	ABC	C1J-C2J-C3J	2.45	115.16	110.01
2	A	479	ABC	O2H-C2H-C3H	-2.44	104.62	110.38
2	A	479	ABC	C7I-C1I-N1J	-2.44	107.09	110.68
2	A	479	ABC	O5L-C5L-C6L	-2.43	101.34	106.74
2	A	479	ABC	O5G-C5G-C6G	-2.37	101.49	106.74
2	A	479	ABC	O2J-C2J-C3J	-2.33	104.89	110.38
2	A	479	ABC	O4H-C4H-C3H	2.21	112.85	107.23
2	A	479	ABC	C1G-O5G-C5G	2.19	117.37	113.63
2	A	479	ABC	O6I-C6I-C5I	-2.17	105.92	112.74
2	A	479	ABC	C1K-O5K-C5K	2.12	117.86	113.72

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	479	ABC	C7I-C1I-N1J-C4J
2	A	479	ABC	C7I-C5I-C6I-O6I
2	A	479	ABC	O5G-C1G-O4H-C4H
2	A	479	ABC	C2G-C1G-O4H-C4H
2	A	479	ABC	C4I-C5I-C6I-O6I
2	A	479	ABC	C5I-C4I-O4I-C1H

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/478 (99%)	-0.50	3 (0%) 85 90	8, 17, 33, 54	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	476	SER	2.4
1	A	471	GLY	2.3
1	A	470	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ABC	A	479	64/64	0.96	0.06	5,15,30,34	0
3	CA	A	480	1/1	0.98	0.03	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.