



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 11:06 PM UTC

PDB ID : 3TB6 / pdb_00003tb6
Title : Structure of the effector-binding domain of arabinose repressor AraR from *Bacillus subtilis*
Authors : Rezacova, P.; Prochazkova, K.
Deposited on : 2011-08-05
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	 73% 19% • 5%
1	B	298	 75% 16% • • 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Arabinose metabolism transcriptional repressor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	8	0
			2269	1440	381	434	14			
1	B	283	Total	C	N	O	S	0	6	0
			2263	1436	377	436	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	GLY	-	expression tag	UNP P96711
A	66	ILE	-	expression tag	UNP P96711
A	67	ASP	-	expression tag	UNP P96711
A	68	PRO	-	expression tag	UNP P96711
A	69	PHE	-	expression tag	UNP P96711
A	70	THR	-	expression tag	UNP P96711
B	65	GLY	-	expression tag	UNP P96711
B	66	ILE	-	expression tag	UNP P96711
B	67	ASP	-	expression tag	UNP P96711
B	68	PRO	-	expression tag	UNP P96711
B	69	PHE	-	expression tag	UNP P96711
B	70	THR	-	expression tag	UNP P96711

- Molecule 2 is beta-L-arabinopyranose (CCD ID: ARB) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	132	Total 132	O 132	0	0
4	B	113	Total 113	O 113	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Arabinose metabolism transcriptional repressor

Chain A: 



- Molecule 1: Arabinose metabolism transcriptional repressor

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.07Å 106.33Å 111.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.91 – 2.21 29.91 – 2.21	Depositor EDS
% Data completeness (in resolution range)	86.7 (29.91-2.21) 87.0 (29.91-2.21)	Depositor EDS
R_{merge}	0.57	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.186 , 0.231 0.183 , 0.228	Depositor DCC
R_{free} test set	1660 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4809	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ARB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	6/2334 (0.3%)	0.95	3/3149 (0.1%)
1	B	0.70	3/2322 (0.1%)	0.93	3/3132 (0.1%)
All	All	0.78	9/4656 (0.2%)	0.94	6/6281 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	LYS	CE-NZ	12.51	1.86	1.49
1	A	337	HIS	CE1-NE2	12.49	1.45	1.32
1	A	338	LYS	CD-CE	11.78	1.87	1.52
1	A	337	HIS	CG-CD2	8.86	1.45	1.35
1	A	337	HIS	CG-ND1	7.45	1.46	1.38
1	B	259	GLU	CD-OE1	7.18	1.39	1.25
1	A	338	LYS	C-O	7.18	1.33	1.24
1	B	293	GLU	C-N	5.47	1.40	1.33
1	B	259	GLU	CD-OE2	5.44	1.35	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	338	LYS	CD-CE-NZ	-7.75	87.11	111.90
1	A	338	LYS	CG-CD-CE	-7.57	93.90	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	141	VAL	CB-CA-C	-7.44	99.05	110.50
1	B	135	HIS	N-CA-C	6.97	120.12	108.20
1	B	141	VAL	CB-CA-C	-5.92	101.69	110.33
1	B	141	VAL	N-CA-C	5.38	115.87	108.12

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	133	SER	Peptide
1	B	134	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2269	0	2315	43	0
1	B	2263	0	2295	39	0
2	A	10	0	10	0	0
2	B	10	0	10	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	132	0	0	3	0
4	B	113	0	0	5	0
All	All	4809	0	4646	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:LYS:CD	1:A:338:LYS:CE	1.87	1.51
1:A:338:LYS:CE	1:A:338:LYS:NZ	1.86	1.36
1:A:205:MET:HE1	1:A:254:VAL:HG13	1.15	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:MET:HE1	1:A:254:VAL:CG1	1.97	0.95
1:A:205:MET:CE	1:A:254:VAL:HG13	2.02	0.90
1:B:158:ASN:HB3	4:B:563:HOH:O	1.72	0.89
1:A:125[B]:ARG:HG3	1:A:152:ASN:OD1	1.74	0.86
1:A:338:LYS:CE	1:A:338:LYS:CG	2.59	0.80
1:B:133:SER:O	1:B:134:GLN:HG2	1.82	0.79
1:A:262:SER:HA	1:A:265:MET:HE2	1.64	0.79
1:B:226:HIS:HE1	4:B:585:HOH:O	1.69	0.74
1:A:283:MET:HE3	1:A:287:MET:HE1	1.71	0.72
1:B:205:MET:HE1	1:B:254:VAL:HG13	1.72	0.70
1:A:303:SER:OG	1:A:305:PHE:HB2	1.91	0.70
1:A:129:GLU:HG3	4:A:541:HOH:O	1.91	0.70
1:A:283:MET:HE3	1:A:287:MET:CE	2.22	0.70
1:B:205:MET:HE3	1:B:207:ILE:HD11	1.73	0.70
1:B:303:SER:OG	1:B:305:PHE:HB2	1.93	0.68
1:A:338:LYS:CD	1:A:338:LYS:NZ	2.58	0.67
1:A:79:ASN:N	1:A:81:THR:HG1	1.95	0.65
1:A:283:MET:HE3	1:A:284:LEU:HD13	1.79	0.64
1:A:170:ASN:C	1:A:182:THR:HG22	2.25	0.60
1:B:132:LEU:C	1:B:134:GLN:H	2.11	0.59
1:B:132:LEU:O	1:B:134:GLN:N	2.36	0.58
1:B:262:SER:HA	1:B:265:MET:HE2	1.86	0.58
1:B:172:SER:HB3	1:B:180:SER:OG	2.05	0.56
1:A:304[B]:HIS:NE2	1:B:308:ILE:HD11	2.21	0.56
1:B:197:LEU:HD13	1:B:204:MET:HE1	1.87	0.56
1:A:284:LEU:HD12	1:A:287:MET:HE3	1.88	0.56
1:B:213:THR:HG21	4:B:538:HOH:O	2.06	0.56
1:A:205:MET:HE2	1:A:269:ILE:HD13	1.88	0.55
1:A:130:ASN:O	1:A:134:GLN:NE2	2.41	0.54
1:A:281:ILE:HG21	1:A:311:VAL:HG21	1.89	0.53
1:B:158:ASN:ND2	4:B:562:HOH:O	2.41	0.52
1:A:224:GLN:O	1:A:228:GLU:HG2	2.11	0.51
1:A:319:PRO:HG2	1:A:323:LEU:HB2	1.92	0.50
1:B:132:LEU:C	1:B:134:GLN:N	2.69	0.50
1:B:313:LEU:HA	1:B:356:SER:HB3	1.93	0.49
1:A:141:VAL:O	1:A:168:MET:HA	2.12	0.49
1:B:300:TYR:O	1:B:301:ASP:HB2	2.13	0.49
1:A:290:LYS:HB3	1:A:293:GLU:HG2	1.95	0.48
1:A:208:PHE:O	1:A:240:THR:HA	2.13	0.48
1:A:165:PRO:HG3	1:A:335:LEU:HD13	1.96	0.48
1:B:182:THR:HB	4:B:547:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:341:LYS:HG2	1:B:343[A]:GLU:H	1.77	0.47
1:A:283:MET:HG2	1:A:287:MET:HE2	1.96	0.47
1:B:341:LYS:HG2	1:B:343[B]:GLU:H	1.77	0.47
1:B:205:MET:HE1	1:B:254:VAL:CG1	2.43	0.47
1:B:312:LYS:HB3	1:B:354:ARG:HD3	1.96	0.47
1:B:245:GLU:HG2	1:B:249:LYS:HD3	1.97	0.46
1:A:79:ASN:O	1:A:80:LYS:HB2	2.16	0.46
1:A:158:ASN:HB3	4:A:604:HOH:O	2.16	0.46
1:B:205:MET:HE2	1:B:269:ILE:CG2	2.46	0.45
1:B:209:LYS:O	1:B:215:GLY:HA3	2.16	0.45
1:B:86:THR:O	1:B:116:SER:HA	2.16	0.45
1:A:269:ILE:HB	1:A:297:ILE:HD13	1.99	0.44
1:A:273:ASN:HA	1:A:300:TYR:HB3	1.99	0.44
1:B:132:LEU:O	1:B:134:GLN:O	2.35	0.44
1:B:244:GLU:H	1:B:244:GLU:CD	2.25	0.44
1:A:134:GLN:OE1	1:B:134:GLN:HB2	2.18	0.44
1:A:182:THR:O	1:A:346:ILE:HA	2.17	0.44
1:A:343:GLU:HB2	4:A:553:HOH:O	2.17	0.44
1:B:157:LEU:HD13	1:B:176:LEU:CD2	2.48	0.43
1:B:85:LEU:HD23	1:B:141:VAL:CG1	2.48	0.43
1:A:131:LEU:HD12	1:A:131:LEU:HA	1.93	0.43
1:B:197:LEU:HG	1:B:229:ARG:HD2	1.99	0.43
1:A:303:SER:HG	1:A:305:PHE:HB2	1.83	0.43
1:B:205:MET:CE	1:B:207:ILE:HD11	2.44	0.43
1:A:281:ILE:HD13	1:A:311:VAL:HG21	2.02	0.42
1:A:134:GLN:CD	1:B:134:GLN:HB2	2.45	0.41
1:A:283:MET:CE	1:A:284:LEU:HD13	2.47	0.41
1:A:205:MET:HE3	1:A:207:ILE:HD11	2.03	0.41
1:B:273:ASN:HA	1:B:300:TYR:HB3	2.01	0.41
1:A:284:LEU:HA	1:A:287:MET:HE3	2.02	0.41
1:A:104:TYR:CE2	1:A:108:GLN:HG3	2.56	0.41
1:B:83:GLY:N	1:B:136:ILE:HD12	2.36	0.40
1:B:222:PHE:O	1:B:226:HIS:HD2	2.05	0.40
1:B:341:LYS:HD2	1:B:343[B]:GLU:OE1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/298 (97%)	278 (96%)	7 (2%)	4 (1%)	9	7
1	B	287/298 (96%)	278 (97%)	7 (2%)	2 (1%)	18	18
All	All	576/596 (97%)	556 (96%)	14 (2%)	6 (1%)	11	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	ASP
1	A	301	ASP
1	B	133	SER
1	B	301	ASP
1	A	135	HIS
1	A	247	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/262 (98%)	239 (93%)	19 (7%)	13	13
1	B	256/262 (98%)	234 (91%)	22 (9%)	10	9
All	All	514/524 (98%)	473 (92%)	41 (8%)	11	11

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	SER
1	A	105	LEU
1	A	107	GLU
1	A	131	LEU
1	A	134	GLN
1	A	136	ILE
1	A	141	VAL
1	A	159	LEU
1	A	249	LYS
1	A	251	LEU
1	A	263	LYS
1	A	284	LEU
1	A	289	LEU
1	A	293	GLU
1	A	313	LEU
1	A	335	LEU
1	A	341	LYS
1	A	360	LEU
1	A	361	ASN
1	B	105	LEU
1	B	129[A]	GLU
1	B	129[B]	GLU
1	B	131	LEU
1	B	132	LEU
1	B	136	ILE
1	B	141	VAL
1	B	157	LEU
1	B	159	LEU
1	B	197	LEU
1	B	213	THR
1	B	244	GLU
1	B	251	LEU
1	B	278	LEU
1	B	282	ASP
1	B	284	LEU
1	B	293	GLU
1	B	308	ILE
1	B	310	GLU
1	B	351	LEU
1	B	359	LYS
1	B	360	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	162	ASN
1	B	135	HIS
1	B	158	ASN
1	B	226	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401	-	5,5,5	0.56	0	5,5,5	0.71	0
2	ARB	A	400	-	10,10,10	0.90	0	14,14,14	1.35	4 (28%)
3	GOL	B	401	-	5,5,5	0.41	0	5,5,5	0.69	0
2	ARB	B	400	-	10,10,10	0.68	0	14,14,14	0.95	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401	-	-	2/4/4/4	-
2	ARB	A	400	-	-	-	0/1/1/1
2	ARB	B	400	-	-	-	0/1/1/1
3	GOL	B	401	-	-	4/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	ARB	O4-C4-C3	2.79	115.94	110.15
2	B	400	ARB	C5-O5-C1	2.53	117.72	112.46
2	A	400	ARB	C5-C4-C3	-2.37	106.19	109.64
2	A	400	ARB	O3-C3-C4	2.27	114.68	110.05
2	A	400	ARB	C5-O5-C1	2.12	116.86	112.46

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	GOL	O1-C1-C2-C3
3	B	401	GOL	O1-C1-C2-C3
3	B	401	GOL	C1-C2-C3-O3
3	B	401	GOL	O2-C2-C3-O3
3	A	401	GOL	O1-C1-C2-O2
3	B	401	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/298 (94%)	-0.23	1 (0%) 88 87	26, 47, 81, 110	10 (3%)
1	B	283/298 (94%)	0.01	4 (1%) 73 72	26, 53, 103, 134	7 (2%)
All	All	566/596 (94%)	-0.11	5 (0%) 81 79	26, 50, 90, 134	17 (3%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	361	ASN	3.0
1	B	360	LEU	2.5
1	B	133	SER	2.3
1	B	135	HIS	2.3
1	B	202	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	401	6/6	0.78	0.20	59,64,66,67	0
3	GOL	B	401	6/6	0.91	0.10	58,60,62,63	0
2	ARB	A	400	10/10	0.97	0.06	35,39,40,40	0
2	ARB	B	400	10/10	0.97	0.05	41,43,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.