



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 03:26 AM UTC

PDB ID : 3TBS / pdb_00003tbs
Title : CRYSTAL STRUCTURE OF THE MURINE CLASS I MAJOR HISTOCOMPATIBILITY COMPLEX H-2DB IN COMPLEX WITH LCMV-DERIVED GP33 ALTERED PEPTIDE ligand (V3P,Y4A)
Authors : Duru, A.D.; Allerbring, E.B.; Uchtenhagen, H.; Mazumdar, P.A.; Badia-Martinez, D.; Madhurantakam, C.; Sandalova, T.; Nygren, P.; Achour, A.
Deposited on : 2011-08-08
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	9% 76% 18% • •
1	D	276	11% 76% 18% • •
2	B	99	3% 85% 14% •
2	E	99	2% 81% 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	9	 89%11%
3	F	9	 89%11%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	0	0
			2229	1409	394	417	9			
1	D	271	Total	C	N	O	S	0	0	0
			2217	1402	392	414	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			
2	E	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			

- Molecule 3 is a protein called GLYCOPROTEIN G1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			66	42	11	12	1			
3	F	9	Total	C	N	O	S	0	0	0
			66	42	11	12	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	3	PRO	VAL	engineered mutation	UNP P07399
C	4	ALA	TYR	engineered mutation	UNP P07399
C	9	MET	CYS	engineered mutation	UNP P07399
F	3	PRO	VAL	engineered mutation	UNP P07399
F	4	ALA	TYR	engineered mutation	UNP P07399
F	9	MET	CYS	engineered mutation	UNP P07399

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	46	Total 46	O 46	0	0
6	B	27	Total 27	O 27	0	0
6	C	5	Total 5	O 5	0	0
6	D	41	Total 41	O 41	0	0
6	E	25	Total 25	O 25	0	0
6	F	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



● Molecule 3: GLYCOPROTEIN G1



● Molecule 3: GLYCOPROTEIN G1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.84Å 125.12Å 101.32Å 90.00° 128.15° 90.00°	Depositor
Resolution (Å)	49.21 – 2.49 49.21 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.6 (49.21-2.49) 99.6 (49.21-2.49)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.230 , 0.270 0.233 , 0.265	Depositor DCC
R_{free} test set	2006 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6389	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	1/2293 (0.0%)	0.82	1/3111 (0.0%)
1	D	0.72	1/2281 (0.0%)	0.84	2/3095 (0.1%)
2	B	0.80	0/847	0.84	0/1148
2	E	0.72	0/847	0.82	0/1148
3	C	0.70	0/67	0.68	0/88
3	F	0.91	0/67	0.99	0/88
All	All	0.73	2/6402 (0.0%)	0.83	3/8678 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	181	ARG	CG-CD	-9.29	1.24	1.52
1	A	181	ARG	CG-CD	-6.70	1.32	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	181	ARG	CB-CG-CD	7.59	128.76	111.30
1	D	181	ARG	CB-CG-CD	6.79	126.93	111.30
1	D	178	THR	N-CA-C	-5.19	104.94	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2229	0	2097	44	0
1	D	2217	0	2079	46	0
2	B	821	0	796	9	0
2	E	821	0	796	13	0
3	C	66	0	68	1	0
3	F	66	0	68	1	0
4	A	5	0	0	0	0
4	D	5	0	0	1	0
5	A	6	0	8	1	0
5	D	6	0	8	0	0
6	A	46	0	0	2	0
6	B	27	0	0	4	0
6	C	5	0	0	0	0
6	D	41	0	0	1	0
6	E	25	0	0	0	0
6	F	3	0	0	0	0
All	All	6389	0	5920	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:LEU:C	1:A:179:LEU:HD23	1.62	1.21
1:A:141:GLN:HA	1:A:141:GLN:HE21	1.08	1.16
2:B:99:MET:HE2	6:B:114:HOH:O	1.52	1.07
1:A:179:LEU:C	1:A:179:LEU:CD2	2.30	1.04
1:A:179:LEU:HD23	1:A:179:LEU:O	1.65	0.93
1:D:127:ASN:OD1	1:D:134:THR:HG22	1.75	0.86
1:A:141:GLN:HA	1:A:141:GLN:NE2	1.90	0.83
1:A:141:GLN:HE21	1:A:141:GLN:CA	1.88	0.79
1:A:176:ASN:O	1:A:177:ALA:HB3	1.84	0.77
1:A:176:ASN:O	1:A:177:ALA:CB	2.30	0.77
5:A:340:GOL:O1	5:A:340:GOL:O3	2.05	0.74
1:A:176:ASN:HD22	1:A:176:ASN:N	1.86	0.74
2:B:97:ARG:HG2	6:B:115:HOH:O	1.89	0.71
1:A:127:ASN:OD1	1:A:134:THR:HG22	1.91	0.70
1:D:155:HIS:HB3	3:F:6:PHE:CZ	2.28	0.68
1:A:179:LEU:HD23	1:A:180:LEU:N	2.10	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:35:ARG:NH2	2:E:54:MET:O	2.29	0.65
1:A:249:VAL:HG22	1:A:257:TYR:CE2	2.33	0.63
1:D:127:ASN:OD1	1:D:134:THR:CG2	2.47	0.63
1:A:155:HIS:HB3	3:C:6:PHE:CZ	2.34	0.63
1:A:177:ALA:C	1:A:179:LEU:N	2.56	0.62
1:D:234:ARG:HD3	2:E:10:TYR:CE2	2.36	0.61
1:D:141:GLN:NE2	1:D:141:GLN:HA	2.16	0.61
1:D:234:ARG:HD3	2:E:10:TYR:CZ	2.36	0.61
1:D:141:GLN:CA	1:D:141:GLN:HE21	2.16	0.58
1:A:252:GLY:N	1:A:254:GLU:OE1	2.37	0.58
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.39	0.57
1:D:90:GLY:N	4:D:339:SO4:O3	2.28	0.57
1:A:179:LEU:CD2	1:A:180:LEU:N	2.64	0.57
1:D:249:VAL:HG22	1:D:257:TYR:CE2	2.39	0.57
1:A:176:ASN:N	1:A:176:ASN:ND2	2.48	0.57
1:A:234:ARG:HD3	2:B:10:TYR:CE2	2.41	0.56
1:D:141:GLN:HA	1:D:141:GLN:HE21	1.71	0.55
1:D:252:GLY:N	1:D:254:GLU:OE1	2.40	0.55
1:A:234:ARG:HD3	2:B:10:TYR:CZ	2.44	0.53
1:D:175:GLY:O	1:D:177:ALA:HB3	2.10	0.52
1:D:234:ARG:NH2	2:E:99:MET:O	2.33	0.51
1:A:19:GLU:HA	1:A:19:GLU:OE1	2.10	0.51
1:D:141:GLN:NE2	1:D:141:GLN:CA	2.73	0.51
1:D:48:ARG:NH2	2:E:53:ASP:OD2	2.45	0.50
1:D:35:ARG:CZ	2:E:53:ASP:CB	2.90	0.50
1:D:185:PRO:HD2	1:D:266:LEU:HD13	1.93	0.50
1:D:29:ASP:HB2	1:D:179:LEU:HD21	1.94	0.49
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.48	0.49
1:D:219:LEU:O	1:D:219:LEU:HG	2.12	0.49
1:D:271:THR:C	1:D:272:LEU:HD23	2.38	0.49
1:A:219:LEU:O	1:A:219:LEU:HG	2.13	0.49
1:D:35:ARG:NH2	2:E:53:ASP:HB3	2.27	0.49
1:A:78:LEU:CD1	1:A:95:LEU:HB2	2.44	0.48
1:D:141:GLN:HE21	1:D:141:GLN:N	2.11	0.48
1:D:35:ARG:CZ	2:E:53:ASP:HB3	2.42	0.48
1:D:111:ARG:HG2	1:D:113:TYR:CE1	2.49	0.48
1:D:19:GLU:HA	1:D:19:GLU:OE1	2.14	0.47
1:A:185:PRO:HD2	1:A:266:LEU:HD13	1.96	0.47
1:D:179:LEU:HD23	1:D:179:LEU:HA	1.73	0.47
1:A:111:ARG:HG2	1:A:113:TYR:CE1	2.50	0.46
1:D:81:LEU:HD13	1:D:118:TYR:CD1	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:ASN:ND2	1:A:176:ASN:H	2.13	0.46
1:A:78:LEU:HD13	1:A:95:LEU:HB2	1.96	0.46
1:A:127:ASN:OD1	1:A:134:THR:CG2	2.62	0.46
1:D:6:ARG:NH2	1:D:113:TYR:CE1	2.84	0.45
1:A:264:GLU:HG2	6:A:353:HOH:O	2.16	0.45
1:D:178:THR:O	1:D:178:THR:OG1	2.34	0.45
1:D:189:VAL:HG23	1:D:272:LEU:HD12	1.99	0.45
1:A:75:ARG:CZ	6:A:348:HOH:O	2.64	0.44
2:B:3:LYS:HE3	6:B:125:HOH:O	2.17	0.44
1:A:271:THR:C	1:A:272:LEU:HD23	2.43	0.44
1:D:78:LEU:CD1	1:D:95:LEU:HB2	2.48	0.44
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.53	0.43
1:D:12:VAL:HG22	1:D:94:THR:HG22	1.98	0.43
2:B:23:LEU:HD23	2:B:39:MET:HE2	2.00	0.43
1:A:202:ARG:HD3	1:A:244:TRP:CE3	2.54	0.43
1:A:106:ASP:O	1:A:107:TRP:HB2	2.18	0.43
2:B:99:MET:CE	6:B:114:HOH:O	2.32	0.43
6:D:349:HOH:O	2:E:3:LYS:HE3	2.19	0.43
1:A:196:LYS:O	1:A:198:GLU:N	2.52	0.42
1:D:33:PHE:C	1:D:48:ARG:HB2	2.44	0.42
1:A:238:ASP:OD1	1:A:238:ASP:C	2.63	0.42
1:D:179:LEU:HD22	1:D:179:LEU:O	2.19	0.42
1:A:6:ARG:NH2	1:A:113:TYR:CE1	2.88	0.42
1:A:74:PHE:HZ	1:A:97:GLN:HE21	1.67	0.42
1:A:203:CYS:HB2	1:A:217:TRP:CZ2	2.54	0.42
1:D:202:ARG:HD3	1:D:244:TRP:CE3	2.55	0.42
1:D:194:ARG:HG3	1:D:195:SER:N	2.34	0.42
1:D:249:VAL:HG22	1:D:257:TYR:CZ	2.55	0.42
1:A:189:VAL:HG23	1:A:272:LEU:HD12	2.01	0.41
1:A:33:PHE:C	1:A:48:ARG:HB2	2.45	0.41
1:A:262:TYR:CG	1:D:108:ARG:HD3	2.55	0.41
1:D:35:ARG:CZ	2:E:53:ASP:HB2	2.50	0.41
2:E:79:ALA:HB2	2:E:94:TYR:CD1	2.55	0.41
1:A:32:GLU:OE2	1:A:48:ARG:HD2	2.20	0.41
1:A:234:ARG:NH2	2:B:99:MET:O	2.38	0.41
1:D:231:VAL:HB	2:E:8:GLN:HE22	1.86	0.41
1:D:11:ALA:HA	1:D:21:ARG:O	2.21	0.41
1:D:106:ASP:O	1:D:107:TRP:HB2	2.21	0.40
1:D:179:LEU:HD22	1:D:179:LEU:C	2.45	0.40
1:D:203:CYS:HB2	1:D:217:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/276 (96%)	259 (97%)	6 (2%)	1 (0%)	30	49
1	D	265/276 (96%)	257 (97%)	7 (3%)	1 (0%)	30	49
2	B	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
2	E	97/99 (98%)	93 (96%)	3 (3%)	1 (1%)	12	24
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	739/768 (96%)	715 (97%)	21 (3%)	3 (0%)	30	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	197	GLY
1	D	197	GLY
2	E	47	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/234 (98%)	202 (88%)	27 (12%)	5	11
1	D	226/234 (97%)	202 (89%)	24 (11%)	6	14
2	B	94/94 (100%)	85 (90%)	9 (10%)	8	17
2	E	94/94 (100%)	85 (90%)	9 (10%)	8	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	655/668 (98%)	586 (90%)	69 (10%)	6	14

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	17	LEU
1	A	39	ASP
1	A	72	GLN
1	A	78	LEU
1	A	82	LEU
1	A	114	LEU
1	A	134	THR
1	A	141	GLN
1	A	149	GLN
1	A	154	GLU
1	A	166	GLU
1	A	176	ASN
1	A	179	LEU
1	A	181	ARG
1	A	194	ARG
1	A	215	LEU
1	A	218	GLN
1	A	219	LEU
1	A	225	THR
1	A	230	LEU
1	A	234	ARG
1	A	247	VAL
1	A	249	VAL
1	A	254	GLU
1	A	266	LEU
1	A	272	LEU
2	B	1	ILE
2	B	4	THR
2	B	29	GLN
2	B	64	ILE
2	B	75	THR
2	B	77	THR
2	B	81	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	93	VAL
2	B	97	ARG
1	D	15	PRO
1	D	17	LEU
1	D	72	GLN
1	D	78	LEU
1	D	82	LEU
1	D	114	LEU
1	D	134	THR
1	D	141	GLN
1	D	149	GLN
1	D	154	GLU
1	D	166	GLU
1	D	178	THR
1	D	179	LEU
1	D	194	ARG
1	D	215	LEU
1	D	218	GLN
1	D	219	LEU
1	D	230	LEU
1	D	234	ARG
1	D	247	VAL
1	D	249	VAL
1	D	254	GLU
1	D	266	LEU
1	D	272	LEU
2	E	1	ILE
2	E	4	THR
2	E	29	GLN
2	E	64	ILE
2	E	75	THR
2	E	77	THR
2	E	81	ARG
2	E	93	VAL
2	E	97	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	97	GLN
1	A	141	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	149	GLN
1	A	176	ASN
1	A	192	HIS
1	A	220	ASN
2	B	13	HIS
2	B	67	HIS
3	C	5	ASN
1	D	72	GLN
1	D	97	GLN
1	D	141	GLN
1	D	149	GLN
1	D	192	HIS
1	D	220	ASN
1	D	255	GLN
2	E	8	GLN
2	E	67	HIS
3	F	5	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	340	-	5,5,5	0.41	0	5,5,5	0.22	0
4	SO4	D	339	-	4,4,4	0.26	0	6,6,6	0.16	0
4	SO4	A	339	-	4,4,4	0.24	0	6,6,6	0.41	0
5	GOL	D	340	-	5,5,5	0.48	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	340	-	-	4/4/4/4	-
5	GOL	D	340	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	340	GOL	O1-C1-C2-C3
5	A	340	GOL	C1-C2-C3-O3
5	D	340	GOL	C1-C2-C3-O3
5	A	340	GOL	O1-C1-C2-O2
5	A	340	GOL	O2-C2-C3-O3
5	D	340	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	340	GOL	1	0
4	D	339	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/276 (98%)	0.53	24 (8%) 15 14	28, 51, 94, 110	0
1	D	271/276 (98%)	0.67	30 (11%) 10 8	30, 55, 94, 114	0
2	B	99/99 (100%)	0.16	3 (3%) 52 48	29, 43, 65, 73	0
2	E	99/99 (100%)	0.22	2 (2%) 65 61	35, 46, 60, 66	0
3	C	9/9 (100%)	-0.25	0 100 100	36, 37, 42, 44	0
3	F	9/9 (100%)	-0.27	0 100 100	31, 34, 44, 49	0
All	All	759/768 (98%)	0.47	59 (7%) 19 17	28, 49, 90, 114	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	180	LEU	6.2
1	A	180	LEU	5.2
1	A	179	LEU	5.1
1	D	177	ALA	4.3
1	A	174	ASN	3.9
1	D	175	GLY	3.8
1	D	178	THR	3.8
1	A	196	LYS	3.8
1	A	230	LEU	3.8
1	D	196	LYS	3.6
1	A	220	ASN	3.6
1	D	251	LEU	3.6
1	D	219	LEU	3.5
1	D	57	PRO	3.4
1	A	16	GLY	3.4
1	D	183	ASP	3.3
1	A	17	LEU	3.3
1	A	177	ALA	3.3
2	B	1	ILE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	THR	3.3
1	D	230	LEU	3.3
1	D	1	GLY	3.3
2	E	1	ILE	3.3
1	D	179	LEU	3.1
1	D	182	THR	3.0
1	A	251	LEU	2.9
1	D	195	SER	2.9
1	A	222	GLU	2.9
1	D	59	TYR	2.8
1	D	224	LEU	2.8
1	D	275	GLU	2.8
1	D	89	ALA	2.8
1	A	253	LYS	2.7
1	A	275	GLU	2.6
1	A	258	THR	2.6
1	A	274	TRP	2.5
1	A	219	LEU	2.5
1	D	16	GLY	2.5
1	A	195	SER	2.5
1	A	175	GLY	2.4
1	A	249	VAL	2.4
1	D	220	ASN	2.4
1	D	72	GLN	2.4
1	A	224	LEU	2.4
1	D	253	LYS	2.4
1	D	222	GLU	2.4
2	B	74	GLU	2.3
1	A	176	ASN	2.3
1	D	51	TRP	2.2
1	D	193	PRO	2.2
1	D	199	VAL	2.2
1	D	105	SER	2.2
1	A	197	GLY	2.2
1	D	75	ARG	2.1
2	B	48	LYS	2.1
2	E	19	LYS	2.1
1	A	181	ARG	2.1
1	D	88	SER	2.0
1	D	181	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	D	340	6/6	0.62	0.27	80,83,83,84	0
5	GOL	A	340	6/6	0.87	0.13	63,63,64,64	0
4	SO4	D	339	5/5	0.91	0.15	73,73,74,74	0
4	SO4	A	339	5/5	0.96	0.09	54,55,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.