



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2026 – 10:20 PM UTC

PDB ID : 3TCF / pdb_00003tcf
Title : Crystal structure of E. coli OppA complexed with endogenous ligands
Authors : Klepsch, M.M.; Kovermann, M.; Low, C.; Balbach, J.; de Gier, J.W.; Slotboom, D.J.; Berntsson, R.P.-A.
Deposited on : 2011-08-09
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.












Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	524	78% 19% ..
1	B	524	5% 85% 13% .
1	C	524	3% 86% 12% ..
1	D	524	2% 85% 12% ..
1	E	524	85% 13% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	524	
1	G	524	
1	H	524	
2	I	3	
2	J	3	
2	K	3	
2	L	3	
2	M	3	
2	N	3	
2	O	3	
2	P	3	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 35615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic oligopeptide-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	0	2	0
			4137	2636	695	796	10			
1	B	517	Total	C	N	O	S	0	2	0
			4136	2636	694	796	10			
1	C	517	Total	C	N	O	S	0	1	0
			4129	2632	693	794	10			
1	D	517	Total	C	N	O	S	0	1	0
			4129	2632	693	794	10			
1	E	517	Total	C	N	O	S	0	2	0
			4137	2636	694	797	10			
1	F	517	Total	C	N	O	S	0	4	0
			4155	2649	696	800	10			
1	G	517	Total	C	N	O	S	0	2	0
			4135	2635	694	796	10			
1	H	517	Total	C	N	O	S	0	1	0
			4129	2632	693	794	10			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	expression tag	UNP P23843
A	544	HIS	-	expression tag	UNP P23843
A	545	HIS	-	expression tag	UNP P23843
A	546	HIS	-	expression tag	UNP P23843
A	547	HIS	-	expression tag	UNP P23843
A	548	HIS	-	expression tag	UNP P23843
A	549	HIS	-	expression tag	UNP P23843
B	26	MET	-	expression tag	UNP P23843
B	544	HIS	-	expression tag	UNP P23843
B	545	HIS	-	expression tag	UNP P23843
B	546	HIS	-	expression tag	UNP P23843
B	547	HIS	-	expression tag	UNP P23843
B	548	HIS	-	expression tag	UNP P23843

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	549	HIS	-	expression tag	UNP P23843
C	26	MET	-	expression tag	UNP P23843
C	544	HIS	-	expression tag	UNP P23843
C	545	HIS	-	expression tag	UNP P23843
C	546	HIS	-	expression tag	UNP P23843
C	547	HIS	-	expression tag	UNP P23843
C	548	HIS	-	expression tag	UNP P23843
C	549	HIS	-	expression tag	UNP P23843
D	26	MET	-	expression tag	UNP P23843
D	544	HIS	-	expression tag	UNP P23843
D	545	HIS	-	expression tag	UNP P23843
D	546	HIS	-	expression tag	UNP P23843
D	547	HIS	-	expression tag	UNP P23843
D	548	HIS	-	expression tag	UNP P23843
D	549	HIS	-	expression tag	UNP P23843
E	26	MET	-	expression tag	UNP P23843
E	544	HIS	-	expression tag	UNP P23843
E	545	HIS	-	expression tag	UNP P23843
E	546	HIS	-	expression tag	UNP P23843
E	547	HIS	-	expression tag	UNP P23843
E	548	HIS	-	expression tag	UNP P23843
E	549	HIS	-	expression tag	UNP P23843
F	26	MET	-	expression tag	UNP P23843
F	544	HIS	-	expression tag	UNP P23843
F	545	HIS	-	expression tag	UNP P23843
F	546	HIS	-	expression tag	UNP P23843
F	547	HIS	-	expression tag	UNP P23843
F	548	HIS	-	expression tag	UNP P23843
F	549	HIS	-	expression tag	UNP P23843
G	26	MET	-	expression tag	UNP P23843
G	544	HIS	-	expression tag	UNP P23843
G	545	HIS	-	expression tag	UNP P23843
G	546	HIS	-	expression tag	UNP P23843
G	547	HIS	-	expression tag	UNP P23843
G	548	HIS	-	expression tag	UNP P23843
G	549	HIS	-	expression tag	UNP P23843
H	26	MET	-	expression tag	UNP P23843
H	544	HIS	-	expression tag	UNP P23843
H	545	HIS	-	expression tag	UNP P23843
H	546	HIS	-	expression tag	UNP P23843
H	547	HIS	-	expression tag	UNP P23843
H	548	HIS	-	expression tag	UNP P23843

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	549	HIS	-	expression tag	UNP P23843

- Molecule 2 is a protein called Endogenous peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	J	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	K	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	L	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	M	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	N	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	O	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	P	3	Total	C	N	O	0	0	0
			15	9	3	3			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	360	Total	O	0	0
			360	360		
3	B	268	Total	O	0	0
			268	268		
3	C	285	Total	O	0	0
			285	285		
3	D	265	Total	O	0	0
			265	265		
3	E	351	Total	O	0	0
			351	351		
3	F	344	Total	O	0	0
			344	344		
3	G	305	Total	O	0	0
			305	305		
3	H	221	Total	O	0	0
			221	221		
3	I	2	Total	O	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	J	2	Total 2	O 2	0	0
3	K	1	Total 1	O 1	0	0
3	M	2	Total 2	O 2	0	0
3	P	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Periplasmic oligopeptide-binding protein




- Molecule 1: Periplasmic oligopeptide-binding protein





- Molecule 1: Periplasmic oligopeptide-binding protein









• Molecule 1: Periplasmic oligopeptide-binding protein




• Molecule 1: Periplasmic oligopeptide-binding protein




• Molecule 1: Periplasmic oligopeptide-binding protein




- Molecule 1: Periplasmic oligopeptide-binding protein



- Molecule 1: Periplasmic oligopeptide-binding protein



- Molecule 2: Endogenous peptide



- Molecule 2: Endogenous peptide

Chain J:  67% 33%

 X1 X2 X3

- Molecule 2: Endogenous peptide

Chain K:  67% 33%

 X1 X2 X3

- Molecule 2: Endogenous peptide

Chain L:  67% 33%

 X1 X2 X3

- Molecule 2: Endogenous peptide

Chain M:  33% 67%

 X1 X2 X3

- Molecule 2: Endogenous peptide

Chain N:  67% 33%

 X1 X2 X3

- Molecule 2: Endogenous peptide

Chain O:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Endogenous peptide

Chain P:  67% 33%

 X1 X2 X3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.62Å 202.60Å 208.95Å 90.00° 95.71° 90.00°	Depositor
Resolution (Å)	48.37 – 2.00 48.37 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.9 (48.37-2.00) 99.2 (48.37-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.229 , 0.267 0.230 , 0.267	Depositor DCC
R_{free} test set	17271 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.225	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	35615	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.69 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.6804e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	2/4245 (0.0%)	0.98	8/5790 (0.1%)
1	B	0.85	0/4244	0.84	4/5789 (0.1%)
1	C	0.85	0/4237	0.83	0/5779
1	D	0.82	0/4237	0.86	2/5779 (0.0%)
1	E	0.87	0/4245	0.86	1/5790 (0.0%)
1	F	0.90	0/4264	0.87	4/5817 (0.1%)
1	G	0.86	0/4243	0.88	3/5788 (0.1%)
1	H	0.83	0/4237	0.83	2/5779 (0.0%)
All	All	0.87	2/33952 (0.0%)	0.87	24/46311 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	397	HIS	CG-CD2	5.62	1.42	1.35
1	A	60	VAL	CA-CB	5.48	1.57	1.54

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	110	ASP	N-CA-C	7.22	120.20	111.82
1	A	110	ASP	N-CA-C	7.14	121.09	112.38
1	H	207	GLN	CA-C-N	6.29	127.71	119.84
1	H	207	GLN	C-N-CA	6.29	127.71	119.84
1	B	110	ASP	N-CA-C	6.22	119.71	111.75

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	483	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4137	0	4046	101	0
1	B	4136	0	4047	59	0
1	C	4129	0	4041	74	0
1	D	4129	0	4041	55	0
1	E	4137	0	4044	69	0
1	F	4155	0	4061	88	0
1	G	4135	0	4046	112	0
1	H	4129	0	4041	102	0
2	I	15	0	5	2	0
2	J	15	0	5	1	0
2	K	15	0	5	1	0
2	L	15	0	5	1	0
2	M	15	0	5	2	0
2	N	15	0	5	1	0
2	O	15	0	5	0	0
2	P	15	0	5	1	0
3	A	360	0	0	37	0
3	B	268	0	0	12	0
3	C	285	0	0	26	0
3	D	265	0	0	9	0
3	E	351	0	0	20	0
3	F	344	0	0	36	0
3	G	305	0	0	33	0
3	H	221	0	0	41	0
3	I	2	0	0	1	0
3	J	2	0	0	0	0
3	K	1	0	0	0	0
3	M	2	0	0	0	0
3	P	2	0	0	1	0
All	All	35615	0	32407	667	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 667 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:147:ILE:HG21	3:H:3221:HOH:O	1.16	1.33
1:E:158:THR:HG23	3:E:3234:HOH:O	1.21	1.30
1:G:184:LEU:HD22	3:G:3263:HOH:O	1.27	1.29
1:A:457:LEU:HD13	3:A:3253:HOH:O	1.11	1.27
1:F:65:ILE:HG22	3:F:3274:HOH:O	1.26	1.26

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/524 (99%)	500 (97%)	15 (3%)	2 (0%)	30	27
1	B	517/524 (99%)	504 (98%)	12 (2%)	1 (0%)	43	42
1	C	516/524 (98%)	497 (96%)	18 (4%)	1 (0%)	43	42
1	D	516/524 (98%)	501 (97%)	14 (3%)	1 (0%)	43	42
1	E	517/524 (99%)	500 (97%)	16 (3%)	1 (0%)	43	42
1	F	519/524 (99%)	503 (97%)	15 (3%)	1 (0%)	43	42
1	G	517/524 (99%)	503 (97%)	13 (2%)	1 (0%)	43	42
1	H	516/524 (98%)	498 (96%)	16 (3%)	2 (0%)	30	27
All	All	4135/4192 (99%)	4006 (97%)	119 (3%)	10 (0%)	43	42

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	251	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	251	PRO
1	G	251	PRO
1	A	251	PRO
1	A	394	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/458 (99%)	439 (97%)	14 (3%)	35	37
1	B	453/458 (99%)	445 (98%)	8 (2%)	51	58
1	C	452/458 (99%)	445 (98%)	7 (2%)	57	64
1	D	452/458 (99%)	436 (96%)	16 (4%)	32	32
1	E	453/458 (99%)	445 (98%)	8 (2%)	51	58
1	F	455/458 (99%)	444 (98%)	11 (2%)	43	47
1	G	453/458 (99%)	437 (96%)	16 (4%)	32	32
1	H	452/458 (99%)	431 (95%)	21 (5%)	24	22
All	All	3623/3664 (99%)	3522 (97%)	101 (3%)	39	41

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	329	VAL
1	G	221	ASP
1	H	484	ASP
1	F	347[B]	TYR
1	G	57	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 85 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	39	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	92	ASN
1	G	55	HIS
1	G	280	GLN
1	H	207	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	517/524 (98%)	0.19	7 (1%) 73 73	8, 22, 36, 51	2 (0%)
1	B	517/524 (98%)	0.46	26 (5%) 34 33	9, 28, 47, 61	2 (0%)
1	C	517/524 (98%)	0.47	18 (3%) 47 46	12, 29, 42, 53	1 (0%)
1	D	517/524 (98%)	0.49	12 (2%) 61 60	13, 30, 42, 53	1 (0%)
1	E	517/524 (98%)	0.28	7 (1%) 73 73	10, 25, 36, 44	2 (0%)
1	F	517/524 (98%)	0.40	16 (3%) 51 50	9, 25, 41, 57	4 (0%)
1	G	517/524 (98%)	0.71	37 (7%) 21 20	11, 30, 51, 64	2 (0%)
1	H	517/524 (98%)	1.42	140 (27%) 1 1	22, 38, 67, 79	1 (0%)
2	I	0/3	-	-	-	-
2	J	0/3	-	-	-	-
2	K	0/3	-	-	-	-
2	L	0/3	-	-	-	-
2	M	0/3	-	-	-	-
2	N	0/3	-	-	-	-
2	O	0/3	-	-	-	-
2	P	0/3	-	-	-	-
All	All	4136/4216 (98%)	0.55	263 (6%) 25 24	8, 28, 48, 79	15 (0%)

The worst 5 of 263 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	126[A]	SER	6.4
1	H	97	VAL	6.2
1	H	193	VAL	5.7
1	H	209	GLY	4.7
1	D	330	ASN	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.