



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 09:18 PM UTC

PDB ID : 6TC5 / pdb_00006tc5
Title : Phytochromobilin-adducted PAS-GAF bidomain of Sorghum bicolor phyB
Authors : Nagano, S.; Guan, K.; Shenkutie, S.M.; Hughes, J.E.
Deposited on : 2019-11-05
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	354	9% 77% 12% 11%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	315	2430	1545	435	433	17	0	1	1

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	111	MET	-	initiating methionine	UNP Q6S527
AAA	112	ALA	-	expression tag	UNP Q6S527
AAA	460	HIS	-	expression tag	UNP Q6S527
AAA	461	HIS	-	expression tag	UNP Q6S527
AAA	462	HIS	-	expression tag	UNP Q6S527
AAA	463	HIS	-	expression tag	UNP Q6S527
AAA	464	HIS	-	expression tag	UNP Q6S527

- Molecule 2 is 3-[5-[[[(3 {R},4 {R})-3-ethyl-4-methyl-5-oxidanylidene-3,4-dihydropyrrol-2-yl]methyl]-2-[[5-[(4-ethyl-3-methyl-5-oxidanylidene-pyrrol-2-yl)methyl]-3-(3-hydroxy-3-oxopropyl)-4-methyl-1 {H}-pyrrol-2-yl]methyl]-4-methyl-1 {H}-pyrrol-3-yl]propanoic acid (CCD ID: O6E) (formula: C₃₃H₃₈N₄O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 3 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	O	S	0	0
			4	2	1	1		
3	AAA	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	150	Total 150	O 150	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phytochrome



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	133.88Å 133.88Å 46.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.82 – 2.10 43.82 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (43.82-2.10) 99.0 (43.82-2.10)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.197 , 0.230 0.205 , 0.236	Depositor DCC
R_{free} test set	1401 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.042 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2631	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BME, O6E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.08	0/2485	1.41	7/3375 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	235	ARG	CB-CA-C	9.96	126.72	110.78
1	AAA	170	PRO	O-C-N	-6.35	118.39	121.31
1	AAA	224	ASP	CA-CB-CG	5.92	118.52	112.60
1	AAA	306	ARG	CG-CD-NE	-5.26	100.43	112.00
1	AAA	308	ASP	N-CA-C	-5.10	107.01	113.18
1	AAA	243	ILE	CA-C-N	5.04	126.99	120.44
1	AAA	243	ILE	C-N-CA	5.04	126.99	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2430	0	2435	21	0
2	AAA	43	0	0	0	0
3	AAA	8	0	10	0	0
4	AAA	150	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2631	0	2445	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:451:LEU:O	1:AAA:452:ALA:N	1.92	1.03
1:AAA:188:VAL:O	4:AAA:601:HOH:O	1.89	0.88
1:AAA:394:SER:N	1:AAA:409:ALA:O	2.20	0.74
1:AAA:147:GLU:OE2	4:AAA:602:HOH:O	2.14	0.65
1:AAA:291:TYR:CZ	1:AAA:299:GLY:HA3	2.34	0.63
1:AAA:192:ARG:CB	4:AAA:601:HOH:O	2.47	0.62
1:AAA:192:ARG:N	4:AAA:601:HOH:O	1.91	0.62
1:AAA:243:ILE:O	1:AAA:247:VAL:HG23	2.06	0.55
1:AAA:427:PRO:HB2	1:AAA:429:PRO:HD2	1.92	0.50
1:AAA:194:PHE:O	4:AAA:603:HOH:O	2.20	0.49
1:AAA:286:ASP:CG	1:AAA:307:ARG:HH21	2.21	0.48
1:AAA:237:GLU:HB3	1:AAA:241:LEU:HD23	1.95	0.48
1:AAA:286:ASP:CG	1:AAA:307:ARG:NH2	2.72	0.48
1:AAA:291:TYR:CE2	1:AAA:299:GLY:HA3	2.50	0.47
1:AAA:238:ASP:OD1	1:AAA:241:LEU:HB2	2.16	0.45
1:AAA:355:MET:HE1	1:AAA:359:LEU:HD13	2.00	0.44
1:AAA:425:CYS:SG	1:AAA:425:CYS:O	2.76	0.43
1:AAA:133:ALA:HB3	1:AAA:144:ALA:HB3	2.02	0.41
1:AAA:340:ALA:HA	1:AAA:424:ARG:O	2.21	0.40
1:AAA:385:SER:HA	1:AAA:418:CYS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	308/354 (87%)	291 (94%)	17 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	260/294 (88%)	249 (96%)	11 (4%)	26	28

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	117	SER
1	AAA	185	SER
1	AAA	213	SER
1	AAA	236	THR
1	AAA	237	GLU
1	AAA	255	ARG
1	AAA	263	LEU
1	AAA	333	GLN
1	AAA	337	ARG
1	AAA	356	SER
1	AAA	410	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	BME	AAA	502	1	3,3,3	0.13	0	2,2,2	0.49	0
3	BME	AAA	501	1	3,3,3	0.40	0	2,2,2	0.95	0
2	O6E	AAA	500	1	46,46,46	2.31	17 (36%)	62,67,67	1.82	16 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BME	AAA	502	1	-	1/1/1/1	-
3	BME	AAA	501	1	-	1/1/1/1	-
2	O6E	AAA	500	1	-	6/26/74/74	0/4/4/4

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	500	O6E	CAV-CBB	-5.09	1.37	1.45
2	AAA	500	O6E	CAG-CAB	5.02	1.54	1.30
2	AAA	500	O6E	CBD-CAX	4.67	1.56	1.52
2	AAA	500	O6E	CAW-CBC	-4.59	1.36	1.45
2	AAA	500	O6E	CBL-CBH	-4.48	1.36	1.45
2	AAA	500	O6E	CAB-CBI	-4.00	1.36	1.47
2	AAA	500	O6E	CAU-CBE	3.61	1.58	1.50
2	AAA	500	O6E	CBJ-CBN	3.13	1.57	1.50
2	AAA	500	O6E	OAZ-CAL	-2.91	1.21	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	500	O6E	CAX-NBP	-2.83	1.34	1.37
2	AAA	500	O6E	OBG-CAM	-2.78	1.21	1.30
2	AAA	500	O6E	CAI-CAM	2.75	1.56	1.50
2	AAA	500	O6E	CBI-CBM	-2.60	1.36	1.47
2	AAA	500	O6E	CBO-CBK	-2.58	1.37	1.42
2	AAA	500	O6E	CAR-CBH	2.57	1.56	1.50
2	AAA	500	O6E	CAY-CBE	-2.46	1.35	1.41
2	AAA	500	O6E	CAS-CBC	2.06	1.55	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	500	O6E	CBD-CAX-NBP	5.99	113.27	108.29
2	AAA	500	O6E	CAF-CAA-CBB	-4.12	101.15	112.53
2	AAA	500	O6E	OBQ-CAX-CBD	-3.81	123.14	126.17
2	AAA	500	O6E	CBI-CBM-NAJ	3.36	110.35	106.13
2	AAA	500	O6E	CAW-NAJ-CBM	-2.92	107.08	110.66
2	AAA	500	O6E	CAQ-CAY-CBE	-2.77	120.74	127.53
2	AAA	500	O6E	CAA-CBB-CBH	-2.74	122.73	127.87
2	AAA	500	O6E	CAQ-CAY-NAN	2.74	131.47	125.29
2	AAA	500	O6E	CAV-CBB-CBH	2.71	109.66	106.73
2	AAA	500	O6E	CBB-CAV-NAE	-2.59	106.38	110.04
2	AAA	500	O6E	CAO-CBO-NAN	2.41	130.73	125.29
2	AAA	500	O6E	CAO-CAV-NAE	2.40	129.64	124.60
2	AAA	500	O6E	CAP-CBL-CBH	2.26	130.67	124.87
2	AAA	500	O6E	CBI-CBC-CAW	2.25	110.43	107.92
2	AAA	500	O6E	CAO-CBO-CBK	-2.23	122.44	127.22
2	AAA	500	O6E	CBM-CBI-CBC	-2.16	105.15	107.92

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AAA	501	BME	O1-C1-C2-S2
3	AAA	502	BME	O1-C1-C2-S2
2	AAA	500	O6E	CAW-CAP-CBL-CBH
2	AAA	500	O6E	CAW-CAP-CBL-NAE
2	AAA	500	O6E	CBL-CAP-CAW-NAJ
2	AAA	500	O6E	CAA-CAF-CAL-OAZ
2	AAA	500	O6E	CAA-CAF-CAL-OBF
2	AAA	500	O6E	CAD-CAI-CAM-OBG

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	AAA	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	AAA	451:LEU	C	452:ALA	N	3.12

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	315/354 (88%)	0.30	31 (9%) 13 13	22, 36, 76, 107	1 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	394	SER	5.7
1	AAA	114	ALA	5.0
1	AAA	241	LEU	4.2
1	AAA	236	THR	4.0
1	AAA	243	ILE	3.9
1	AAA	239	PRO	3.9
1	AAA	329	PHE	3.8
1	AAA	409	ALA	3.7
1	AAA	436	PHE	3.6
1	AAA	451	LEU	3.6
1	AAA	452	ALA	3.6
1	AAA	410	MET	3.4
1	AAA	168	ALA	3.3
1	AAA	450	GLN	3.1
1	AAA	249	SER	3.1
1	AAA	248	GLN	2.7
1	AAA	235	ARG	2.6
1	AAA	155	LEU	2.5
1	AAA	242	SER	2.5
1	AAA	244	ALA	2.4
1	AAA	240	ALA	2.3
1	AAA	247	VAL	2.3
1	AAA	252	LEU	2.3
1	AAA	169	PRO	2.3
1	AAA	262	ALA	2.2
1	AAA	333	GLN	2.2
1	AAA	156	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	245	GLY	2.2
1	AAA	453	HIS	2.2
1	AAA	308	ASP	2.0
1	AAA	264	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BME	AAA	502	4/4	0.90	0.17	63,68,72,73	0
2	O6E	AAA	500	43/43	0.95	0.07	26,31,39,42	0
3	BME	AAA	501	4/4	0.97	0.08	40,41,43,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around O6E AAA 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.