



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:45 AM UTC

PDB ID : 5TDP / pdb_00005tdp
Title : Crystal structure of the Fab fragment of anti-HER2 antibody 4D5 with re-designed heavy and light chain interfaces
Authors : Yin, Y.; Carter, P.J.
Deposited on : 2016-09-19
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	27% 90% 9%
1	C	214	9% 93% 6%
2	B	223	6% 86% 7% 7%
2	D	223	3% 90% 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-HER2 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	3	0
			1654	1034	279	333	8			
1	C	213	Total	C	N	O	S	0	0	0
			1636	1025	276	329	6			

- Molecule 2 is a protein called anti-HER2 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	0	7	0
			1616	1018	273	315	10			
2	D	214	Total	C	N	O	S	0	6	0
			1648	1040	277	321	10			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	59	Total	O	0	0
			59	59		
4	B	134	Total	O	0	0
			134	134		
4	C	106	Total	O	0	0
			106	106		
4	D	180	Total	O	0	0
			180	180		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: anti-HER2 Fab Light Chain




- Molecule 1: anti-HER2 Fab Light Chain



- Molecule 2: anti-HER2 Fab Heavy Chain



- Molecule 2: anti-HER2 Fab Heavy Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.04Å 79.75Å 85.74Å 113.78° 93.07° 102.41°	Depositor
Resolution (Å)	43.70 – 1.72 43.70 – 1.72	Depositor EDS
% Data completeness (in resolution range)	95.0 (43.70-1.72) 95.1 (43.70-1.72)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 1.72Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.220 , 0.252 0.221 , 0.250	Depositor DCC
R_{free} test set	4732 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7039	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1690	0.76	2/2295 (0.1%)
1	C	0.41	0/1672	0.72	0/2271
2	B	0.48	0/1653	0.72	0/2253
2	D	0.49	0/1688	0.77	0/2301
All	All	0.44	0/6703	0.74	2/9120 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	VAL	CA-C-N	5.50	125.50	119.89
1	A	58	VAL	C-N-CA	5.50	125.50	119.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1654	0	1606	11	0
1	C	1636	0	1595	7	0
2	B	1616	0	1571	10	0
2	D	1648	0	1595	7	0
3	A	6	0	8	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	59	0	0	1	0
4	B	134	0	0	1	0
4	C	106	0	0	0	0
4	D	180	0	0	2	0
All	All	7039	0	6375	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:ASP:HA	4:A:401:HOH:O	1.82	0.80
1:A:119:PRO:HG2	3:A:301:GOL:H32	1.68	0.75
2:B:144:ILE:HG12	2:B:190:THR:HG22	1.75	0.69
1:A:209:PHE:HB2	3:A:301:GOL:H2	1.78	0.65
2:B:206[A]:ASN:HD21	2:B:208:LYS:HE3	1.60	0.65
2:D:144:ILE:HG12	2:D:190:THR:HG22	1.83	0.59
2:B:206[A]:ASN:ND2	2:B:208:LYS:HE3	2.19	0.56
1:C:142:ARG:CZ	1:C:163:VAL:HG21	2.40	0.51
2:D:124:LYS:HG2	4:D:324:HOH:O	2.11	0.50
1:A:176:PHE:CD2	2:B:173:SER:HB3	2.46	0.49
1:A:35:TRP:CZ3	1:A:88[B]:CYS:HB2	2.48	0.49
2:D:62:ASP:HA	2:D:65:LYS:HD3	1.97	0.47
2:D:123:THR:HG22	2:D:210:SER:HB3	1.95	0.47
1:C:4:MET:HE3	1:C:23:CYS:SG	2.55	0.46
2:B:106:ALA:N	4:B:305:HOH:O	2.48	0.46
1:C:163:VAL:HG22	1:C:175:LEU:HD12	1.98	0.46
2:B:12[A]:VAL:HG21	2:B:18:LEU:HD22	1.99	0.45
1:A:66:ARG:HG3	1:A:71:PHE:CE2	2.52	0.45
2:D:67:ARG:HG3	4:D:398:HOH:O	2.17	0.44
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.99	0.43
1:A:25:ALA:O	1:A:69:THR:OG1	2.36	0.43
1:A:3:GLN:H	1:A:26:SER:HG	1.65	0.43
1:C:107:LYS:HA	1:C:140:TYR:OH	2.19	0.43
2:B:83:MET:HE2	2:B:86:LEU:HD21	2.01	0.43
1:A:3:GLN:HB3	1:A:26:SER:HB3	2.00	0.42
2:B:4:LEU:HD22	2:B:22:CYS:SG	2.60	0.42
1:A:142:ARG:CZ	1:A:163:VAL:HG21	2.49	0.42
2:B:196:LEU:HD23	2:B:196:LEU:HA	1.91	0.42
1:C:37:GLN:HB2	1:C:47:LEU:HD11	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:213:LYS:HB2	2:D:213:LYS:HE2	1.76	0.42
2:B:30:LYS:HE3	2:B:30:LYS:HB2	1.86	0.41
2:D:12[B]:VAL:HG21	2:D:18:LEU:HD22	2.02	0.41
1:C:167:ASP:OD2	1:C:169:LYS:N	2.54	0.40
1:C:140:TYR:CG	1:C:141:PRO:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/214 (100%)	207 (97%)	7 (3%)	0	100	100
1	C	211/214 (99%)	204 (97%)	7 (3%)	0	100	100
2	B	208/223 (93%)	205 (99%)	3 (1%)	0	100	100
2	D	216/223 (97%)	214 (99%)	2 (1%)	0	100	100
All	All	849/874 (97%)	830 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/187 (101%)	186 (98%)	3 (2%)	55	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	186/187 (100%)	185 (100%)	1 (0%)	81	71
2	B	179/183 (98%)	177 (99%)	2 (1%)	65	50
2	D	181/183 (99%)	178 (98%)	3 (2%)	53	31
All	All	735/740 (99%)	726 (99%)	9 (1%)	61	46

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	27	GLN
1	A	171	SER
2	B	50	ARG
2	B	157	VAL
1	C	137	ASN
2	D	157	VAL
2	D	176	VAL
2	D	206	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	ASN
1	A	138	ASN
1	A	147	GLN
1	C	147	GLN
2	D	171	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	301	-	5,5,5	0.24	0	5,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	GOL	O1-C1-C2-C3
3	A	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	GOL	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/214 (99%)	1.41	58 (27%) 1 3	13, 37, 60, 86	3 (1%)
1	C	213/214 (99%)	0.82	20 (9%) 14 19	14, 28, 45, 66	0
2	B	208/223 (93%)	0.39	13 (6%) 26 31	9, 20, 39, 54	7 (3%)
2	D	214/223 (95%)	0.35	6 (2%) 55 63	7, 19, 33, 47	6 (2%)
All	All	848/874 (97%)	0.74	97 (11%) 10 14	7, 25, 50, 86	16 (1%)

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	83	PHE	7.9
1	C	83	PHE	7.7
1	A	154	LEU	6.1
1	A	108	ARG	5.4
1	A	15	VAL	5.4
1	A	106	ILE	5.3
1	A	80	PRO	5.1
1	C	169	LYS	4.4
1	A	157	GLY	4.1
1	A	78	LEU	4.1
2	B	134	SER	4.0
1	A	84	ALA	4.0
1	C	93	THR	3.9
1	C	108	ARG	3.9
2	D	32	THR	3.8
1	C	105	GLU	3.8
1	C	69	THR	3.8
1	A	59	PRO	3.6
1	A	156	SER	3.6
1	A	142	ARG	3.6
1	A	16	GLY	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	220	PRO	3.5
2	D	219	GLU	3.4
1	A	2	ILE	3.4
2	B	99	TRP	3.3
1	A	69	THR	3.3
1	A	77	SER	3.3
1	A	169	LYS	3.2
1	A	92	TYR	3.1
1	A	147	GLN	3.1
1	A	60	SER	3.1
2	B	197	GLY	3.1
2	B	142	THR	3.1
2	D	176	VAL	3.1
1	A	25	ALA	3.1
1	A	54	LEU	3.0
1	C	3	GLN	3.0
2	B	199	GLN	3.0
1	C	2	ILE	3.0
2	B	190	THR	2.8
1	A	61	ARG	2.8
1	A	17	ASP	2.8
1	A	126	LYS	2.7
1	A	205	VAL	2.7
2	B	112	GLN	2.7
1	A	62	PHE	2.7
2	D	99	TRP	2.7
2	D	142	THR	2.7
1	A	168	SER	2.7
1	A	214	CYS	2.7
1	A	76	SER	2.6
1	A	194[A]	CYS	2.6
1	A	14	SER	2.6
2	B	194	SER	2.6
1	C	168	SER	2.6
1	C	80	PRO	2.6
1	A	58	VAL	2.5
1	C	106	ILE	2.5
2	B	141	GLY	2.5
1	A	109	THR	2.5
2	B	43	LYS	2.5
1	A	79	GLN	2.5
1	C	214	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	129	THR	2.5
1	A	100	GLN	2.4
1	C	94	THR	2.4
1	C	142	ARG	2.4
1	A	73	LEU	2.4
1	A	3	GLN	2.4
1	A	110	VAL	2.4
1	A	49	TYR	2.4
1	A	5	THR	2.3
1	A	107	LYS	2.3
2	D	198	THR	2.3
2	B	3[A]	GLN	2.3
1	A	85	THR	2.3
1	A	53	PHE	2.3
1	A	19	VAL	2.3
1	C	15	VAL	2.3
1	C	145	LYS	2.2
1	A	94	THR	2.2
1	A	129	THR	2.2
1	A	26	SER	2.2
1	C	12	SER	2.2
1	A	201	LEU	2.2
1	C	60	SER	2.2
1	A	23	CYS	2.2
1	A	202	SER	2.1
1	A	81	GLU	2.1
1	C	92	TYR	2.1
1	A	13	ALA	2.1
1	A	144	ALA	2.1
2	B	30	LYS	2.1
1	A	42	LYS	2.1
1	A	197	THR	2.1
1	A	95	PRO	2.0
1	A	146	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	301	6/6	0.81	0.15	35,46,46,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.