



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 07:24 PM UTC

PDB ID : 7TDV / pdb_00007tdv
Title : Crystal structure of S. aureus glutamine synthetase in Met-Sox-P/ADP transition state complex
Authors : Schumacher, M.A.
Deposited on : 2022-01-03
Resolution : 2.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2995 (2.94-2.90)
Clashscore	190562	3213 (2.94-2.90)
Ramachandran outliers	187476	3128 (2.94-2.90)
Sidechain outliers	187428	3130 (2.94-2.90)
RSRZ outliers	180081	2995 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	 66% 32% .
1	B	449	 69% 29% .
1	C	449	 69% 29% .
1	D	449	 56% 42% .
1	E	449	 60% 37% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	449	 A horizontal bar chart showing the quality of the chain. The bar is divided into two segments: a green segment on the left representing 66% and a yellow segment on the right representing 33%. A small grey square is at the far right end of the bar, and a small black dot is at the very end of the bar.

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 21932 atoms, of which 132 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamine synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3531	2249	591	674	17			
1	C	443	Total	C	N	O	S	0	0	0
			3546	2258	596	675	17			
1	D	442	Total	C	N	O	S	0	0	0
			3537	2252	591	677	17			
1	E	442	Total	C	N	O	S	1	0	0
			3533	2251	592	673	17			
1	B	443	Total	C	N	O	S	0	0	0
			3536	2252	592	675	17			
1	H	442	Total	C	N	O	S	0	0	0
			3533	2251	592	673	17			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP E3VXC2
A	-1	SER	-	expression tag	UNP E3VXC2
A	0	HIS	-	expression tag	UNP E3VXC2
A	15	VAL	ALA	conflict	UNP E3VXC2
C	-2	GLY	-	expression tag	UNP E3VXC2
C	-1	SER	-	expression tag	UNP E3VXC2
C	0	HIS	-	expression tag	UNP E3VXC2
C	15	VAL	ALA	conflict	UNP E3VXC2
D	-2	GLY	-	expression tag	UNP E3VXC2
D	-1	SER	-	expression tag	UNP E3VXC2
D	0	HIS	-	expression tag	UNP E3VXC2
D	15	VAL	ALA	conflict	UNP E3VXC2
E	-2	GLY	-	expression tag	UNP E3VXC2
E	-1	SER	-	expression tag	UNP E3VXC2
E	0	HIS	-	expression tag	UNP E3VXC2
E	15	VAL	ALA	conflict	UNP E3VXC2
B	-2	GLY	-	expression tag	UNP E3VXC2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	SER	-	expression tag	UNP E3VXC2
B	0	HIS	-	expression tag	UNP E3VXC2
B	15	VAL	ALA	conflict	UNP E3VXC2
H	-2	GLY	-	expression tag	UNP E3VXC2
H	-1	SER	-	expression tag	UNP E3VXC2
H	0	HIS	-	expression tag	UNP E3VXC2
H	15	VAL	ALA	conflict	UNP E3VXC2

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	C	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	D	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	E	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	B	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	H	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

- Molecule 3 is L-METHIONINE-S-SULFOXIMINE PHOSPHATE (CCD ID: P3S) (formula: $C_5H_{13}N_2O_6PS$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		
3	C	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		
3	D	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		
3	E	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		
3	B	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		
3	H	1	Total	C	H	N	O	P	S	0	0
			25	5	10	2	6	1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		
4	C	5	Total	Mg	0	0
			5	5		
4	D	3	Total	Mg	0	0
			3	3		
4	E	4	Total	Mg	0	0
			4	4		
4	B	5	Total	Mg	0	0
			5	5		
4	H	2	Total	Mg	0	0
			2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	50	Total	O	0	0
			50	50		
6	C	77	Total	O	0	0
			77	77		
6	D	30	Total	O	0	0
			30	30		
6	E	26	Total	O	0	0
			26	26		
6	B	56	Total	O	0	0
			56	56		
6	H	67	Total	O	0	0
			67	67		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamine synthetase




• Molecule 1: Glutamine synthetase




• Molecule 1: Glutamine synthetase

Chain D:  56% 42%



• Molecule 1: Glutamine synthetase


Chain E:  60% 37%




• Molecule 1: Glutamine synthetase

Chain B:  69% 29%





● Molecule 1: Glutamine synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	154.62Å 154.62Å 299.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.89 – 2.92 48.89 – 2.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.89-2.92) 99.9 (48.89-2.92)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.198 , 0.256 0.201 , 0.256	Depositor DCC
R_{free} test set	2000 reflections (2.37%)	wwPDB-VP
Wilson B-factor (Å ²)	69.5	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21932	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ADP, MG, P3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	2/3613 (0.1%)	0.73	3/4894 (0.1%)
1	B	0.53	0/3618	0.70	0/4901
1	C	0.55	0/3628	0.75	4/4912 (0.1%)
1	D	0.59	2/3619 (0.1%)	0.77	5/4901 (0.1%)
1	E	0.49	1/3615 (0.0%)	0.72	5/4895 (0.1%)
1	H	0.55	1/3615 (0.0%)	0.68	0/4895
All	All	0.54	6/21708 (0.0%)	0.73	17/29398 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	135	PRO	N-CA	17.17	1.69	1.47
1	A	324	VAL	C-N	13.15	1.50	1.33
1	E	302	VAL	C-N	11.66	1.49	1.33
1	D	319	SER	C-N	7.25	1.50	1.33
1	H	325	PRO	C-O	-5.79	1.16	1.23

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	134	GLU	CA-C-N	14.81	138.35	119.84
1	D	134	GLU	C-N-CA	14.81	138.35	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	302	VAL	CA-C-N	12.22	132.37	119.78
1	E	302	VAL	C-N-CA	12.22	132.37	119.78
1	A	324	VAL	CA-C-N	11.63	132.36	119.92

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	13	LYS	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3531	0	3431	140	0
1	B	3536	0	3433	132	2
1	C	3546	0	3455	109	2
1	D	3537	0	3432	188	0
1	E	3533	0	3437	172	1
1	H	3533	0	3437	158	2
2	A	27	12	12	3	0
2	B	27	12	12	2	0
2	C	27	12	12	0	0
2	D	27	12	12	4	0
2	E	27	12	12	0	0
2	H	27	12	12	2	0
3	A	15	10	10	5	0
3	B	15	10	10	1	0
3	C	15	10	10	1	0
3	D	15	10	10	2	0
3	E	15	10	10	2	0
3	H	15	10	10	3	0
4	A	2	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	3	0	0	0	0
4	E	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	2	0	0	0	0
5	H	5	0	0	1	0
6	A	50	0	0	1	0
6	B	56	0	0	0	0
6	C	77	0	0	3	0
6	D	30	0	0	1	0
6	E	26	0	0	0	0
6	H	67	0	0	2	0
All	All	21800	132	20757	853	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 853 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:135:PRO:CA	1:D:135:PRO:N	1.69	1.36
1:C:255:GLY:O	1:C:256:LYS:HD3	1.07	1.24
1:C:7:THR:HG22	1:C:10:ASP:OD2	1.40	1.19
1:E:172:ARG:O	1:E:175:VAL:HG23	1.46	1.14
1:D:136:GLU:OE1	3:D:502:P3S:N	1.83	1.10

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:LYS:NZ	1:B:438:GLU:OE2[7_465]	1.39	0.81
1:E:299:LYS:NZ	1:E:438:GLU:OE1[7_465]	1.70	0.50
1:C:442:TYR:OH	1:H:426:GLU:OE1[7_465]	2.02	0.18
1:C:426:GLU:OE2	1:H:442:TYR:OH[7_465]	2.13	0.07
1:B:299:LYS:CE	1:B:438:GLU:OE2[7_465]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/449 (98%)	416 (94%)	24 (6%)	0	100	100
1	B	441/449 (98%)	412 (93%)	29 (7%)	0	100	100
1	C	441/449 (98%)	417 (95%)	24 (5%)	0	100	100
1	D	440/449 (98%)	421 (96%)	19 (4%)	0	100	100
1	E	440/449 (98%)	415 (94%)	25 (6%)	0	100	100
1	H	440/449 (98%)	418 (95%)	22 (5%)	0	100	100
All	All	2642/2694 (98%)	2499 (95%)	143 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/391 (97%)	381 (100%)	0	100	100
1	B	381/391 (97%)	379 (100%)	2 (0%)	81	93
1	C	383/391 (98%)	381 (100%)	2 (0%)	81	93
1	D	382/391 (98%)	380 (100%)	2 (0%)	81	93
1	E	381/391 (97%)	376 (99%)	5 (1%)	61	84
1	H	381/391 (97%)	380 (100%)	1 (0%)	86	95
All	All	2289/2346 (98%)	2277 (100%)	12 (0%)	81	93

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	423	LYS
1	E	436	GLU
1	H	299	LYS
1	B	90	LYS
1	D	319	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	283	ASN
1	B	242	ASN
1	B	36	ASN
1	B	264	ASN
1	C	413	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 34 ligands modelled in this entry, 21 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	P3S	H	502	4	11,14,14	2.24	4 (36%)	9,21,21	6.66	4 (44%)
2	ADP	C	501	4	28,29,29	1.44	5 (17%)	43,45,45	2.42	21 (48%)
3	P3S	C	502	4	11,14,14	1.87	2 (18%)	9,21,21	5.86	3 (33%)
2	ADP	B	501	4	28,29,29	1.55	4 (14%)	43,45,45	2.35	18 (41%)
3	P3S	D	502	4	11,14,14	1.99	4 (36%)	9,21,21	6.14	4 (44%)
2	ADP	D	501	4	28,29,29	1.57	6 (21%)	43,45,45	1.87	13 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	P3S	E	502	4	11,14,14	2.22	3 (27%)	9,21,21	5.58	2 (22%)
2	ADP	H	501	4	28,29,29	1.75	5 (17%)	43,45,45	2.49	17 (39%)
3	P3S	A	502	4	11,14,14	1.96	4 (36%)	9,21,21	6.61	3 (33%)
2	ADP	A	501	4	28,29,29	1.74	6 (21%)	43,45,45	2.02	12 (27%)
2	ADP	E	501	4	28,29,29	1.53	6 (21%)	43,45,45	1.98	13 (30%)
5	SO4	H	505	-	4,4,4	0.25	0	6,6,6	0.13	0
3	P3S	B	502	4	11,14,14	2.34	3 (27%)	9,21,21	6.53	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	P3S	H	502	4	-	6/9/16/16	-
2	ADP	C	501	4	-	4/16/32/32	0/3/3/3
3	P3S	C	502	4	-	5/9/16/16	-
2	ADP	B	501	4	-	6/16/32/32	0/3/3/3
3	P3S	D	502	4	-	5/9/16/16	-
2	ADP	D	501	4	-	3/16/32/32	0/3/3/3
3	P3S	E	502	4	-	3/9/16/16	-
2	ADP	H	501	4	-	2/16/32/32	0/3/3/3
3	P3S	A	502	4	-	5/9/16/16	-
2	ADP	A	501	4	-	2/16/32/32	0/3/3/3
2	ADP	E	501	4	-	4/16/32/32	0/3/3/3
3	P3S	B	502	4	-	8/9/16/16	-

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ADP	PA-O3A	5.46	1.65	1.59
2	H	501	ADP	PA-O3A	5.15	1.65	1.59
2	D	501	ADP	C5-C4	4.57	1.47	1.39
2	H	501	ADP	C5-C4	4.50	1.47	1.39
3	H	502	P3S	PA-NE	4.50	1.74	1.58

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	502	P3S	OE-SD-CE	-17.49	81.48	109.21
3	B	502	P3S	OE-SD-CE	-17.00	82.26	109.21
3	A	502	P3S	OE-SD-CE	-16.85	82.49	109.21
3	D	502	P3S	OE-SD-CE	-16.21	83.51	109.21
3	C	502	P3S	OE-SD-CE	-15.06	85.32	109.21

There are no chirality outliers.

5 of 53 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	501	ADP	PA-O3A-PB-O3B
2	E	501	ADP	PA-O3A-PB-O2B
2	E	501	ADP	O4'-C4'-C5'-O5'
2	B	501	ADP	PA-O3A-PB-O2B
2	B	501	ADP	C5'-O5'-PA-O2A


There are no ring outliers.

11 monomers are involved in 26 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	502	P3S	3	0
3	C	502	P3S	1	0
2	B	501	ADP	2	0
3	D	502	P3S	2	0
2	D	501	ADP	4	0
3	E	502	P3S	2	0
2	H	501	ADP	2	0
3	A	502	P3S	5	0
2	A	501	ADP	3	0
5	H	505	SO4	1	0
3	B	502	P3S	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.




Ligand P3S C 502




Ligand ADP B 501




Ligand P3S D 502




Ligand ADP D 501




Ligand P3S E 502




Ligand ADP H 501




Ligand P3S A 502



Ligand ADP A 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/449 (98%)	-0.32	2 (0%) 87 83	47, 68, 96, 106	0
1	B	443/449 (98%)	-0.39	1 (0%) 91 89	41, 62, 92, 111	0
1	C	443/449 (98%)	-0.42	0 100 100	42, 59, 87, 105	0
1	D	442/449 (98%)	-0.19	1 (0%) 91 89	47, 74, 102, 112	0
1	E	442/449 (98%)	-0.14	1 (0%) 91 89	51, 79, 103, 122	1 (0%)
1	H	442/449 (98%)	-0.37	0 100 100	40, 61, 88, 102	0
All	All	2654/2694 (98%)	-0.30	5 (0%) 91 89	40, 67, 96, 122	1 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	GLU	3.1
1	E	426	GLU	3.1
1	A	86	ALA	2.7
1	B	86	ALA	2.5
1	D	124	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	D	505	1/1	0.45	0.30	104,104,104,104	0
4	MG	E	506	1/1	0.56	0.32	89,89,89,89	0
4	MG	C	507	1/1	0.72	0.18	81,81,81,81	0
4	MG	B	507	1/1	0.75	0.32	82,82,82,82	0
4	MG	E	505	1/1	0.80	0.21	73,73,73,73	0
5	SO4	H	505	5/5	0.87	0.15	187,187,188,188	0
4	MG	H	503	1/1	0.90	0.18	43,43,43,43	0
4	MG	D	503	1/1	0.92	0.12	57,57,57,57	0
4	MG	C	506	1/1	0.92	0.07	90,90,90,90	0
4	MG	B	505	1/1	0.92	0.06	85,85,85,85	0
3	P3S	D	502	15/15	0.93	0.09	62,72,86,86	0
4	MG	A	503	1/1	0.94	0.17	61,61,61,61	0
4	MG	C	503	1/1	0.94	0.12	48,48,48,48	0
2	ADP	D	501	27/27	0.94	0.08	59,78,96,101	0
2	ADP	A	501	27/27	0.95	0.08	66,75,91,92	0
3	P3S	E	502	15/15	0.95	0.09	65,78,93,95	0
3	P3S	H	502	15/15	0.95	0.10	52,64,79,82	0
2	ADP	E	501	27/27	0.95	0.09	65,77,96,97	0
2	ADP	H	501	27/27	0.95	0.08	43,56,71,77	0
3	P3S	A	502	15/15	0.96	0.07	61,67,81,81	0
3	P3S	C	502	15/15	0.96	0.07	44,48,58,58	0
2	ADP	B	501	27/27	0.96	0.06	49,57,71,73	0
4	MG	E	503	1/1	0.96	0.13	63,63,63,63	0
2	ADP	C	501	27/27	0.96	0.06	45,49,60,61	0
3	P3S	B	502	15/15	0.97	0.07	53,67,79,82	0
4	MG	B	503	1/1	0.98	0.10	48,48,48,48	0
4	MG	B	504	1/1	0.99	0.04	51,51,51,51	0
4	MG	E	504	1/1	0.99	0.03	66,66,66,66	0
4	MG	B	506	1/1	0.99	0.03	53,53,53,53	0
4	MG	D	504	1/1	0.99	0.02	60,60,60,60	0
4	MG	C	505	1/1	0.99	0.06	47,47,47,47	0
4	MG	H	504	1/1	0.99	0.06	53,53,53,53	0
4	MG	C	504	1/1	0.99	0.09	49,49,49,49	0
4	MG	A	504	1/1	1.00	0.04	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P3S D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around P3S E 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around P3S H 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP E 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP H 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around P3S A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around P3S C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around P3S B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.