



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:59 AM UTC

PDB ID : 3TE5 / pdb_00003te5
Title : structure of the regulatory fragment of sacchromyces cerevisiae ampk in complex with NADH
Authors : Mayer, F.V.; Heath, R.; Underwood, E.; Sanders, M.J.; Carmena, D.; McCartney, R.; Leiper, F.C.; Xiao, B.; Jing, C.; Walker, P.A.; Haire, L.F.; Ogrodowicz, R.; Martin, S.R.; Schmidt, M.C.; Gamblin, S.J.; Carling, D.
Deposited on : 2011-08-12
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	179	11% 66% 10% • 23%
2	B	113	27% 79% 12% • 9%
3	C	323	7% 83% 12% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbon catabolite-derepressing protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	0	0
			1102	722	184	191	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	455	GLY	-	expression tag	UNP P06782
A	456	PRO	-	expression tag	UNP P06782

- Molecule 2 is a protein called SNF1 protein kinase subunit beta-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	103	Total	C	N	O	S	0	0	0
			825	536	133	154	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	303	MET	-	expression tag	UNP P34164

- Molecule 3 is a protein called Nuclear protein SNF4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	313	Total	C	N	O	S	0	0	0
			2438	1555	412	461	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	MET	-	expression tag	UNP P12904
C	2	ALA	-	expression tag	UNP P12904

- [illegible]


- Molecule 5 is water.

WORLDWIDE
 **PDB**
PROTEIN DATA BANK


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbon catabolite-derepressing protein kinase



- Molecule 2: SNF1 protein kinase subunit beta-2



- Molecule 3: Nuclear protein SNF4



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	95.73Å 240.24Å 79.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.50 25.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.00-2.50) 99.7 (25.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.241 , 0.281 0.234 , 0.272	Depositor DCC
R_{free} test set	1617 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	48.1	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4623	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1131	0.73	0/1533
2	B	0.47	0/848	0.80	1/1167 (0.1%)
3	C	0.45	0/2472	0.78	0/3353
All	All	0.44	0/4451	0.77	1/6053 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	336	SER	N-CA-C	5.51	118.04	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1102	0	1113	16	0
2	B	825	0	807	8	0
3	C	2438	0	2487	31	0
4	C	44	0	27	3	0
5	A	38	0	0	1	0
5	B	40	0	0	0	0
5	C	136	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4623	0	4434	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:16:GLN:HG3	3:C:187:ILE:HG13	1.72	0.71
3:C:242:VAL:O	3:C:246:ILE:HG22	1.98	0.63
3:C:93:PHE:HD1	3:C:254:LEU:HD13	1.64	0.62
1:A:611:SER:HB2	3:C:161:ILE:CG2	2.35	0.56
3:C:250:ILE:CG2	3:C:253:ASP:HB2	2.37	0.54
3:C:253:ASP:HA	3:C:256:LEU:HD13	1.91	0.52
1:A:505:TRP:HH2	2:B:392:CYS:HG	1.58	0.52
3:C:93:PHE:CD1	3:C:254:LEU:HD13	2.45	0.52
1:A:575:ILE:HD11	1:A:581:LEU:HB2	1.93	0.51
3:C:16:GLN:HG2	3:C:185:LEU:HA	1.93	0.51
2:B:402:LYS:HD2	3:C:38:VAL:HA	1.94	0.50
2:B:335:THR:HG23	2:B:337:TRP:CD1	2.47	0.49
3:C:82:THR:O	3:C:86:ILE:HG12	2.12	0.49
3:C:149:PRO:HB2	3:C:151:ILE:CD1	2.42	0.49
3:C:250:ILE:HG23	3:C:253:ASP:HB2	1.94	0.49
3:C:19:ALA:O	3:C:23:ILE:HG12	2.12	0.49
1:A:483:LEU:HG	1:A:490:ALA:HB1	1.94	0.49
2:B:410:THR:HG22	2:B:411:PRO:HD2	1.96	0.48
1:A:493:ILE:CG2	3:C:263:MET:HB2	2.45	0.47
3:C:214:GLN:O	3:C:218:GLN:HB2	2.14	0.47
1:A:610:PHE:N	5:A:38:HOH:O	2.47	0.47
3:C:140:MET:HE1	3:C:171:ILE:HD12	1.97	0.47
1:A:623:ILE:HD11	2:B:381:LEU:HD21	1.98	0.46
3:C:286:ASP:OD1	3:C:289:ARG:NH2	2.46	0.46
2:B:402:LYS:HG2	3:C:33:TYR:CE1	2.51	0.45
1:A:610:PHE:HZ	2:B:379:ASN:OD1	2.00	0.45
3:C:246:ILE:HA	5:C:435:HOH:O	2.17	0.45
1:A:522:ILE:HA	1:A:626:LEU:HD21	1.99	0.45
1:A:547:ARG:HG2	1:A:565:LEU:HB3	1.99	0.44
1:A:617:HIS:O	1:A:620:THR:HB	2.18	0.44
3:C:284:ILE:HG23	3:C:296:PHE:CE2	2.53	0.44
3:C:16:GLN:NE2	5:C:409:HOH:O	2.51	0.44
1:A:514:TYR:HB3	1:A:515:PRO:HD2	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:168:GLN:HE22	3:C:312:SER:HB3	1.82	0.43
1:A:611:SER:HB2	3:C:161:ILE:HG21	2.01	0.43
1:A:611:SER:HB2	3:C:161:ILE:HG22	2.01	0.42
3:C:254:LEU:HD12	3:C:255:SER:N	2.35	0.41
1:A:624:MET:HE1	2:B:386:ILE:HD11	2.02	0.41
3:C:312:SER:HB3	4:C:324:NAI:H52N	2.02	0.41
3:C:221:VAL:HG12	4:C:324:NAI:H51A	2.02	0.41
3:C:86:ILE:HG23	3:C:213:ILE:HG23	2.02	0.41
3:C:262:LEU:O	3:C:265:ARG:HB2	2.20	0.41
4:C:324:NAI:H8A	4:C:324:NAI:O2A	2.20	0.41
1:A:482:MET:HB3	3:C:263:MET:HE1	2.02	0.41
3:C:45:LEU:HD21	3:C:57:VAL:HG11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/179 (74%)	128 (97%)	4 (3%)	0	100	100
2	B	99/113 (88%)	96 (97%)	3 (3%)	0	100	100
3	C	309/323 (96%)	303 (98%)	5 (2%)	1 (0%)	36	55
All	All	540/615 (88%)	527 (98%)	12 (2%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	249	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/162 (74%)	118 (99%)	1 (1%)	73	88
2	B	94/108 (87%)	89 (95%)	5 (5%)	20	42
3	C	277/297 (93%)	261 (94%)	16 (6%)	18	38
All	All	490/567 (86%)	468 (96%)	22 (4%)	24	49

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	620	THR
2	B	306	THR
2	B	351	ILE
2	B	359	GLN
2	B	401	GLN
2	B	410	THR
3	C	16	GLN
3	C	21	GLU
3	C	65	SER
3	C	110	LEU
3	C	130	SER
3	C	140	MET
3	C	151	ILE
3	C	170	ARG
3	C	204	GLN
3	C	217	THR
3	C	218	GLN
3	C	242	VAL
3	C	250	ILE
3	C	255	SER
3	C	263	MET
3	C	266	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	485	GLN
3	C	60	GLN
3	C	61	ASN
3	C	168	GLN
3	C	183	HIS
3	C	218	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAI	C	324	-	47,48,48	1.73	12 (25%)	64,73,73	1.73	12 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAI	C	324	-	-	9/29/72/72	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	324	NAI	C4N-C3N	-4.92	1.40	1.50
4	C	324	NAI	C7N-C3N	-3.65	1.40	1.48
4	C	324	NAI	C2A-N1A	3.58	1.40	1.33
4	C	324	NAI	C4N-C5N	-3.33	1.40	1.49
4	C	324	NAI	C2A-N3A	3.27	1.39	1.33
4	C	324	NAI	C5A-N7A	-2.77	1.34	1.39
4	C	324	NAI	C5A-C4A	-2.64	1.34	1.39
4	C	324	NAI	C5A-C6A	-2.47	1.34	1.41
4	C	324	NAI	C6N-C5N	2.33	1.40	1.33
4	C	324	NAI	C8A-N9A	-2.17	1.33	1.37
4	C	324	NAI	C4A-N9A	-2.14	1.33	1.37
4	C	324	NAI	PA-O3	2.06	1.61	1.59

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	324	NAI	N3A-C2A-N1A	-7.31	117.52	128.58
4	C	324	NAI	N9A-C8A-N7A	-4.07	108.16	113.94
4	C	324	NAI	C5A-C4A-N3A	-3.96	121.26	126.72
4	C	324	NAI	C2A-N3A-C4A	3.36	120.03	111.83
4	C	324	NAI	O4D-C1D-N1N	3.19	114.16	108.08
4	C	324	NAI	C5A-N7A-C8A	2.85	107.92	103.45
4	C	324	NAI	C5A-C6A-N6A	-2.62	116.80	123.29
4	C	324	NAI	C4A-N9A-C8A	2.35	108.21	105.74
4	C	324	NAI	C4A-C5A-N7A	-2.33	107.92	110.58
4	C	324	NAI	C4A-N9A-C1B	-2.29	121.28	126.63
4	C	324	NAI	N3A-C4A-N9A	2.22	130.94	127.17
4	C	324	NAI	O4D-C1D-C2D	-2.03	102.26	106.62

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	324	NAI	C5B-O5B-PA-O1A
4	C	324	NAI	C5D-O5D-PN-O2N
4	C	324	NAI	O4B-C4B-C5B-O5B
4	C	324	NAI	O4D-C1D-N1N-C2N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	C	324	NAI	C2B-C1B-N9A-C8A
4	C	324	NAI	C3B-C4B-C5B-O5B
4	C	324	NAI	C5D-O5D-PN-O3
4	C	324	NAI	C2B-C1B-N9A-C4A
4	C	324	NAI	O4B-C1B-N9A-C8A

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	324	NAI	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/179 (77%)	0.82	19 (13%) 6 5	32, 58, 77, 78	0
2	B	103/113 (91%)	1.37	30 (29%) 1 1	34, 60, 86, 89	0
3	C	313/323 (96%)	0.40	21 (6%) 24 21	26, 40, 60, 72	0
All	All	554/615 (90%)	0.68	70 (12%) 8 6	26, 45, 77, 89	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	7	ASP	5.4
2	B	368	GLY	5.0
3	C	248	GLY	5.0
2	B	361	GLN	4.9
2	B	369	ALA	4.7
2	B	359	GLN	4.6
2	B	355	TYR	4.5
2	B	356	TYR	4.5
1	A	562	ILE	4.2
3	C	251	TYR	4.1
2	B	326	LEU	4.1
2	B	358	THR	4.0
2	B	371	PRO	4.0
2	B	352	LEU	3.8
1	A	498	THR	3.6
3	C	249	GLY	3.6
3	C	254	LEU	3.5
1	A	497	VAL	3.4
2	B	357	ALA	3.4
1	A	488	PRO	3.4
3	C	155	GLU	3.4
1	A	577	THR	3.3
2	B	351	ILE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	253	ASP	3.3
3	C	156	GLU	3.3
1	A	487	SER	3.2
2	B	364	GLU	3.2
1	A	491	SER	3.2
2	B	350	VAL	3.1
3	C	9	GLN	3.0
2	B	365	ASN	3.0
2	B	335	THR	3.0
3	C	41	ARG	3.0
3	C	124	THR	2.9
2	B	362	PHE	2.9
2	B	303	MET	2.9
3	C	154	ASP	2.8
3	C	14	ILE	2.8
1	A	550	TYR	2.8
1	A	464	ASP	2.8
1	A	496	LEU	2.7
3	C	255	SER	2.7
2	B	366	ASN	2.6
3	C	93	PHE	2.5
1	A	502	LYS	2.5
2	B	354	LYS	2.5
1	A	501	SER	2.5
3	C	120	ASP	2.5
2	B	337	TRP	2.5
2	B	353	ASN	2.5
2	B	349	ASN	2.4
1	A	486	GLY	2.4
3	C	153	GLN	2.4
3	C	250	ILE	2.4
3	C	21	GLU	2.4
3	C	92	TYR	2.3
1	A	628	VAL	2.3
3	C	158	HIS	2.2
2	B	370	LEU	2.2
2	B	363	ASN	2.2
2	B	347	LEU	2.2
2	B	372	ILE	2.2
1	A	495	PRO	2.2
2	B	339	THR	2.2
1	A	490	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	412	ILE	2.0
1	A	485	GLN	2.0
1	A	493	ILE	2.0
2	B	309	ILE	2.0
1	A	563	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAI	C	324	44/44	0.93	0.09	40,42,43,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAI C 324:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.