



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:07 AM UTC

PDB ID : 5TE7 / pdb_00005te7
Title : Crystal Structure of Broadly Neutralizing VRC01-class Antibody N6 in Complex with HIV-1 Clade C Strain DU172.17 gp120 Core
Authors : Zhou, T.; Kwong, P.D.
Deposited on : 2016-09-20
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	3% 98% .
2	L	210	% 98% .
3	G	350	9% 88% 8% .

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11993 atoms, of which 5532 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of N6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	225	Total	C	H	N	O	S	0	0	0
			3347	1079	1644	296	321	7			


- Molecule 2 is a protein called Light chain of N6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	210	Total	C	H	N	O	S	0	0	0
			3193	1018	1562	285	323	5			

- Molecule 3 is a protein called HIV-1 gp120 core.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	G	336	Total	C	H	N	O	S	0	0	0
			4787	1661	2145	464	496	21			

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	H	1	Total	C	H	N	O	S	
			32	8	17	2	4	1	
4	G	1	Total	C	H	N	O	S	
			32	8	17	2	4	1	

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	H	1	Total	C	H	O		
			10	2	6	2	0	0
5	H	1	Total	C	H	O		
			10	2	6	2	0	0
5	L	1	Total	C	H	O		
			10	2	6	2	0	0
5	L	1	Total	C	H	O		
			10	2	6	2	0	0
5	L	1	Total	C	H	O		
			10	2	6	2	0	0
5	G	1	Total	C	H	O		
			10	2	6	2	0	0
5	G	1	Total	C	H	O		
			10	2	6	2	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	L	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	137	Total	O	0	0
			137	137		
7	L	133	Total	O	0	0
			133	133		
7	G	45	Total	O	0	0
			45	45		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Heavy chain of N6



- Molecule 2: Light chain of N6



- Molecule 3: HIV-1 gp120 core



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.67Å 65.55Å 241.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.52 – 2.15 44.52 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.6 (44.52-2.15) 92.0 (44.52-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.190 , 0.228 0.195 , 0.232	Depositor DCC
R_{free} test set	2023 reflections (3.83%)	wwPDB-VP
Wilson B-factor (Å ²)	42.2	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11993	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.15	0/1749	0.32	0/2383
2	L	0.16	0/1669	0.33	0/2264
3	G	0.14	1/2697 (0.0%)	0.25	0/3652
All	All	0.15	1/6115 (0.0%)	0.29	0/8299

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	252	LYS	C-N	5.02	1.45	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1703	1644	1670	4	0
2	L	1631	1562	1571	2	0
3	G	2642	2145	2592	15	0
4	G	15	17	17	0	0
4	H	15	17	17	0	0
5	G	8	12	12	1	0
5	H	8	12	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	12	18	18	0	0
6	G	98	91	91	0	0
6	L	14	14	13	0	0
7	G	45	0	0	0	0
7	H	137	0	0	3	1
7	L	133	0	0	1	1
All	All	6461	5532	6013	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:17:ASP:OD1	7:L:401:HOH:O	1.98	0.80
3:G:293:GLU:OE2	3:G:337:LYS:NZ	2.17	0.78
1:H:83:LYS:NZ	7:H:402:HOH:O	2.31	0.61
3:G:358:THR:OG1	3:G:465:THR:HG22	2.02	0.59
3:G:389:ASP:HB3	3:G:395:TYR:CE2	2.39	0.58
3:G:346:LYS:HG3	3:G:359:ILE:HB	1.86	0.56
3:G:269:GLU:OE1	3:G:348:LYS:NZ	2.41	0.53
3:G:462:LYS:HG2	3:G:464:ASP:H	1.74	0.52
3:G:425:ARG:HB3	5:G:509:EDO:H21	1.92	0.52
1:H:66:ARG:NH2	1:H:86:ASP:OD2	2.34	0.51
3:G:462:LYS:HE3	3:G:464:ASP:OD1	2.10	0.51
3:G:230:ASN:HB3	3:G:233:PHE:HB2	1.94	0.50
3:G:120:VAL:HG22	3:G:202:THR:HG22	1.94	0.50
3:G:395:TYR:O	3:G:396:MET:HB2	2.14	0.48
3:G:373:THR:HB	3:G:385:CYS:O	2.14	0.47
1:H:34:LEU:HD13	7:H:421:HOH:O	2.17	0.45
3:G:334:SER:HB3	3:G:337:LYS:HB2	1.97	0.45
1:H:214:LYS:NZ	7:H:407:HOH:O	2.43	0.44
3:G:460:LYS:HG2	3:G:463:ASN:CG	2.43	0.44
3:G:264:SER:O	3:G:287:HIS:NE2	2.43	0.43
2:L:76:SER:O	2:L:77:ASP:C	2.64	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:H:436:HOH:O	7:L:483:HOH:O[3_444]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	223/225 (99%)	220 (99%)	3 (1%)	0	100	100
2	L	208/210 (99%)	198 (95%)	9 (4%)	1 (0%)	24	20
3	G	328/350 (94%)	316 (96%)	11 (3%)	1 (0%)	36	34
All	All	759/785 (97%)	734 (97%)	23 (3%)	2 (0%)	36	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	354	PRO
2	L	68	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	185/185 (100%)	185 (100%)	0	100	100
2	L	187/187 (100%)	186 (100%)	1 (0%)	81	87
3	G	296/305 (97%)	294 (99%)	2 (1%)	76	82
All	All	668/677 (99%)	665 (100%)	3 (0%)	84	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	78	LEU
3	G	293	GLU
3	G	342	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	27	GLN
2	L	68	HIS
3	G	82	GLN
3	G	413	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EPE	H	301	-	15,15,15	0.84	1 (6%)	19,20,20	1.88	6 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	G	506	3	14,14,15	0.22	0	17,19,21	0.57	0
5	EDO	G	510	-	3,3,3	0.41	0	2,2,2	0.40	0
5	EDO	H	302	-	3,3,3	0.41	0	2,2,2	0.37	0
5	EDO	L	302	-	3,3,3	0.39	0	2,2,2	0.31	0
6	NAG	G	501	3	14,14,15	0.19	0	17,19,21	0.43	0
6	NAG	G	504	3	14,14,15	0.26	0	17,19,21	0.62	1 (5%)
6	NAG	G	502	3	14,14,15	0.23	0	17,19,21	0.42	0
6	NAG	G	503	3	14,14,15	0.30	0	17,19,21	0.50	0
5	EDO	L	304	-	3,3,3	0.43	0	2,2,2	0.32	0
6	NAG	G	505	3	14,14,15	0.23	0	17,19,21	0.43	0
6	NAG	G	507	3	14,14,15	0.19	0	17,19,21	0.45	0
5	EDO	G	509	-	3,3,3	0.43	0	2,2,2	0.32	0
5	EDO	H	303	-	3,3,3	0.41	0	2,2,2	0.35	0
4	EPE	G	508	-	15,15,15	0.80	1 (6%)	19,20,20	1.54	3 (15%)
5	EDO	L	303	-	3,3,3	0.41	0	2,2,2	0.32	0
6	NAG	L	301	2	14,14,15	0.25	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	H	301	-	-	5/9/19/19	0/1/1/1
6	NAG	G	506	3	-	4/6/23/26	0/1/1/1
5	EDO	G	510	-	-	0/1/1/1	-
5	EDO	H	302	-	-	0/1/1/1	-
5	EDO	L	302	-	-	1/1/1/1	-
6	NAG	G	501	3	-	2/6/23/26	0/1/1/1
6	NAG	G	504	3	-	0/6/23/26	0/1/1/1
6	NAG	G	502	3	-	3/6/23/26	0/1/1/1
6	NAG	G	503	3	-	2/6/23/26	0/1/1/1
5	EDO	L	304	-	-	0/1/1/1	-
6	NAG	G	505	3	-	2/6/23/26	0/1/1/1
6	NAG	G	507	3	-	0/6/23/26	0/1/1/1
5	EDO	G	509	-	-	0/1/1/1	-
5	EDO	H	303	-	-	0/1/1/1	-
4	EPE	G	508	-	-	7/9/19/19	0/1/1/1
5	EDO	L	303	-	-	0/1/1/1	-
6	NAG	L	301	2	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	508	EPE	C10-S	2.79	1.81	1.77
4	H	301	EPE	C10-S	2.74	1.81	1.77

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	301	EPE	C7-N4-C3	3.92	121.68	111.24
4	H	301	EPE	C7-N4-C5	3.79	121.35	111.24
4	G	508	EPE	C5-N4-C3	3.66	116.72	108.84
4	H	301	EPE	C5-N4-C3	3.62	116.64	108.84
4	G	508	EPE	C7-N4-C3	3.37	120.23	111.24
4	H	301	EPE	O1S-S-C10	2.69	110.79	106.73
4	H	301	EPE	O3S-S-C10	2.15	110.21	106.00
4	H	301	EPE	C6-N1-C2	2.09	113.34	108.84
4	G	508	EPE	C7-N4-C5	2.04	116.67	111.24
6	G	504	NAG	C1-O5-C5	2.04	114.91	112.19

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	502	NAG	C1-C2-N2-C7
4	G	508	EPE	C8-C7-N4-C3
6	G	501	NAG	O5-C5-C6-O6
6	G	506	NAG	O5-C5-C6-O6
6	G	501	NAG	C4-C5-C6-O6
6	G	506	NAG	C4-C5-C6-O6
6	L	301	NAG	C8-C7-N2-C2
6	L	301	NAG	O7-C7-N2-C2
6	G	506	NAG	C8-C7-N2-C2
6	G	506	NAG	O7-C7-N2-C2
6	G	503	NAG	O5-C5-C6-O6
6	G	503	NAG	C4-C5-C6-O6
6	G	505	NAG	C4-C5-C6-O6
6	G	505	NAG	O5-C5-C6-O6
4	G	508	EPE	S-C10-C9-N1
4	H	301	EPE	C10-C9-N1-C6
4	G	508	EPE	C9-C10-S-O3S
4	H	301	EPE	N4-C7-C8-O8
4	H	301	EPE	C8-C7-N4-C5
4	H	301	EPE	C9-C10-S-O2S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	G	508	EPE	C9-C10-S-O1S
4	G	508	EPE	C9-C10-S-O2S
4	H	301	EPE	C10-C9-N1-C2
4	G	508	EPE	C10-C9-N1-C2
6	G	502	NAG	O5-C5-C6-O6
6	G	502	NAG	C3-C2-N2-C7
4	G	508	EPE	C10-C9-N1-C6
5	L	302	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	509	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	225/225 (100%)	0.11	6 (2%) 56 60	36, 50, 76, 123	0
2	L	210/210 (100%)	-0.05	2 (0%) 79 82	35, 46, 70, 144	0
3	G	336/350 (96%)	0.82	32 (9%) 14 15	48, 75, 111, 167	0
All	All	771/785 (98%)	0.38	40 (5%) 33 36	35, 59, 104, 167	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	396	MET	5.4
1	H	130	SER	5.0
3	G	73	ALA	4.3
3	G	430	VAL	3.9
3	G	124	GLY	3.9
3	G	84	ILE	3.5
3	G	85	VAL	3.4
3	G	460	LYS	3.3
3	G	232	THR	3.3
3	G	410	GLY	3.2
3	G	45	TRP	3.1
1	H	74	TYR	2.9
3	G	459	GLY	2.9
1	H	216	CYS	2.8
3	G	208	VAL	2.8
1	H	131	THR	2.7
3	G	72	HIS	2.7
3	G	246	GLN	2.7
3	G	289	ASN	2.7
3	G	198	GLY	2.6
3	G	386	TYR	2.6
3	G	119	CYS	2.5
3	G	414	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	G	296	CYS	2.4
3	G	354	PRO	2.4
3	G	261	LEU	2.3
3	G	244	THR	2.3
3	G	201	VAL	2.3
1	H	129	LYS	2.3
2	L	1	TYR	2.3
3	G	82	GLN	2.2
3	G	301	ASN	2.2
3	G	353	PHE	2.1
3	G	336	LYS	2.1
3	G	86	LEU	2.1
1	H	128	SER	2.1
3	G	423	ILE	2.1
3	G	53	PHE	2.1
3	G	233	PHE	2.1
2	L	194	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	G	507	14/15	0.48	0.17	137,175,217,217	0
6	NAG	G	506	14/15	0.64	0.13	96,112,135,136	0
6	NAG	G	502	14/15	0.69	0.16	113,138,166,166	0
6	NAG	G	501	14/15	0.76	0.12	87,109,131,132	0
6	NAG	G	505	14/15	0.79	0.26	112,137,165,168	0
6	NAG	L	301	14/15	0.84	0.22	68,93,118,118	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EPE	G	508	15/15	0.84	0.18	76,91,97,99	0
5	EDO	G	509	4/4	0.84	0.17	67,81,86,86	0
6	NAG	G	504	14/15	0.87	0.11	75,81,97,98	0
5	EDO	L	304	4/4	0.88	0.12	63,76,79,79	0
4	EPE	H	301	15/15	0.89	0.19	79,109,144,147	0
6	NAG	G	503	14/15	0.89	0.10	51,73,85,92	0
5	EDO	L	303	4/4	0.89	0.16	55,67,70,73	0
5	EDO	G	510	4/4	0.90	0.17	86,103,104,104	0
5	EDO	L	302	4/4	0.92	0.16	68,82,92,95	0
5	EDO	H	302	4/4	0.94	0.10	45,54,57,59	0
5	EDO	H	303	4/4	0.96	0.09	45,54,59,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.