



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:31 AM UTC

PDB ID : 3TG3 / pdb_00003tg3
Title : Crystal structure of the MAPK binding domain of MKP7
Authors : Zhang, Y.Y.; Liu, X.; Wu, J.W.; Wang, Z.X.
Deposited on : 2011-08-17
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	
1	B	142	
1	C	142	
1	D	142	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity protein phosphatase 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	128	Total	C	N	O	S	0	0	0
			994	628	167	193	6			
1	B	128	Total	C	N	O	S	0	0	0
			984	621	164	193	6			
1	C	128	Total	C	N	O	S	0	0	0
			991	626	166	193	6			
1	D	127	Total	C	N	O	S	0	0	0
			979	618	163	192	6			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	139	LEU	-	expression tag	UNP Q9BY84
A	140	GLU	-	expression tag	UNP Q9BY84
A	141	HIS	-	expression tag	UNP Q9BY84
A	142	HIS	-	expression tag	UNP Q9BY84
A	143	HIS	-	expression tag	UNP Q9BY84
A	144	HIS	-	expression tag	UNP Q9BY84
A	145	HIS	-	expression tag	UNP Q9BY84
A	146	HIS	-	expression tag	UNP Q9BY84
B	139	LEU	-	expression tag	UNP Q9BY84
B	140	GLU	-	expression tag	UNP Q9BY84
B	141	HIS	-	expression tag	UNP Q9BY84
B	142	HIS	-	expression tag	UNP Q9BY84
B	143	HIS	-	expression tag	UNP Q9BY84
B	144	HIS	-	expression tag	UNP Q9BY84
B	145	HIS	-	expression tag	UNP Q9BY84
B	146	HIS	-	expression tag	UNP Q9BY84
C	139	LEU	-	expression tag	UNP Q9BY84
C	140	GLU	-	expression tag	UNP Q9BY84
C	141	HIS	-	expression tag	UNP Q9BY84
C	142	HIS	-	expression tag	UNP Q9BY84
C	143	HIS	-	expression tag	UNP Q9BY84

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	144	HIS	-	expression tag	UNP Q9BY84
C	145	HIS	-	expression tag	UNP Q9BY84
C	146	HIS	-	expression tag	UNP Q9BY84
D	139	LEU	-	expression tag	UNP Q9BY84
D	140	GLU	-	expression tag	UNP Q9BY84
D	141	HIS	-	expression tag	UNP Q9BY84
D	142	HIS	-	expression tag	UNP Q9BY84
D	143	HIS	-	expression tag	UNP Q9BY84
D	144	HIS	-	expression tag	UNP Q9BY84
D	145	HIS	-	expression tag	UNP Q9BY84
D	146	HIS	-	expression tag	UNP Q9BY84

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	23	Total	O	0	0
			23	23		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	21	Total	O	0	0
			21	21		
3	D	21	Total	O	0	0
			21	21		

- Molecule 1: Dual specificity protein phosphatase 16





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.50Å 47.48Å 64.51Å 91.18° 97.29° 96.81°	Depositor
Resolution (Å)	28.04 – 2.67 28.04 – 2.67	Depositor EDS
% Data completeness (in resolution range)	86.5 (28.04-2.67) 94.7 (28.04-2.67)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 2.68Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.239 , 0.288 0.247 , 0.299	Depositor DCC
R_{free} test set	623 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.544	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 76.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4042	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8456e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/1005 (0.1%)	0.95	0/1354
1	B	0.80	0/994	1.04	3/1340 (0.2%)
1	C	0.68	1/1002 (0.1%)	0.92	1/1352 (0.1%)
1	D	0.52	0/989	0.90	1/1334 (0.1%)
All	All	0.68	2/3990 (0.1%)	0.95	5/5380 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	51	SER	CA-C	-5.88	1.45	1.52
1	C	52	LYS	CA-C	-5.02	1.46	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	HIS	N-CA-C	-8.42	102.10	111.28
1	B	132	PHE	CA-C-N	6.75	127.02	119.32
1	B	132	PHE	C-N-CA	6.75	127.02	119.32
1	D	35	VAL	CB-CA-C	-5.11	105.33	111.88
1	C	35	VAL	CB-CA-C	-5.08	105.47	111.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	994	0	1008	60	0
1	B	984	0	993	48	0
1	C	991	0	1002	60	0
1	D	979	0	988	53	0
2	C	4	0	6	1	0
2	D	4	0	6	2	0
3	A	21	0	0	3	0
3	B	23	0	0	3	0
3	C	21	0	0	5	0
3	D	21	0	0	5	0
All	All	4042	0	4003	207	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ALA:O	1:A:74:LYS:HG3	1.34	1.27
1:C:77:VAL:CG1	1:C:78:ASP:H	1.63	1.10
1:C:77:VAL:HG13	1:C:78:ASP:N	1.59	1.05
1:A:67:GLU:HG2	1:A:71:HIS:NE2	1.74	1.03
1:B:49:ASN:HD22	1:B:49:ASN:H	1.05	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	124/142 (87%)	112 (90%)	10 (8%)	2 (2%)	7 18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	124/142 (87%)	114 (92%)	8 (6%)	2 (2%)	7	18
1	C	124/142 (87%)	114 (92%)	8 (6%)	2 (2%)	7	18
1	D	123/142 (87%)	112 (91%)	9 (7%)	2 (2%)	7	18
All	All	495/568 (87%)	452 (91%)	35 (7%)	8 (2%)	7	18

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	ASP
1	B	80	ASP
1	C	80	ASP
1	D	80	ASP
1	D	23	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/129 (89%)	102 (89%)	13 (11%)	5	13
1	B	113/129 (88%)	105 (93%)	8 (7%)	13	30
1	C	115/129 (89%)	103 (90%)	12 (10%)	7	15
1	D	113/129 (88%)	104 (92%)	9 (8%)	11	25
All	All	456/516 (88%)	414 (91%)	42 (9%)	8	19

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	70	GLN
1	D	35	VAL
1	C	77	VAL
1	C	130	ARG
1	D	49	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	9	GLN
1	D	49	ASN
1	D	59	GLN
1	B	59	GLN
1	B	70	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	D	3318	-	3,3,3	0.48	0	2,2,2	0.22	0
2	EDO	C	3318	-	3,3,3	0.47	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	3318	-	-	1/1/1/1	-
2	EDO	C	3318	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3318	EDO	O1-C1-C2-O2
2	D	3318	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3318	EDO	2	0
2	C	3318	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	128/142 (90%)	0.49	10 (7%)	19 16	7, 36, 86, 127	0
1	B	128/142 (90%)	0.64	13 (10%)	12 10	12, 35, 86, 129	0
1	C	128/142 (90%)	0.68	17 (13%)	7 6	12, 38, 89, 117	0
1	D	127/142 (89%)	0.63	17 (13%)	7 6	9, 37, 98, 152	0
All	All	511/568 (89%)	0.61	57 (11%)	10 8	7, 37, 91, 152	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	66	THR	4.4
1	C	63	VAL	4.1
1	B	82	SER	4.1
1	D	63	VAL	4.0
1	D	62	LYS	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	C	3318	4/4	0.77	0.16	24,25,33,35	0
2	EDO	D	3318	4/4	0.89	0.11	17,22,26,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.