



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:53 PM UTC

PDB ID : 7TGQ / pdb_00007tgq
Title : Zinc finger antiviral protein (ZAP) central domain bound to ADP-ribose
Authors : Pornillos, O.P.
Deposited on : 2022-01-09
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	704	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2959 atoms, of which 1393 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Zinc finger CCCH-type antiviral protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	182	2812	924	1366	250	260	12	0	0	0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ADENOSINE-5-DIPHOSPHORIBOSE (CCD ID: APR) (formula: C₁₅H₂₃N₅O₁₄P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
3	A	1	57	15	21	5	14	2	0	0

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			7	2	3	2		
4	A	1	Total	C	H	O	0	0
			7	2	3	2		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	75	Total	O	0	0
			75	75		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Zinc finger CCCH-type antiviral protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.96Å 89.96Å 52.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.71 – 2.00 43.71 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.9 (43.71-2.00) 94.7 (43.71-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.238 , 0.273 (Not available) , 0.285	Depositor DCC
R_{free} test set	843 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2959	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT, APR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/1489	0.58	0/2025

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1446	1366	1366	16	0
2	A	1	0	0	0	0
3	A	36	21	21	0	0
4	A	8	6	6	5	0
5	A	75	0	0	0	0
All	All	1566	1393	1393	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:572:TYR:HB3	1:A:581:CYS:SG	2.18	0.83
1:A:637:TYR:CE1	1:A:641:LEU:HD11	2.37	0.60
1:A:637:TYR:HE1	1:A:641:LEU:HD11	1.74	0.52
1:A:684:THR:HB	4:A:703:ACT:H1	1.92	0.51
1:A:692:GLN:HG2	4:A:704:ACT:H3	1.93	0.50
1:A:541:LEU:HD23	1:A:585:PRO:HG2	1.92	0.50
1:A:581:CYS:O	1:A:584:PHE:N	2.46	0.48
1:A:670:ASN:C	1:A:670:ASN:OD1	2.57	0.47
1:A:695:LYS:NZ	4:A:704:ACT:H1	2.30	0.47
1:A:691:VAL:CG1	4:A:704:ACT:H2	2.45	0.46
1:A:691:VAL:HG12	4:A:704:ACT:H2	1.98	0.46
1:A:557:LYS:HD3	1:A:685:PHE:CE2	2.53	0.44
1:A:541:LEU:HB3	1:A:585:PRO:HG2	2.00	0.42
1:A:662:SER:O	1:A:666:MET:N	2.52	0.42
1:A:573:THR:O	1:A:581:CYS:SG	2.78	0.41
1:A:550:GLU:OE1	1:A:550:GLU:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/202 (88%)	172 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/187 (84%)	157 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	A	704	-	3,3,3	0.92	0	3,3,3	1.53	1 (33%)
4	ACT	A	703	-	3,3,3	1.32	0	3,3,3	1.62	1 (33%)
3	APR	A	702	-	39,39,39	1.80	10 (25%)	56,60,60	1.80	14 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	APR	A	702	-	-	12/22/54/54	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	APR	PA-O3A	4.08	1.63	1.59
3	A	702	APR	C8-N7	3.62	1.38	1.31
3	A	702	APR	C2-N1	3.47	1.40	1.33
3	A	702	APR	C1D-C2D	3.33	1.56	1.52
3	A	702	APR	PB-O1B	3.14	1.61	1.50
3	A	702	APR	PB-O3A	3.03	1.62	1.59
3	A	702	APR	C2-N3	2.98	1.39	1.33
3	A	702	APR	PA-O1A	2.77	1.60	1.50
3	A	702	APR	C4-N3	2.66	1.39	1.34
3	A	702	APR	C6-N6	2.30	1.40	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	APR	N3-C2-N1	-5.48	120.29	128.58
3	A	702	APR	C5-C4-N3	-5.08	119.72	126.72
3	A	702	APR	N9-C8-N7	-3.79	108.56	113.94
3	A	702	APR	N3-C4-N9	3.51	133.13	127.17
3	A	702	APR	C2-N3-C4	3.26	119.80	111.83
3	A	702	APR	C5-N7-C8	2.88	107.98	103.45
3	A	702	APR	O4D-C1D-C2D	2.81	108.57	104.67
3	A	702	APR	O3A-PA-O1A	-2.74	102.45	110.70
3	A	702	APR	C1D-C2D-C3D	2.54	105.42	102.29
3	A	702	APR	C4-N9-C8	2.32	108.17	105.74
4	A	704	ACT	OXT-C-O	2.25	130.37	122.03
4	A	703	ACT	OXT-C-O	2.18	130.12	122.03
3	A	702	APR	C6-C5-C4	2.16	120.13	117.18
3	A	702	APR	C4-C5-N7	-2.13	108.14	110.58
3	A	702	APR	C5-C6-N6	-2.09	118.10	123.29
3	A	702	APR	O4'-C1'-N9	2.02	111.96	108.09

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	702	APR	C5'-O5'-PA-O1A
3	A	702	APR	C5'-O5'-PA-O2A
3	A	702	APR	C5'-O5'-PA-O3A
3	A	702	APR	PB-O3A-PA-O5'
3	A	702	APR	C5D-O5D-PB-O3A
3	A	702	APR	C5D-O5D-PB-O1B
3	A	702	APR	C5D-O5D-PB-O2B
3	A	702	APR	C3'-C4'-C5'-O5'
3	A	702	APR	O4'-C4'-C5'-O5'
3	A	702	APR	C2'-C1'-N9-C8
3	A	702	APR	C4D-C5D-O5D-PB
3	A	702	APR	O4'-C1'-N9-C8

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	704	ACT	4	0
4	A	703	ACT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/202 (90%)	1.14	53 (29%) 1 1	15, 35, 66, 95	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	509	SER	6.6
1	A	614	GLU	6.5
1	A	572	TYR	5.9
1	A	571	SER	5.6
1	A	541	LEU	5.0
1	A	519	LYS	4.5
1	A	584	PHE	4.3
1	A	582	ASP	4.3
1	A	570	GLY	4.3
1	A	543	GLY	4.1
1	A	583	SER	4.0
1	A	542	ILE	3.6
1	A	524	ASN	3.5
1	A	580	SER	3.4
1	A	573	THR	3.3
1	A	616	GLY	3.2
1	A	617	THR	3.2
1	A	540	MET	3.1
1	A	550	GLU	3.1
1	A	569	VAL	3.1
1	A	544	LYS	3.0
1	A	585	PRO	3.0
1	A	527	CYS	2.9
1	A	575	ASN	2.9
1	A	523	LEU	2.8
1	A	615	SER	2.8
1	A	581	CYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	647	ARG	2.7
1	A	637	TYR	2.7
1	A	673	SER	2.7
1	A	578	VAL	2.6
1	A	600	ASN	2.6
1	A	672	ALA	2.6
1	A	568	SER	2.6
1	A	630	ASN	2.5
1	A	641	LEU	2.5
1	A	612	LYS	2.5
1	A	664	GLN	2.4
1	A	547	THR	2.4
1	A	566	LEU	2.4
1	A	545	THR	2.3
1	A	623	GLU	2.3
1	A	674	LYS	2.3
1	A	574	ILE	2.2
1	A	549	PHE	2.1
1	A	577	ARG	2.1
1	A	644	SER	2.1
1	A	510	GLU	2.1
1	A	522	PRO	2.1
1	A	565	HIS	2.1
1	A	525	GLY	2.1
1	A	528	SER	2.1
1	A	586	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	APR	A	702	36/36	0.85	0.15	23,54,96,103	0
4	ACT	A	704	4/4	0.92	0.10	18,22,29,31	0
4	ACT	A	703	4/4	0.93	0.10	21,24,29,38	0
2	ZN	A	701	1/1	0.98	0.04	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.