



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:07 AM EDT

PDB ID : 1THY / pdb_00001thy
Title : REFINED STRUCTURES OF SUBSTRATE-BOUND AND PHOSPHATE-BOUND THYMIDYLATE SYNTHASE FROM LACTOBACILLUS CASEI
Authors : Finer-Moore, J.; Stroud, R.
Deposited on : 1993-04-02
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UMP	A	317	X	-	-	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2590	1677	438	467	8	0	0	0

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (CCD ID: UMP) (formula: C₉H₁₃N₂O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	20	9	2	8	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	30	Total 30 O 30	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: THYMIDYLATE SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	78.80Å 78.80Å 230.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.90 68.24 – 2.91	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.90) 88.0 (68.24-2.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.53 (at 2.90Å)	Xtrriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.155 , (Not available) 0.154 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtrriage
Anisotropy	0.145	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 74.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2640	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.22	23/2674 (0.9%)	1.95	81/3634 (2.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	106	HIS	CD2-NE2	-7.30	1.29	1.37
1	A	199	HIS	CD2-NE2	-7.05	1.30	1.37
1	A	253	HIS	CD2-NE2	-6.78	1.30	1.37
1	A	66	HIS	CD2-NE2	-6.62	1.30	1.37
1	A	77	HIS	CD2-NE2	-6.62	1.30	1.37
1	A	97	HIS	CD2-NE2	-6.62	1.30	1.37
1	A	80	HIS	CD2-NE2	-6.57	1.30	1.37
1	A	289	HIS	CD2-NE2	-6.42	1.30	1.37
1	A	242	HIS	CD2-NE2	-6.20	1.31	1.37
1	A	238	HIS	CD2-NE2	-6.20	1.31	1.37
1	A	264	HIS	CD2-NE2	-6.16	1.31	1.37
1	A	18	HIS	CD2-NE2	-6.10	1.31	1.37
1	A	259	HIS	CD2-NE2	-6.07	1.31	1.37
1	A	34	HIS	CD2-NE2	-6.03	1.31	1.37
1	A	154	HIS	CD2-NE2	-5.88	1.31	1.37
1	A	199	HIS	CG-ND1	-5.86	1.31	1.38
1	A	174	HIS	CD2-NE2	-5.76	1.31	1.37
1	A	119	HIS	CD2-NE2	-5.74	1.31	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	25	HIS	CD2-NE2	-5.72	1.31	1.37
1	A	238	HIS	CG-ND1	-5.45	1.32	1.38
1	A	229	ASN	CA-CB	-5.26	1.45	1.53
1	A	286	PRO	CA-CB	-5.24	1.46	1.53
1	A	77	HIS	CG-ND1	-5.00	1.32	1.38

All (81) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	ASP	CA-CB-CG	11.78	124.38	112.60
1	A	71	ILE	CA-CB-CG1	-9.70	93.92	110.40
1	A	221	ASP	CA-CB-CG	9.56	122.16	112.60
1	A	22	ASP	CA-CB-CG	8.85	121.45	112.60
1	A	149	GLN	OE1-CD-NE2	-8.79	113.81	122.60
1	A	18	HIS	CA-CB-CG	-8.74	105.06	113.80
1	A	196	PRO	O-C-N	-8.60	117.11	121.15
1	A	94	ASP	N-CA-C	8.34	121.30	111.71
1	A	106	HIS	CB-CG-CD2	-8.23	120.50	131.20
1	A	71	ILE	CA-CB-CG2	8.00	124.09	110.50
1	A	289	HIS	CA-CB-CG	-7.95	105.85	113.80
1	A	97	HIS	N-CA-C	-7.82	94.15	107.99
1	A	77	HIS	CA-CB-CG	-7.36	106.44	113.80
1	A	160	THR	CA-CB-OG1	-7.29	98.66	109.60
1	A	154	HIS	CB-CG-CD2	-7.27	121.74	131.20
1	A	66	HIS	CB-CG-CD2	-7.17	121.88	131.20
1	A	141	ASP	CA-CB-CG	7.17	119.77	112.60
1	A	149	GLN	CG-CD-NE2	7.13	127.10	116.40
1	A	80	HIS	CB-CG-CD2	-7.10	121.97	131.20
1	A	303	ASN	CB-CA-C	-7.04	102.81	111.82
1	A	160	THR	N-CA-CB	-7.03	99.00	110.81
1	A	93	SER	N-CA-C	6.96	119.06	109.18
1	A	104	PHE	N-CA-C	6.93	119.71	111.33
1	A	2	LEU	N-CA-C	6.73	119.44	111.71
1	A	270	GLU	N-CA-C	-6.70	103.66	110.97
1	A	120	GLU	CB-CG-CD	6.54	123.73	112.60
1	A	295	ASP	N-CA-CB	-6.51	99.42	111.53
1	A	289	HIS	CB-CG-CD2	-6.39	122.89	131.20
1	A	132	ASP	CA-CB-CG	6.28	118.88	112.60
1	A	178	ARG	N-CA-C	-6.27	104.95	112.59
1	A	83	ASP	N-CA-C	6.26	118.66	111.02
1	A	95	GLU	CB-CG-CD	6.15	123.05	112.60
1	A	154	HIS	CB-CG-ND1	6.10	131.85	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	294	PHE	N-CA-C	-6.09	102.33	110.55
1	A	106	HIS	CB-CG-ND1	6.08	131.82	122.70
1	A	34	HIS	CB-CG-CD2	-5.96	123.46	131.20
1	A	82	TRP	CE2-CD2-CE3	5.93	124.73	118.80
1	A	160	THR	CB-CA-C	5.92	120.02	109.65
1	A	310	ILE	N-CA-C	-5.89	99.86	108.11
1	A	4	GLN	CA-C-O	-5.84	113.26	118.63
1	A	18	HIS	CB-CG-CD2	-5.83	123.62	131.20
1	A	83	ASP	CA-CB-CG	5.79	118.39	112.60
1	A	242	HIS	CA-CB-CG	5.76	119.56	113.80
1	A	238	HIS	CB-CG-CD2	-5.76	123.72	131.20
1	A	106	HIS	CA-CB-CG	-5.74	108.06	113.80
1	A	151	ARG	N-CA-C	5.72	120.25	112.88
1	A	174	HIS	CB-CG-CD2	-5.66	123.85	131.20
1	A	274	ARG	CB-CG-CD	5.66	124.31	111.30
1	A	285	ASN	CA-C-N	5.60	125.44	119.28
1	A	285	ASN	C-N-CA	5.60	125.44	119.28
1	A	181	ILE	CA-C-O	-5.44	114.50	121.40
1	A	178	ARG	CB-CG-CD	-5.43	98.81	111.30
1	A	77	HIS	CB-CG-CD2	-5.36	124.23	131.20
1	A	37	ARG	N-CA-CB	-5.36	101.80	110.81
1	A	313	PRO	N-CA-C	5.35	119.49	111.14
1	A	263	ASN	OD1-CG-ND2	-5.33	117.27	122.60
1	A	34	HIS	CB-CG-ND1	5.33	130.69	122.70
1	A	269	LYS	CB-CG-CD	-5.32	99.07	111.30
1	A	195	LEU	CA-C-N	5.30	123.47	119.66
1	A	195	LEU	C-N-CA	5.30	123.47	119.66
1	A	270	GLU	CA-CB-CG	5.29	124.68	114.10
1	A	150	TRP	CG-CD2-CE3	5.24	139.14	133.90
1	A	314	VAL	N-CA-C	-5.22	101.95	108.84
1	A	71	ILE	N-CA-CB	-5.21	100.45	110.78
1	A	248	VAL	N-CA-CB	-5.21	101.37	110.49
1	A	153	TRP	CE2-CD2-CG	-5.20	100.97	107.20
1	A	153	TRP	CG-CD2-CE3	5.18	139.08	133.90
1	A	97	HIS	CB-CG-CD2	-5.15	124.51	131.20
1	A	23	ARG	NE-CZ-NH2	-5.14	114.58	119.20
1	A	185	TRP	CE2-CD2-CG	-5.13	101.04	107.20
1	A	199	HIS	CB-CG-CD2	-5.13	124.54	131.20
1	A	150	TRP	CG-CD1-NE1	-5.10	103.58	110.20
1	A	94	ASP	CA-C-O	-5.08	114.36	120.10
1	A	185	TRP	CG-CD2-CE3	5.07	138.97	133.90
1	A	223	PHE	N-CA-C	5.06	116.65	111.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	LYS	N-CA-CB	-5.04	102.70	110.01
1	A	103	ASP	CA-C-N	5.03	127.27	120.38
1	A	103	ASP	C-N-CA	5.03	127.27	120.38
1	A	37	ARG	O-C-N	-5.01	117.24	123.36
1	A	85	TRP	CE2-CD2-CG	-5.00	101.19	107.20
1	A	160	THR	CA-CB-CG2	5.00	119.00	110.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	23	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2590	0	2496	36	0
2	A	20	0	10	3	0
3	A	30	0	0	2	0
All	All	2640	0	2506	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:CYS:SG	2:A:317:UMP:H6	2.07	0.77
1:A:274:ARG:HG2	1:A:307:TYR:CE2	2.31	0.65
1:A:217:GLN:HE22	1:A:229:ASN:ND2	1.97	0.62
1:A:177:SER:HB3	1:A:180:LEU:HD13	1.85	0.59
1:A:5:PRO:HB3	1:A:36:MET:HB2	1.86	0.57
1:A:142:LEU:HB2	1:A:145:VAL:HG21	1.87	0.56
1:A:274:ARG:HG2	1:A:307:TYR:CD2	2.41	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:HG2	1:A:76:GLN:HE21	1.71	0.54
1:A:221:ASP:HB3	2:A:317:UMP:HN3	1.72	0.53
1:A:243:GLU:HB2	1:A:291:ILE:HG23	1.89	0.53
1:A:101:MET:HE3	1:A:114:PHE:CZ	2.45	0.52
1:A:66:HIS:HD2	3:A:345:HOH:O	1.93	0.50
1:A:112:PRO:HG2	1:A:114:PHE:HB2	1.92	0.50
1:A:198:CYS:SG	2:A:317:UMP:C6	2.99	0.50
1:A:207:ASN:HB3	3:A:346:HOH:O	2.10	0.50
1:A:283:GLN:HB3	1:A:300:LYS:HB2	1.94	0.49
1:A:23:ARG:N	1:A:23:ARG:HD3	2.28	0.49
1:A:175:PRO:HB2	1:A:206:VAL:HG11	1.96	0.48
1:A:90:TRP:HH2	1:A:95:GLU:OE1	1.98	0.46
1:A:142:LEU:HB2	1:A:145:VAL:CG2	2.45	0.46
1:A:90:TRP:CZ3	1:A:95:GLU:HB3	2.50	0.46
1:A:265:LEU:O	1:A:269:LYS:HD2	2.17	0.45
1:A:180:LEU:HB3	1:A:204:PHE:HB2	1.99	0.44
1:A:222:ILE:HD11	1:A:258:ALA:HB1	2.00	0.44
1:A:40:LEU:HB2	1:A:248:VAL:HG13	2.00	0.44
1:A:223:PHE:HD1	1:A:268:ILE:HD13	1.82	0.44
1:A:72:ARG:O	1:A:76:GLN:HG3	2.19	0.43
1:A:72:ARG:HH22	1:A:133:ASP:HA	1.84	0.42
1:A:175:PRO:O	1:A:206:VAL:HB	2.19	0.42
1:A:186:ASN:O	1:A:190:VAL:HG13	2.20	0.42
1:A:106:HIS:O	1:A:109:GLN:HB2	2.20	0.42
1:A:285:ASN:HA	1:A:286:PRO:HD3	1.86	0.42
1:A:11:LYS:O	1:A:14:LEU:HB2	2.20	0.41
1:A:185:TRP:HB2	1:A:200:THR:HG23	2.03	0.41
1:A:195:LEU:HD12	1:A:195:LEU:HA	1.81	0.41
1:A:101:MET:HE3	1:A:114:PHE:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/316 (99%)	292 (93%)	21 (7%)	1 (0%)	36 65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	112	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	277/278 (100%)	243 (88%)	34 (12%)	4 15

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	9	LEU
1	A	62	LEU
1	A	65	LEU
1	A	71	ILE
1	A	89	LYS
1	A	101	MET
1	A	103	ASP
1	A	110	LYS
1	A	113	GLU
1	A	119	HIS
1	A	128	ARG
1	A	130	LEU
1	A	133	ASP
1	A	144	LEU
1	A	145	VAL
1	A	160	THR
1	A	164	LEU
1	A	179	ARG
1	A	180	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	187	PRO
1	A	215	LEU
1	A	236	LEU
1	A	239	LEU
1	A	243	GLU
1	A	246	LEU
1	A	248	VAL
1	A	262	VAL
1	A	265	LEU
1	A	277	ARG
1	A	278	PRO
1	A	284	LEU
1	A	301	LEU
1	A	302	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	66	HIS
1	A	76	GLN
1	A	170	GLN
1	A	229	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UMP	A	317	-	21,21,21	1.38	4 (19%)	30,31,31	3.23	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	317	-	1/1/4/4	3/10/22/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	317	UMP	C6-N1	-3.18	1.30	1.38
2	A	317	UMP	P-OP3	-2.26	1.46	1.54
2	A	317	UMP	C5'-C4'	2.20	1.58	1.51
2	A	317	UMP	P-OP2	-2.07	1.47	1.54

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	317	UMP	O4'-C1'-N1	16.27	136.74	107.86
2	A	317	UMP	O5'-C5'-C4'	2.88	118.80	108.99
2	A	317	UMP	C1'-N1-C6	-2.85	115.92	121.53
2	A	317	UMP	C4'-O4'-C1'	-2.29	104.06	109.51
2	A	317	UMP	O5'-P-OP1	2.25	112.52	106.44
2	A	317	UMP	C2'-C3'-C4'	-2.23	98.28	102.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	317	UMP	C1'

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	317	UMP	O4'-C1'-N1-C6
2	A	317	UMP	O4'-C1'-N1-C2
2	A	317	UMP	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	317	UMP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/316 (100%)	-0.90	10 (3%) 50 41	4, 19, 60, 60	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	25	HIS	5.0
1	A	100	ASP	4.0
1	A	111	ASP	3.3
1	A	109	GLN	3.3
1	A	102	THR	3.3
1	A	113	GLU	2.7
1	A	106	HIS	2.6
1	A	112	PRO	2.5
1	A	97	HIS	2.4
1	A	103	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	UMP	A	317	20/20	0.97	0.08	6,18,27,31	15

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.