



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 12:55 AM UTC

PDB ID : 5TH2 / pdb\_00005th2  
Title : Cetuximab Fab in complex with L5Q mediotope variant  
Authors : Bzymek, K.P.; Williams, J.C.  
Deposited on : 2016-09-28  
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 91% 9%
1	C	213	 92% 8%
2	B	221	 90% 7% .
2	D	221	 93% 6%
3	E	12	 100%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	12	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '8%', a green segment in the middle labeled '67%', and a yellow segment on the right labeled '33%'.</p>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cetuximab Fab, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	213	Total	C	N	O	S	0	11	0
			1708	1057	291	355	5			
1	C	213	Total	C	N	O	S	0	8	0
			1676	1045	286	340	5			

- Molecule 2 is a protein called cetuximab Fab, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	216	Total	C	N	O	S	0	5	0
			1659	1056	274	323	6			
2	D	220	Total	C	N	O	S	0	6	0
			1688	1074	278	330	6			

- Molecule 3 is a protein called L5Q meditope.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	12	Total	C	N	O	S	0	0	0
			102	60	21	19	2			
3	F	12	Total	C	N	O	S	0	0	0
			102	60	21	19	2			

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	218	Total O 218 218	0	0
5	B	185	Total O 185 185	0	0
5	C	233	Total O 233 233	0	0
5	D	190	Total O 190 190	0	0
5	E	12	Total O 12 12	0	0
5	F	12	Total O 12 12	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cetuximab Fab, light chain

Chain A:  91% 9%





- Molecule 1: cetuximab Fab, light chain

Chain C:  92% 8%



- Molecule 2: cetuximab Fab, heavy chain

Chain B:  90% 7% .



- Molecule 2: cetuximab Fab, heavy chain

Chain D:  93% 6%



- Molecule 3: L5Q meditope

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: L5Q meditope

Chain F:  8% 67% 33%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.19Å 83.12Å 212.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.77 – 1.84 44.77 – 1.84	Depositor EDS
% Data completeness (in resolution range)	92.3 (44.77-1.84) 92.3 (44.77-1.84)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 1.84Å)	Xtrriage
Refinement program	PHENIX 1.10_2155	Depositor
R, $R_{free}$	0.161 , 0.184 0.162 , 0.185	Depositor DCC
$R_{free}$ test set	4595 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtrriage
Anisotropy	0.295	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 53.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7800	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1751	0.50	0/2377
1	C	0.27	0/1728	0.51	0/2345
2	B	0.26	0/1711	0.52	0/2337
2	D	0.26	0/1747	0.50	0/2385
3	E	0.28	0/102	0.52	0/132
3	F	0.31	0/102	0.63	0/132
All	All	0.27	0/7141	0.51	0/9708

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1708	0	1634	16	0
1	C	1676	0	1633	14	0
2	B	1659	0	1623	9	0
2	D	1688	0	1661	10	0
3	E	102	0	101	0	0
3	F	102	0	101	4	0
4	A	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	5	0	0	0	0
4	C	5	0	0	1	0
5	A	218	0	0	3	0
5	B	185	0	0	2	3
5	C	233	0	0	5	1
5	D	190	0	0	2	3
5	E	12	0	0	0	0
5	F	12	0	0	0	0
All	All	7800	0	6753	48	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1:ASP:OD2	5:C:401:HOH:O	1.99	0.79
1:A:45[B]:ARG:NH1	5:A:401:HOH:O	2.15	0.74
4:C:301:PO4:O2	5:C:402:HOH:O	2.06	0.72
1:C:161:GLU:HG3	5:C:406:HOH:O	1.93	0.67
1:C:165:GLU:OE1	5:C:403:HOH:O	2.13	0.66
1:C:123[B]:GLU:HG2	5:D:340:HOH:O	1.99	0.62
1:A:161:GLU:HG3	5:A:407:HOH:O	2.03	0.59
2:B:11:LEU:HD11	2:B:118:SER:HB3	1.85	0.58
2:D:154:GLU:OE1	3:F:6:SER:OG	2.17	0.57
2:B:67:LEU:HD11	2:B:80:PHE:CE2	2.40	0.56
1:A:145:LYS:HB3	1:A:197[B]:THR:HG22	1.86	0.56
1:C:10:ILE:HD11	3:F:12:CYS:HB2	1.91	0.52
2:D:129:PRO:HD3	2:D:215:LYS:HE2	1.90	0.52
1:A:145:LYS:HB3	1:A:197[B]:THR:CG2	2.40	0.52
2:D:11:LEU:HD11	2:D:118:SER:HB3	1.91	0.52
1:C:9:VAL:HG23	1:C:10:ILE:HG13	1.90	0.51
2:B:215:LYS:NZ	5:B:406:HOH:O	2.43	0.51
1:C:166:GLN:HG3	1:C:173:TYR:CZ	2.47	0.50
5:C:433:HOH:O	2:D:149:LYS:HE2	2.11	0.50
1:A:45[B]:ARG:NH1	5:A:406:HOH:O	2.44	0.49
1:C:33[A]:ILE:HG21	1:C:71:PHE:CD1	2.48	0.49
1:A:197[A]:THR:HG22	1:A:204:PRO:HB3	1.94	0.48
2:D:92[A]:ILE:HD12	3:F:3:PHE:HZ	1.79	0.47
1:C:80:SER:HA	1:C:106:LEU:HD22	1.96	0.47
2:B:7:SER:HB3	5:B:555:HOH:O	2.14	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:ILE:HD13	1:A:166:GLN:HB3	1.97	0.47
2:B:13[A]:GLN:OE1	2:B:119:ALA:HA	2.14	0.47
1:A:80:SER:HA	1:A:106:LEU:HD22	1.98	0.46
2:B:146[B]:CYS:HB3	2:B:202:CYS:SG	2.56	0.44
1:C:145:LYS:HB2	1:C:197:THR:HB	2.00	0.44
2:D:72:ASP:OD2	2:D:75:LYS:HD3	2.16	0.44
1:A:11:LEU:CD1	1:C:8:PRO:HB3	2.48	0.43
1:A:37:GLN:HB2	1:A:47:LEU:HD11	2.00	0.43
1:A:166:GLN:HG3	1:A:173:TYR:CZ	2.53	0.43
2:B:132:PRO:HB3	2:B:143:ALA:O	2.18	0.43
1:C:120:PRO:HD3	1:C:132:VAL:HG22	2.01	0.43
2:D:146[B]:CYS:SG	2:D:202:CYS:SG	3.16	0.43
1:A:2:ILE:HD12	1:A:27:GLN:HG2	2.01	0.43
2:D:123:LYS:NZ	5:D:312:HOH:O	2.52	0.42
1:A:145:LYS:HB3	1:A:197[A]:THR:OG1	2.19	0.42
1:A:123[A]:GLU:O	1:A:126:LYS:HG2	2.18	0.42
1:C:106:LEU:HD12	1:C:106:LEU:HA	1.74	0.42
1:A:19:VAL:HG22	1:A:75:ILE:HB	2.02	0.42
2:B:100:THR:OG1	2:B:103:ASP:HB2	2.21	0.41
1:C:116:PHE:HD2	2:D:136:SER:HA	1.86	0.41
2:B:146[B]:CYS:SG	2:B:202:CYS:SG	3.19	0.41
2:D:92[A]:ILE:HD12	3:F:3:PHE:CZ	2.56	0.41
1:A:123[B]:GLU:O	1:A:126:LYS:HG2	2.20	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:548:HOH:O	5:D:431:HOH:O[4_555]	1.78	0.42
5:B:469:HOH:O	5:D:483:HOH:O[3_544]	2.03	0.17
5:C:445:HOH:O	5:C:571:HOH:O[3_444]	2.06	0.14
5:B:548:HOH:O	5:D:414:HOH:O[4_555]	2.18	0.02

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/213 (104%)	217 (98%)	5 (2%)	0	100	100
1	C	219/213 (103%)	213 (97%)	6 (3%)	0	100	100
2	B	217/221 (98%)	216 (100%)	1 (0%)	0	100	100
2	D	224/221 (101%)	223 (100%)	1 (0%)	0	100	100
3	E	10/12 (83%)	10 (100%)	0	0	100	100
3	F	10/12 (83%)	10 (100%)	0	0	100	100
All	All	902/892 (101%)	889 (99%)	13 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/188 (105%)	196 (99%)	2 (1%)	68	58
1	C	194/188 (103%)	194 (100%)	0	100	100
2	B	190/191 (100%)	187 (98%)	3 (2%)	55	39
2	D	195/191 (102%)	194 (100%)	1 (0%)	81	75
3	E	12/12 (100%)	12 (100%)	0	100	100
3	F	12/12 (100%)	11 (92%)	1 (8%)	10	2
All	All	801/782 (102%)	794 (99%)	7 (1%)	70	60

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	VAL
1	A	33	ILE
2	B	100	THR
2	B	105	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	155	PRO
2	D	105	GLU
3	F	11	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	ASN
1	A	41	ASN
1	A	92	ASN
1	A	93	ASN
2	B	3	GLN
1	C	32	ASN
1	C	38	GLN
1	C	92	ASN
1	C	160	GLN
2	D	39	GLN
2	D	70	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PO4	B	301	-	4,4,4	0.91	0	6,6,6	0.53	0
4	PO4	A	301	-	4,4,4	1.04	0	6,6,6	0.64	0
4	PO4	C	301	-	4,4,4	1.06	0	6,6,6	0.40	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	301	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/213 (100%)	-0.33	1 (0%) 87 92	12, 27, 45, 71	11 (5%)
1	C	213/213 (100%)	-0.25	1 (0%) 87 92	12, 27, 48, 87	8 (3%)
2	B	216/221 (97%)	-0.14	2 (0%) 81 87	14, 29, 49, 76	5 (2%)
2	D	220/221 (99%)	-0.20	1 (0%) 87 92	14, 29, 48, 73	6 (2%)
3	E	12/12 (100%)	0.10	0 100 100	25, 33, 51, 65	0
3	F	12/12 (100%)	0.03	1 (8%) 17 18	24, 32, 53, 70	0
All	All	886/892 (99%)	-0.22	6 (0%) 84 90	12, 28, 49, 87	30 (3%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	139	GLY	4.2
1	C	213	ALA	3.9
2	B	133	SER	3.8
1	A	213	ALA	3.6
3	F	12	CYS	3.4
2	D	137	THR	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	A	301	5/5	0.92	0.13	31,44,55,55	5
4	PO4	B	301	5/5	0.93	0.08	40,44,49,49	5
4	PO4	C	301	5/5	0.93	0.12	20,35,39,39	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.