



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 01:19 AM UTC

PDB ID : 9THO / pdb_00009tho
Title : Crystal structure of the adduct formed upon reaction of [V(IV)O(acetylacetonate)₂] with human serum transferrin with Fe(III) bound at the C-lobe only
Authors : Paolillo, M.; Ferraro, G.; Banneville, A.S.; Cornaciu, I.; Pica, A.; Merlino, A.
Deposited on : 2025-12-03
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	679	10% 83% 14% ..

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 5529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	670	5270	3306	916	1001	47	0	8	0

- Molecule 2 is BICARBONATE ION (CCD ID: BCT) (formula: CHO_3).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		
2	AAA	1	Total	C	O	0	0
			4	1	3		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			6	3	3		
3	AAA	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	Fe	0	0
			1	1		

- Molecule 5 is MALONIC ACID (CCD ID: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is bis(oxidanyl)-oxidanylidene-(oxidanyl(oxidanylidene)vanadio)oxy-vanadium (CCD ID: A1JJT) (formula: $H_3O_6V_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	O	V	0	0
			8	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	AAA	187	Total 187	O 187	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serotransferrin



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	136.13Å 157.69Å 107.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.85 – 2.55 78.85 – 2.55	Depositor EDS
% Data completeness (in resolution range)	96.2 (78.85-2.55) 96.2 (78.85-2.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.55Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.198 , 0.249 0.203 , 0.250	Depositor DCC
R_{free} test set	1828 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5529	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FE, A1JJT, BCT, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.08	0/5387	1.44	3/7276 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	AAA	646	THR	CA-CB-OG1	-5.36	101.57	109.60
1	AAA	452	THR	CA-C-N	5.00	131.10	121.54
1	AAA	452	THR	C-N-CA	5.00	131.10	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	157	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	5270	0	5078	64	0
2	AAA	44	0	0	1	0
3	AAA	12	0	16	0	0
4	AAA	1	0	0	0	0
5	AAA	7	0	2	0	0
6	AAA	8	0	0	0	0
7	AAA	187	0	0	5	0
All	All	5529	0	5096	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:425:GLY:HA2	1:AAA:584:ASN:OD1	1.64	0.97
1:AAA:663:ARG:HD3	1:AAA:671:LEU:HD21	1.75	0.69
1:AAA:374:THR:HG21	1:AAA:395:PHE:CD2	2.31	0.64
1:AAA:163:ASP:OD2	1:AAA:166:ASP:OD2	2.17	0.63
1:AAA:538:VAL:HB	1:AAA:539:PRO:HD3	1.79	0.63
1:AAA:3:ASP:CG	1:AAA:5:THR:HG22	2.29	0.58
1:AAA:26:MET:CE	1:AAA:274:PHE:CD2	2.86	0.58
1:AAA:421:THR:HG21	7:AAA:968:HOH:O	2.04	0.58
1:AAA:663:ARG:CD	1:AAA:671:LEU:HD21	2.34	0.58
1:AAA:188:TYR:CZ	1:AAA:206:LYS:HE2	2.39	0.57
1:AAA:23:ARG:HG3	1:AAA:37:VAL:O	2.04	0.57
1:AAA:524:LEU:HB2	1:AAA:531:ALA:HB2	1.86	0.57
1:AAA:188:TYR:OH	1:AAA:206:LYS:HE2	2.05	0.56
1:AAA:109:MET:HE1	1:AAA:243:LEU:HD21	1.87	0.56
1:AAA:26:MET:CE	1:AAA:274:PHE:HD2	2.20	0.54
1:AAA:109:MET:HE2	1:AAA:226:LEU:HB3	1.88	0.54
1:AAA:600:ILE:O	1:AAA:604:GLN:HG2	2.09	0.53
1:AAA:374:THR:HG21	1:AAA:395:PHE:CE2	2.44	0.53
1:AAA:95:TYR:OH	1:AAA:246:VAL:HG11	2.09	0.53
1:AAA:462:ILE:HB	1:AAA:463:PRO:HD3	1.91	0.53
1:AAA:86:GLY:O	1:AAA:300:HIS:CD2	2.62	0.52
1:AAA:95:TYR:CZ	1:AAA:246:VAL:HG11	2.44	0.52
1:AAA:350:HIS:CE1	7:AAA:811:HOH:O	2.65	0.50
1:AAA:563:CYS:HB2	1:AAA:565:ASP:OD1	2.10	0.50
1:AAA:79:PRO:HB3	1:AAA:250:THR:HG21	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:26:MET:HE2	1:AAA:274:PHE:CE2	2.47	0.50
1:AAA:460:TRP:O	1:AAA:464:MET:HB2	2.12	0.49
1:AAA:4:LYS:HA	1:AAA:262:LEU:HD21	1.93	0.49
1:AAA:26:MET:HE1	1:AAA:274:PHE:CD2	2.48	0.49
1:AAA:109:MET:CE	1:AAA:243:LEU:HD21	2.41	0.49
1:AAA:354:LYS:HD3	1:AAA:630:LEU:HD22	1.95	0.48
1:AAA:26:MET:HE1	1:AAA:274:PHE:HD2	1.79	0.48
1:AAA:109:MET:HE2	1:AAA:226:LEU:CB	2.43	0.48
1:AAA:377:CYS:HB3	1:AAA:389:MET:SD	2.54	0.47
1:AAA:452:THR:OG1	1:AAA:459:GLY:HA3	2.15	0.47
1:AAA:538:VAL:HG11	1:AAA:571:VAL:HG21	1.97	0.47
1:AAA:26:MET:HE2	1:AAA:274:PHE:HE2	1.78	0.46
1:AAA:418:CYS:O	1:AAA:420:ASP:N	2.49	0.46
1:AAA:624:SER:OG	1:AAA:633:ASP:OD1	2.30	0.46
1:AAA:551:ALA:HA	1:AAA:554:LEU:CD1	2.45	0.46
1:AAA:281:GLU:HG3	1:AAA:282:PHE:N	2.31	0.46
1:AAA:4:LYS:O	1:AAA:35:PRO:HA	2.18	0.44
1:AAA:26:MET:CE	1:AAA:274:PHE:CE2	3.00	0.44
1:AAA:382:MET:SD	1:AAA:402:CYS:HB3	2.58	0.43
1:AAA:69:ASP:OD1	1:AAA:327:ARG:NH2	2.49	0.43
1:AAA:579:LEU:HD12	7:AAA:954:HOH:O	2.18	0.43
1:AAA:276:LYS:HD2	2:AAA:712:BCT:O1	2.18	0.43
1:AAA:407:VAL:HG12	1:AAA:594:GLU:HG3	2.00	0.43
1:AAA:483:GLY:HA2	1:AAA:497:LEU:HD12	2.01	0.42
1:AAA:574:TYR:CD1	1:AAA:575:ALA:N	2.87	0.42
1:AAA:677:ARG:NH1	7:AAA:817:HOH:O	2.51	0.42
1:AAA:392:ASP:HA	1:AAA:585:HIS:CD2	2.55	0.42
1:AAA:68:TYR:CG	1:AAA:323[A]:ILE:HD13	2.55	0.42
1:AAA:68:TYR:CD2	1:AAA:323[A]:ILE:HD13	2.55	0.41
1:AAA:493:SER:HA	1:AAA:496:LYS:HE2	2.02	0.41
1:AAA:551:ALA:HA	1:AAA:554:LEU:HD12	2.02	0.41
1:AAA:624:SER:HB3	1:AAA:629:LEU:HB2	2.01	0.41
1:AAA:383:ASN:OD1	1:AAA:383:ASN:C	2.63	0.41
1:AAA:418:CYS:C	1:AAA:420:ASP:N	2.79	0.41
1:AAA:452:THR:HA	1:AAA:516:GLY:O	2.20	0.41
1:AAA:352:ARG:O	1:AAA:352:ARG:HD2	2.21	0.41
1:AAA:535:HIS:CE1	1:AAA:536:GLN:OE1	2.74	0.40
1:AAA:354:LYS:NZ	7:AAA:823:HOH:O	2.54	0.40
1:AAA:535:HIS:HB2	1:AAA:574:TYR:CG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	672/679 (99%)	620 (92%)	48 (7%)	4 (1%)	21	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	419	GLU
1	AAA	436	ALA
1	AAA	614	ASP
1	AAA	634	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	572/572 (100%)	540 (94%)	32 (6%)	19	29

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	5	THR
1	AAA	29	VAL
1	AAA	71	TYR
1	AAA	89	GLU
1	AAA	182	LEU
1	AAA	249	HIS
1	AAA	260	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	AAA	262	LEU
1	AAA	291	LYS
1	AAA	327	ARG
1	AAA	361	ASN
1	AAA	365	LYS
1	AAA	407	VAL
1	AAA	420	ASP
1	AAA	438	ASP
1	AAA	454	VAL
1	AAA	477	ASP
1	AAA	489	LYS
1	AAA	496	LYS
1	AAA	556[A]	GLU
1	AAA	556[B]	GLU
1	AAA	563	CYS
1	AAA	571	VAL
1	AAA	607	LEU
1	AAA	624	SER
1	AAA	627	LYS
1	AAA	629	LEU
1	AAA	630	LEU
1	AAA	632	ARG
1	AAA	637	CYS
1	AAA	648	GLU
1	AAA	666	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 1 is monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	AAA	713	-	5,5,5	0.20	0	5,5,5	0.47	0
2	BCT	AAA	702	-	3,3,3	1.36	0	2,3,3	1.24	0
2	BCT	AAA	709	-	3,3,3	0.54	0	2,3,3	1.55	0
5	MLA	AAA	715	4	6,6,6	0.49	0	7,7,7	1.85	3 (42%)
6	A1JJT	AAA	716	1	0,7,7	-	-	-		
2	BCT	AAA	710	-	3,3,3	1.03	0	2,3,3	1.38	0
2	BCT	AAA	704	-	3,3,3	0.84	0	2,3,3	1.59	1 (50%)
2	BCT	AAA	711	-	3,3,3	0.78	0	2,3,3	1.89	1 (50%)
3	GOL	AAA	708	-	5,5,5	0.35	0	5,5,5	0.69	0
2	BCT	AAA	707	-	3,3,3	1.40	0	2,3,3	1.81	1 (50%)
2	BCT	AAA	712	-	3,3,3	0.93	0	2,3,3	1.25	0
2	BCT	AAA	706	-	3,3,3	0.42	0	2,3,3	1.71	1 (50%)
2	BCT	AAA	703	-	3,3,3	0.84	0	2,3,3	1.39	0
2	BCT	AAA	701	-	3,3,3	0.79	0	2,3,3	1.44	0
2	BCT	AAA	705	-	3,3,3	0.72	0	2,3,3	1.80	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	AAA	708	-	-	2/4/4/4	-
3	GOL	AAA	713	-	-	0/4/4/4	-
5	MLA	AAA	715	4	-	0/4/4/4	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	AAA	715	MLA	O3A-C3-C2	-2.62	114.65	122.11
5	AAA	715	MLA	O1B-C1-C2	-2.47	115.07	122.11
2	AAA	711	BCT	O2-C-O1	2.45	125.95	119.68
2	AAA	705	BCT	O2-C-O1	2.32	125.62	119.68
2	AAA	707	BCT	O2-C-O1	2.22	125.36	119.68
2	AAA	704	BCT	O2-C-O1	2.21	125.33	119.68
5	AAA	715	MLA	O3B-C3-C2	2.04	120.83	114.51
2	AAA	706	BCT	O2-C-O1	2.02	124.83	119.68

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AAA	708	GOL	C1-C2-C3-O3
3	AAA	708	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	712	BCT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	670/679 (98%)	0.40	65 (9%) 13 12	5, 27, 73, 120	8 (1%)

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	424	ALA	9.9
1	AAA	422	PRO	7.9
1	AAA	606[A]	HIS	5.9
1	AAA	613	THR	5.0
1	AAA	417	ASN	4.7
1	AAA	412	TYR	4.5
1	AAA	88	LYS	4.5
1	AAA	623	ARG	4.4
1	AAA	611	ASN	4.4
1	AAA	612	VAL	4.2
1	AAA	349	HIS	4.2
1	AAA	338	GLU	4.2
1	AAA	340	LYS	4.0
1	AAA	421	THR	4.0
1	AAA	350	HIS	3.9
1	AAA	333	GLU	3.8
1	AAA	423	GLU	3.7
1	AAA	667	THR	3.7
1	AAA	666	SER	3.7
1	AAA	419	GLU	3.7
1	AAA	363	VAL	3.6
1	AAA	615	CYS	3.5
1	AAA	617	GLY	3.5
1	AAA	619	PHE	3.5
1	AAA	618	ASN	3.3
1	AAA	348	SER	3.1
1	AAA	413	ASN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	602	ARG	3.1
1	AAA	503	LEU	3.0
1	AAA	418	CYS	2.9
1	AAA	643	ASP	2.9
1	AAA	640	LYS	2.8
1	AAA	332	PRO	2.8
1	AAA	607	LEU	2.8
1	AAA	339	CYS	2.8
1	AAA	625	GLU	2.8
1	AAA	608	PHE	2.7
1	AAA	89	GLU	2.7
1	AAA	642	HIS	2.7
1	AAA	365	LYS	2.7
1	AAA	360	VAL	2.6
1	AAA	610	SER	2.6
1	AAA	657	LYS	2.5
1	AAA	438	ASP	2.5
1	AAA	569	LYS	2.5
1	AAA	603	GLN	2.5
1	AAA	597	VAL	2.5
1	AAA	361	ASN	2.5
1	AAA	87	SER	2.5
1	AAA	622	PHE	2.4
1	AAA	3	ASP	2.4
1	AAA	364	GLY	2.4
1	AAA	182	LEU	2.4
1	AAA	442	ASP	2.4
1	AAA	616	SER	2.3
1	AAA	601	LEU	2.3
1	AAA	614	ASP	2.3
1	AAA	644	ARG	2.2
1	AAA	409	ALA	2.2
1	AAA	575	ALA	2.2
1	AAA	420	ASP	2.2
1	AAA	571	VAL	2.1
1	AAA	375[A]	GLU	2.1
1	AAA	634	ASP	2.0
1	AAA	632	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BCT	AAA	711	4/4	0.72	0.26	59,65,70,80	0
2	BCT	AAA	709	4/4	0.79	0.18	36,40,46,49	0
3	GOL	AAA	708	6/6	0.79	0.21	40,48,53,54	0
2	BCT	AAA	706	4/4	0.80	0.19	41,46,47,54	0
2	BCT	AAA	705	4/4	0.81	0.25	65,68,69,72	0
2	BCT	AAA	710	4/4	0.82	0.23	52,53,59,64	0
2	BCT	AAA	712	4/4	0.86	0.18	47,56,59,60	0
2	BCT	AAA	703	4/4	0.87	0.15	45,48,54,56	0
2	BCT	AAA	704	4/4	0.93	0.10	30,30,30,37	0
5	MLA	AAA	715	7/7	0.93	0.10	17,18,20,23	0
3	GOL	AAA	713	6/6	0.94	0.09	20,21,22,23	0
2	BCT	AAA	702	4/4	0.94	0.09	32,37,39,41	0
2	BCT	AAA	707	4/4	0.95	0.08	32,32,33,38	0
6	A1JJT	AAA	716	8/8	0.97	0.12	20,29,39,45	8
2	BCT	AAA	701	4/4	0.98	0.05	22,24,25,25	0
4	FE	AAA	714	1/1	0.99	0.02	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1JJT AAA 716:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.