



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 02:18 PM EDT

PDB ID : 3TIA / pdb_00003tia
Title : Crystal structure of 1957 pandemic H2N2 neuraminidase complexed with laninamivir
Authors : Vavricka, C.J.; Li, Q.; Wu, Y.; Qi, J.; Wang, M.; Liu, Y.; Gao, F.; Liu, J.; Feng, E.; He, J.; Wang, J.; Liu, H.; Jiang, H.; Gao, G.F.
Deposited on : 2011-08-20
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	
1	B	469	
1	C	469	
1	D	469	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	2	50% 50%
2	H	2	50% 50%
2	J	2	50% 50%
3	F	4	75% 25%
3	G	4	75% 25%
3	I	4	75% 25%
3	K	4	75% 25%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 14414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	1	0
			3023	1866	543	591	23			
1	B	388	Total	C	N	O	S	0	0	0
			3017	1863	542	589	23			
1	C	388	Total	C	N	O	S	0	1	0
			3025	1867	544	591	23			
1	D	388	Total	C	N	O	S	0	0	0
			3017	1863	542	589	23			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	G	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	I	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	K	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		

- Molecule 5 is 5-acetamido-2,6-anhydro-4-carbamimidamido-3,4,5-trideoxy-7-O-methyl-D-glycero-D-galacto-non-2-enonic acid (CCD ID: LNV) (formula: C₁₃H₂₂N₄O₇).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			24	13	4	7		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			24	13	4	7		
5	C	1	Total	C	N	O	0	0
			24	13	4	7		
5	D	1	Total	C	N	O	0	0
			24	13	4	7		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	475	Total	O	0	0
			475	475		
7	B	494	Total	O	0	0
			494	494		
7	C	488	Total	O	0	0
			488	488		
7	D	477	Total	O	0	0
			477	477		

Chain D:  72% 10% 17%

MET ASN PRO ASN GLN LYS ILE THR THR ILE GLY SER VAL SER LEU THR ALA THR VAL CYS PHE LEU MET GLN ILE ALA ILE LEU ALA THR THR VAL THR LEU THR THR LEU HIS PHE LYS GLN HIS CYS ASP SER PRO ALA SER ASN GLN VAL MET PRO CYS GLU PRO ILE ILE ILE ARG

ASN ILE THR GLU ILE VAL TYR LEU ASN ASN THR THR ILE GLU LYS ILE ILE CYS PRO LYS VAL V82 G112 Y121 D125 P126 Q131 M142 K143 H144 H155 G164 L169 V174 A177 H184 D185 G186 H191 V192 C193 Q226 F227 S228 N234

V240 S245 R249 I254 L255 H264 I265 Q273 H274 C289 I290 D293 N294 W295 K296 R300 P301 D329 N334 S335 N336 C337 R344 G345 T346 Q347 E369 N387 V392 N393 V424 G461 A462 N463 I469

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%

NAG1 NAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%

NAG1 NAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%

NAG1 NAG2

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  75% 25%

NAG1 NAG2 BMA3 MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  75% 25%

NAG1 NAG2 BMA3 MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  75% 25%

MAG1
MAG2
BMA3
MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  75% 25%

MAG1
MAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.93Å 140.31Å 89.89Å 90.00° 101.54° 90.00°	Depositor
Resolution (Å)	42.99 – 1.80 42.99 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.4 (42.99-1.80) 99.7 (42.99-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.52 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.152 , 0.170 0.153 , 0.171	Depositor DCC
R_{free} test set	10154 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	11.0	Xtriage
Anisotropy	0.494	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.449 for l,-k,h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	14414	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LNV, MAN, BMA, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/3096	0.72	0/4203
1	B	0.34	0/3090	0.71	1/4195 (0.0%)
1	C	0.33	0/3098	0.72	1/4206 (0.0%)
1	D	0.34	0/3090	0.73	1/4195 (0.0%)
All	All	0.34	0/12374	0.72	3/16799 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	226	GLN	N-CA-C	5.98	117.88	111.36
1	B	226	GLN	N-CA-C	5.38	117.22	111.36
1	C	226	GLN	N-CA-C	5.31	117.15	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3023	0	2836	31	0
1	B	3017	0	2832	41	0
1	C	3025	0	2837	47	0
1	D	3017	0	2832	44	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	28	0	25	0	0
2	H	28	0	25	1	0
2	J	28	0	25	0	0
3	F	50	0	43	0	0
3	G	50	0	43	0	0
3	I	50	0	43	0	0
3	K	50	0	43	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	24	0	0	1	0
5	B	24	0	0	0	0
5	C	24	0	0	0	0
5	D	24	0	0	0	0
6	B	14	0	13	0	0
7	A	475	0	0	2	0
7	B	494	0	0	5	0
7	C	488	0	0	2	0
7	D	477	0	0	3	0
All	All	14414	0	11597	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:226:GLN:HE21	1:B:240:VAL:H	1.22	0.84
1:C:226:GLN:HE21	1:C:240:VAL:H	1.25	0.83
1:C:334:ASN:HA	1:C:387:ASN:HD21	1.44	0.82
1:C:344:ARG:HH12	1:C:369:GLU:CD	1.86	0.81
1:A:144:HIS:HE2	1:D:463:ASN:H	1.23	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/469 (82%)	370 (96%)	17 (4%)	0	100	100
1	B	386/469 (82%)	372 (96%)	14 (4%)	0	100	100
1	C	387/469 (82%)	371 (96%)	16 (4%)	0	100	100
1	D	386/469 (82%)	371 (96%)	15 (4%)	0	100	100
All	All	1546/1876 (82%)	1484 (96%)	62 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/414 (82%)	335 (99%)	4 (1%)	63	57
1	B	338/414 (82%)	334 (99%)	4 (1%)	63	57
1	C	339/414 (82%)	334 (98%)	5 (2%)	57	49
1	D	338/414 (82%)	333 (98%)	5 (2%)	57	49
All	All	1354/1656 (82%)	1336 (99%)	18 (1%)	61	54

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	249	ARG
1	D	424	VAL
1	D	387	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	142	ASN
1	D	142	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	226	GLN
1	D	273	GLN
1	D	432	GLN
1	B	264	HIS
1	B	226	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	2,1	14,14,15	0.59	0	17,19,21	0.86	1 (5%)
2	NAG	E	2	2	14,14,15	0.53	0	17,19,21	0.87	0
3	NAG	F	1	3,1	14,14,15	0.49	0	17,19,21	1.12	3 (17%)
3	NAG	F	2	3	14,14,15	0.48	0	17,19,21	0.72	0
3	BMA	F	3	3	11,11,12	0.52	0	15,15,17	0.61	0
3	MAN	F	4	3	11,11,12	0.65	0	15,15,17	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	G	1	3,1	14,14,15	0.57	0	17,19,21	1.03	2 (11%)
3	NAG	G	2	3	14,14,15	0.46	0	17,19,21	0.66	0
3	BMA	G	3	3	11,11,12	0.55	0	15,15,17	0.55	0
3	MAN	G	4	3	11,11,12	0.54	0	15,15,17	0.62	0
2	NAG	H	1	2,1	14,14,15	0.57	0	17,19,21	0.81	0
2	NAG	H	2	2	14,14,15	0.55	0	17,19,21	0.62	0
3	NAG	I	1	3,1	14,14,15	0.49	0	17,19,21	1.19	2 (11%)
3	NAG	I	2	3	14,14,15	0.43	0	17,19,21	0.83	0
3	BMA	I	3	3	11,11,12	0.61	0	15,15,17	0.60	0
3	MAN	I	4	3	11,11,12	0.57	0	15,15,17	0.63	0
2	NAG	J	1	2,1	14,14,15	0.57	0	17,19,21	0.73	0
2	NAG	J	2	2	14,14,15	0.57	0	17,19,21	0.91	1 (5%)
3	NAG	K	1	3,1	14,14,15	0.51	0	17,19,21	0.93	1 (5%)
3	NAG	K	2	3	14,14,15	0.46	0	17,19,21	0.64	0
3	BMA	K	3	3	11,11,12	0.61	0	15,15,17	0.57	0
3	MAN	K	4	3	11,11,12	0.53	0	15,15,17	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
3	MAN	G	4	3	-	0/2/19/22	0/1/1/1
2	NAG	H	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	2/6/23/26	0/1/1/1
3	NAG	I	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
3	MAN	I	4	3	-	0/2/19/22	0/1/1/1
2	NAG	J	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	J	2	2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	K	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	K	2	3	-	0/6/23/26	0/1/1/1
3	BMA	K	3	3	-	0/2/19/22	0/1/1/1
3	MAN	K	4	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	1	NAG	C1-O5-C5	3.07	116.30	112.19
3	I	1	NAG	C2-N2-C7	-2.73	119.24	122.90
3	F	1	NAG	C1-O5-C5	2.71	115.82	112.19
3	F	1	NAG	C2-N2-C7	-2.46	119.60	122.90
3	G	1	NAG	C2-N2-C7	-2.45	119.62	122.90

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	1	NAG	C8-C7-N2-C2
2	H	2	NAG	O5-C5-C6-O6
3	F	1	NAG	O7-C7-N2-C2
2	H	2	NAG	C4-C5-C6-O6
3	I	1	NAG	C8-C7-N2-C2


There are no ring outliers.


1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	LNV	C	801	-	23,24,24	3.46	9 (39%)	22,33,33	1.94	9 (40%)
5	LNV	B	801	-	23,24,24	3.51	10 (43%)	22,33,33	2.08	6 (27%)
6	NAG	B	601	1	14,14,15	0.56	0	17,19,21	0.91	1 (5%)
5	LNV	D	801	-	23,24,24	3.49	10 (43%)	22,33,33	2.01	6 (27%)
5	LNV	A	801	-	23,24,24	3.31	11 (47%)	22,33,33	2.03	8 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LNV	C	801	-	-	1/24/40/40	0/1/1/1
5	LNV	B	801	-	-	0/24/40/40	0/1/1/1
6	NAG	B	601	1	-	0/6/23/26	0/1/1/1
5	LNV	D	801	-	-	0/24/40/40	0/1/1/1
5	LNV	A	801	-	-	0/24/40/40	0/1/1/1

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	801	LNV	C3-C2	10.52	1.51	1.33
5	B	801	LNV	C3-C2	10.51	1.51	1.33
5	C	801	LNV	C3-C2	10.42	1.51	1.33
5	A	801	LNV	C3-C2	10.30	1.51	1.33
5	D	801	LNV	O8-C8	-6.55	1.29	1.43

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	801	LNV	O6-C2-C3	-5.63	118.32	124.62
5	D	801	LNV	O6-C2-C3	-5.18	118.82	124.62
5	A	801	LNV	O6-C2-C3	-4.96	119.07	124.62
5	C	801	LNV	O6-C2-C3	-4.49	119.59	124.62
5	D	801	LNV	C6-C5-N5	4.05	117.36	110.91

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	801	LNV	O1A-C1-C2-C3


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	801	LNV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/469 (82%)	-1.59	0 100 100	4, 11, 23, 53	1 (0%)
1	B	388/469 (82%)	-1.56	0 100 100	3, 11, 24, 55	0
1	C	388/469 (82%)	-1.59	0 100 100	4, 11, 24, 54	1 (0%)
1	D	388/469 (82%)	-1.57	0 100 100	4, 11, 24, 50	0
All	All	1552/1876 (82%)	-1.57	0 100 100	3, 11, 24, 55	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	H	2	14/15	0.96	0.09	75,81,97,98	0
2	NAG	E	2	14/15	0.97	0.09	66,74,89,91	0
2	NAG	J	2	14/15	0.97	0.07	41,61,77,78	0
2	NAG	E	1	14/15	0.99	0.03	22,31,40,45	0
2	NAG	J	1	14/15	0.99	0.04	24,34,39,43	0
2	NAG	H	1	14/15	0.99	0.03	18,29,41,46	0
3	NAG	F	2	14/15	0.99	0.02	12,15,19,20	0
3	BMA	F	3	11/12	0.99	0.02	12,17,22,22	0
3	MAN	F	4	11/12	0.99	0.03	16,20,23,25	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BMA	G	3	11/12	0.99	0.03	10,13,16,20	0
3	MAN	G	4	11/12	0.99	0.02	15,17,21,22	0
3	BMA	I	3	11/12	0.99	0.02	11,15,19,21	0
3	NAG	K	2	14/15	0.99	0.03	9,12,16,17	0
3	BMA	K	3	11/12	0.99	0.02	9,14,18,19	0
3	MAN	K	4	11/12	0.99	0.02	15,16,21,24	0
3	NAG	I	2	14/15	1.00	0.02	12,14,21,22	0
3	NAG	G	2	14/15	1.00	0.02	10,12,16,18	0
3	MAN	I	4	11/12	1.00	0.02	17,19,22,22	0
3	NAG	K	1	14/15	1.00	0.02	9,12,31,33	0
3	NAG	F	1	14/15	1.00	0.02	9,15,34,38	0
3	NAG	G	1	14/15	1.00	0.02	10,12,28,29	0
3	NAG	I	1	14/15	1.00	0.02	8,15,34,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain G:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain K:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	B	601	14/15	0.98	0.06	38,50,60,70	0
4	CA	B	501	1/1	1.00	0.01	9,9,9,9	0
4	CA	C	501	1/1	1.00	0.01	9,9,9,9	0
4	CA	D	501	1/1	1.00	0.01	10,10,10,10	0
5	LNV	A	801	24/24	1.00	0.02	6,9,13,14	0
5	LNV	B	801	24/24	1.00	0.02	5,8,12,13	0
5	LNV	C	801	24/24	1.00	0.01	6,8,13,14	0
5	LNV	D	801	24/24	1.00	0.02	5,8,11,13	0
4	CA	A	501	1/1	1.00	0.01	9,9,9,9	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LNV A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LNV B 801:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LNV C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.