



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 12:52 PM UTC

PDB ID : 3TIB / pdb_00003tib
Title : Crystal structure of 1957 pandemic H2N2 neuraminidase complexed with laninamivir octanoate
Authors : Vavricka, C.J.; Li, Q.; Wu, Y.; Qi, J.; Wang, M.; Liu, Y.; Gao, F.; Liu, J.; Feng, E.; He, J.; Wang, J.; Liu, H.; Jiang, H.; Gao, G.F.
Deposited on : 2011-08-20
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	
1	B	469	
1	C	469	
1	D	469	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	2	 100%
2	G	2	 100%
2	I	2	 100%
2	K	2	 100%
3	F	4	 75% 25%
3	H	4	 75% 25%
3	J	4	 50% 50%
3	L	4	 75% 25%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	0	1	0
			3023	1866	543	591	23			
1	B	388	Total	C	N	O	S	0	1	0
			3023	1866	543	591	23			
1	C	388	Total	C	N	O	S	0	0	0
			3017	1863	542	589	23			
1	D	388	Total	C	N	O	S	0	1	0
			3023	1866	543	591	23			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	K	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	H	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	J	4	Total	C	N	O	0	0	0
			50	28	2	20			
3	L	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		

- Molecule 5 is 5-acetamido-2,6-anhydro-4-carbamimidamido-3,4,5-trideoxy-7-O-methyl-9-O-octanoyl-D-glycero-D-galacto-non-2-enonic acid (CCD ID: LVO) (formula: C₂₁H₃₆N₄O₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			26	14	4	8		
5	B	1	Total	C	N	O	0	0
			26	14	4	8		
5	C	1	Total	C	N	O	0	0
			26	14	4	8		
5	D	1	Total	C	N	O	0	0
			26	14	4	8		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	358	Total	O	0	0
			358	358		
6	B	347	Total	O	0	0
			347	347		
6	C	333	Total	O	0	0
			333	333		
6	D	356	Total	O	0	0
			356	356		

- Molecule 1: Neuraminidase

H264	H273	H274	H284	H295	H296	D304	T325	D329	N334	N335	N336	C337	R344	G348	W361	R364	E369	E376	K378	N387	Q391	V392	G461	L469														
ASN	THR	GLU	ILE	VAL	TYR	LEU	ASN	ASN	THR	THR	ILE	GLU	LYS	GLU	ILE	ALA	CYS	PRO	THR	VAL	CYS	PHE	LEU	HIS	GLN	ASN	GLN	VAL	MET	PRO	CYS	GLU	PRO	ILE	ILE	ILE	GLU	PRO

- Molecule 1: Neuraminidase


A246	Q273	H274	C278	R288	C289	N294	K296	D304	T325	D329	N334	N336	C337	R344	G348	W361	R364	E369	E375	K378	N387	S388	V392	N393	G454	G461	I469																	
ASN	THR	GLU	VAL	LEU	ASN	THR	THR	GLU	LEU	PRO	CYS	VAL	VAL	VAL	VAL	Q131	M142	K143	H144	H155	M161	G164	C175	I176	A177	H184	D185	G186	C193	K199	T202	Q226	V231	V240										
MET	ASN	PRO	ASN	GLN	LYS	ILE	ILE	THR	GLY	THR	VAL	SER	VAL	THR	ALA	THR	THR	VAL	THR	LEU	HIS	PHE	LYS	GLN	HIS	GLU	CYS	ASP	SER	PRO	ALA	SER	ASN	GLN	VAL	MET	PRO	CYS	GLU	PRO	ILE	ILE	GLU	VAL

- Molecule 1: Neuraminidase

[illegible]

- Molecule 1: Neuraminidase

Chain D:  73% 9% 17%




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  75% 25%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  75% 25%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.45Å 140.18Å 90.30Å 90.00° 101.22° 90.00°	Depositor
Resolution (Å)	43.17 – 2.20 43.17 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.1 (43.17-2.20) 99.1 (43.17-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.155 , 0.191 0.161 , 0.196	Depositor DCC
R_{free} test set	5561 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 20.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.467 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13900	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LVO, MAN, BMA, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/3096	0.71	1/4203 (0.0%)
1	B	0.32	0/3096	0.72	1/4203 (0.0%)
1	C	0.32	0/3090	0.71	0/4195
1	D	0.31	0/3096	0.71	1/4203 (0.0%)
All	All	0.32	0/12378	0.72	3/16804 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	GLN	N-CA-C	5.59	117.46	111.36
1	D	226	GLN	N-CA-C	5.49	117.35	111.36
1	A	226	GLN	N-CA-C	5.47	117.32	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3023	0	2836	37	0
1	B	3023	0	2836	32	0
1	C	3017	0	2832	38	0
1	D	3023	0	2836	38	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	28	0	25	0	0
2	G	28	0	25	0	0
2	I	28	0	25	0	0
2	K	28	0	25	0	0
3	F	50	0	43	1	0
3	H	50	0	43	0	0
3	J	50	0	43	1	0
3	L	50	0	43	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	26	0	0	1	0
5	B	26	0	0	1	0
5	C	26	0	0	1	0
5	D	26	0	0	1	0
6	A	358	0	0	5	0
6	B	347	0	0	2	0
6	C	333	0	0	2	0
6	D	356	0	0	4	0
All	All	13900	0	11612	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:226:GLN:HE21	1:C:240:VAL:H	1.16	0.92
1:D:226:GLN:HE21	1:D:240:VAL:H	1.19	0.91
1:A:144:HIS:HE2	1:C:463:ASN:H	1.21	0.89
1:B:226:GLN:HE21	1:B:240:VAL:H	1.19	0.88
1:A:226:GLN:HE21	1:A:240:VAL:H	1.20	0.85

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/469 (82%)	371 (96%)	16 (4%)	0	100	100
1	B	387/469 (82%)	370 (96%)	17 (4%)	0	100	100
1	C	386/469 (82%)	371 (96%)	15 (4%)	0	100	100
1	D	387/469 (82%)	372 (96%)	15 (4%)	0	100	100
All	All	1547/1876 (82%)	1484 (96%)	63 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/414 (82%)	331 (98%)	8 (2%)	43	58
1	B	339/414 (82%)	329 (97%)	10 (3%)	37	51
1	C	338/414 (82%)	331 (98%)	7 (2%)	47	63
1	D	339/414 (82%)	332 (98%)	7 (2%)	47	63
All	All	1355/1656 (82%)	1323 (98%)	32 (2%)	43	58

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	231	VAL
1	D	364	ARG
1	B	231	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	202	THR
1	D	369	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	274	HIS
1	D	356	ASN
1	B	274	HIS
1	B	273	GLN
1	D	387	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	2,1	14,14,15	0.58	0	17,19,21	0.74	0
2	NAG	E	2	2	14,14,15	0.47	0	17,19,21	0.72	0
3	NAG	F	1	3,1	14,14,15	0.51	0	17,19,21	0.79	0
3	NAG	F	2	3	14,14,15	0.50	0	17,19,21	0.74	0
3	BMA	F	3	3	11,11,12	0.59	0	15,15,17	0.73	0
3	MAN	F	4	3	11,11,12	0.66	0	15,15,17	0.65	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	G	1	2,1	14,14,15	0.58	0	17,19,21	0.78	0
2	NAG	G	2	2	14,14,15	0.50	0	17,19,21	0.75	0
3	NAG	H	1	3,1	14,14,15	0.46	0	17,19,21	1.19	2 (11%)
3	NAG	H	2	3	14,14,15	0.46	0	17,19,21	0.73	0
3	BMA	H	3	3	11,11,12	0.62	0	15,15,17	0.64	0
3	MAN	H	4	3	11,11,12	0.63	0	15,15,17	0.71	0
2	NAG	I	1	2,1	14,14,15	0.57	0	17,19,21	0.92	0
2	NAG	I	2	2	14,14,15	0.52	0	17,19,21	0.61	0
3	NAG	J	1	3,1	14,14,15	0.57	0	17,19,21	0.89	1 (5%)
3	NAG	J	2	3	14,14,15	0.48	0	17,19,21	0.74	0
3	BMA	J	3	3	11,11,12	0.61	0	15,15,17	0.63	0
3	MAN	J	4	3	11,11,12	0.54	0	15,15,17	0.65	0
2	NAG	K	1	2,1	14,14,15	0.55	0	17,19,21	0.95	0
2	NAG	K	2	2	14,14,15	0.54	0	17,19,21	0.70	0
3	NAG	L	1	3,1	14,14,15	0.47	0	17,19,21	0.85	1 (5%)
3	NAG	L	2	3	14,14,15	0.52	0	17,19,21	0.73	0
3	BMA	L	3	3	11,11,12	0.62	0	15,15,17	0.73	0
3	MAN	L	4	3	11,11,12	0.59	0	15,15,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	1/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
2	NAG	G	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
3	NAG	H	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	0/2/19/22	0/1/1/1
3	MAN	H	4	3	-	0/2/19/22	0/1/1/1
2	NAG	I	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
3	NAG	J	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	0/6/23/26	0/1/1/1
3	BMA	J	3	3	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	J	4	3	-	0/2/19/22	0/1/1/1
2	NAG	K	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	K	2	2	-	0/6/23/26	0/1/1/1
3	NAG	L	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	0/6/23/26	0/1/1/1
3	BMA	L	3	3	-	0/2/19/22	0/1/1/1
3	MAN	L	4	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	1	NAG	C1-O5-C5	3.53	116.92	112.19
3	J	1	NAG	C1-O5-C5	2.49	115.52	112.19
3	H	1	NAG	C2-N2-C7	-2.12	120.06	122.90
3	L	1	NAG	C2-N2-C7	-2.02	120.19	122.90

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	H	1	NAG	C8-C7-N2-C2
3	H	1	NAG	O7-C7-N2-C2
3	J	3	BMA	C4-C5-C6-O6
3	F	3	BMA	C4-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	3	BMA	1	0
3	F	3	BMA	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	LVO	D	801	-	25,26,33	3.05	11 (44%)	23,35,43	3.20	14 (60%)
5	LVO	C	801	-	25,26,33	3.16	12 (48%)	23,35,43	3.16	11 (47%)
5	LVO	B	801	-	25,26,33	3.25	12 (48%)	23,35,43	3.07	11 (47%)
5	LVO	A	801	-	25,26,33	3.25	12 (48%)	23,35,43	3.32	11 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LVO	D	801	-	-	10/26/42/50	0/1/1/1
5	LVO	C	801	-	-	8/26/42/50	0/1/1/1
5	LVO	B	801	-	-	7/26/42/50	0/1/1/1
5	LVO	A	801	-	-	6/26/42/50	0/1/1/1

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	801	LVO	C3-C2	10.50	1.51	1.33
5	B	801	LVO	C3-C2	10.02	1.50	1.33
5	A	801	LVO	C3-C2	9.86	1.50	1.33
5	D	801	LVO	C3-C2	9.64	1.50	1.33
5	D	801	LVO	O8-C8	-5.70	1.31	1.43

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	801	LVO	O6-C2-C3	-9.36	114.15	124.62
5	A	801	LVO	O6-C2-C3	-8.97	114.58	124.62
5	B	801	LVO	O6-C2-C3	-8.35	115.28	124.62
5	C	801	LVO	O6-C2-C3	-7.93	115.74	124.62
5	C	801	LVO	C6-C5-N5	6.53	121.31	110.91

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	801	LVO	C5-C6-C7-C8
5	A	801	LVO	OAV-CAN-O9-C9
5	B	801	LVO	OAV-CAN-O9-C9
5	C	801	LVO	O1B-C1-C2-C3
5	C	801	LVO	C5-C6-C7-C8


There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	801	LVO	1	0
5	C	801	LVO	1	0
5	B	801	LVO	1	0
5	A	801	LVO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand LVO D 801





Ligand LVO B 801





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/469 (82%)	-1.79	0 100 100	3, 12, 25, 49	1 (0%)
1	B	388/469 (82%)	-1.79	0 100 100	4, 12, 25, 50	1 (0%)
1	C	388/469 (82%)	-1.77	0 100 100	4, 12, 26, 48	0
1	D	388/469 (82%)	-1.78	0 100 100	4, 12, 26, 53	1 (0%)
All	All	1552/1876 (82%)	-1.78	0 100 100	3, 12, 26, 53	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	2	14/15	0.98	0.05	54,63,82,84	0
2	NAG	G	2	14/15	0.98	0.05	56,64,80,83	0
2	NAG	K	2	14/15	0.98	0.06	56,72,80,83	0
2	NAG	E	1	14/15	0.99	0.03	26,35,46,52	0
2	NAG	I	1	14/15	0.99	0.03	23,33,42,48	0
2	NAG	I	2	14/15	0.99	0.04	51,67,78,84	0
2	NAG	K	1	14/15	0.99	0.04	26,33,47,50	0
2	NAG	G	1	14/15	0.99	0.03	24,33,43,47	0
3	BMA	F	3	11/12	0.99	0.03	16,20,26,30	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAN	F	4	11/12	0.99	0.02	15,20,24,25	0
3	BMA	H	3	11/12	0.99	0.02	16,20,26,26	0
3	MAN	H	4	11/12	0.99	0.02	17,20,24,30	0
3	BMA	J	3	11/12	0.99	0.02	11,18,25,26	0
3	MAN	J	4	11/12	0.99	0.02	14,18,21,22	0
3	BMA	L	3	11/12	0.99	0.02	12,17,20,29	0
3	MAN	L	4	11/12	0.99	0.03	16,18,23,23	0
3	NAG	J	1	14/15	1.00	0.01	12,16,32,35	0
3	NAG	J	2	14/15	1.00	0.01	11,14,18,19	0
3	NAG	H	1	14/15	1.00	0.02	12,17,33,33	0
3	NAG	H	2	14/15	1.00	0.01	14,17,22,24	0
3	NAG	L	1	14/15	1.00	0.02	12,16,31,33	0
3	NAG	L	2	14/15	1.00	0.02	10,15,18,20	0
3	NAG	F	1	14/15	1.00	0.02	15,21,33,38	0
3	NAG	F	2	14/15	1.00	0.01	14,16,24,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	LVO	C	801	26/33	0.99	0.03	6,11,23,36	0
4	CA	B	501	1/1	1.00	0.02	11,11,11,11	0
4	CA	C	501	1/1	1.00	0.02	11,11,11,11	0
4	CA	D	501	1/1	1.00	0.01	12,12,12,12	0
5	LVO	A	801	26/33	1.00	0.02	7,11,20,36	0
5	LVO	B	801	26/33	1.00	0.02	7,11,20,34	0
4	CA	A	501	1/1	1.00	0.01	10,10,10,10	0
5	LVO	D	801	26/33	1.00	0.02	8,12,20,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around LVO C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LVO A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around LVO B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LVO D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.