



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:01 PM UTC

PDB ID : 6TIG / pdb_00006tig
Title : Structure of the N terminal domain of Bc2L-C lectin (1-131) in complex with
Globo H (H-type 3) antigen
Authors : Varrot, A.; Bermeo, R.
Deposited on : 2019-11-22
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	134	
1	BBB	134	
1	CCC	134	
2	A	4	
3	B	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	5	 100%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	134	Total	C	N	O	S	0	2	0
			994	639	162	189	4			
1	BBB	134	Total	C	N	O	S	0	5	0
			1027	662	166	195	4			
1	CCC	134	Total	C	N	O	S	0	3	0
			1002	645	162	191	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-2	GLY	-	expression tag	UNP B4EH86
AAA	-1	HIS	-	expression tag	UNP B4EH86
BBB	-2	GLY	-	expression tag	UNP B4EH86
BBB	-1	HIS	-	expression tag	UNP B4EH86
CCC	-2	GLY	-	expression tag	UNP B4EH86
CCC	-1	HIS	-	expression tag	UNP B4EH86

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	A	4	Total	C	N	O	0	0	0
			47	26	1	20			

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	5	Total	C	N	O	0	0	0
			58	32	1	25			
3	C	5	Total	C	N	O	0	0	0
			58	32	1	25			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	104	Total	O	0	5
			109	109		
4	BBB	103	Total	O	0	3
			106	106		
4	CCC	80	Total	O	0	2
			82	82		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lectin

Chain AAA: 



- Molecule 1: Lectin

Chain BBB: 




- Molecule 1: Lectin

Chain CCC: 




- Molecule 2: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose

Chain A: 



- Molecule 3: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose

Chain B: 



- Molecule 3: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose

Chain C:

100%

GAL1
GLA2
NGA3
GAL4
FUC5

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	74.37Å 42.88Å 102.65Å 90.00° 96.01° 90.00°	Depositor
Resolution (Å)	37.10 – 1.90 37.10 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (37.10-1.90) 97.5 (37.10-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.165 , 0.227 0.176 , 0.228	Depositor DCC
R_{free} test set	1552 reflections (6.19%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3483	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, NGA, GLA, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.21	2/1018 (0.2%)	1.26	0/1393
1	BBB	1.20	4/1055 (0.4%)	1.29	2/1441 (0.1%)
1	CCC	1.26	1/1029 (0.1%)	1.32	3/1408 (0.2%)
All	All	1.22	7/3102 (0.2%)	1.29	5/4242 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	72	CYS	C-O	-8.12	1.14	1.23
1	AAA	108	LYS	C-O	5.76	1.30	1.23
1	BBB	41	ILE	C-O	5.44	1.29	1.24
1	BBB	103[A]	VAL	C-O	5.40	1.29	1.24
1	BBB	103[B]	VAL	C-O	5.40	1.29	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	95	ASP	CA-CB-CG	8.33	120.93	112.60
1	CCC	95	ASP	CA-CB-CG	7.66	120.26	112.60
1	BBB	95	ASP	CB-CA-C	-6.66	100.54	110.67
1	CCC	35	ASP	CA-CB-CG	6.61	119.21	112.60
1	CCC	95	ASP	CB-CA-C	-6.14	101.33	110.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	994	0	988	11	0
1	BBB	1027	0	1027	10	0
1	CCC	1002	0	1001	8	1
2	A	47	0	42	0	0
3	B	58	0	51	0	0
3	C	58	0	51	0	0
4	AAA	109	0	0	0	0
4	BBB	106	0	0	0	0
4	CCC	82	0	0	0	0
All	All	3483	0	3160	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:26:LEU:HB2	1:AAA:103[B]:VAL:HG22	1.75	0.68
1:CCC:2:LEU:HD22	1:CCC:126:ILE:HG12	1.80	0.63
1:CCC:26:LEU:HB2	1:CCC:103[B]:VAL:HG22	1.82	0.62
1:BBB:2:LEU:HD22	1:BBB:126:ILE:HG12	1.82	0.62
1:BBB:26:LEU:HB2	1:BBB:103[A]:VAL:HG22	1.81	0.62

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:CCC:32:GLY:O	1:CCC:32:GLY:O[2_556]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	134/134 (100%)	129 (96%)	5 (4%)	0	100	100
1	BBB	137/134 (102%)	133 (97%)	4 (3%)	0	100	100
1	CCC	135/134 (101%)	131 (97%)	4 (3%)	0	100	100
All	All	406/402 (101%)	393 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	106/106 (100%)	103 (97%)	3 (3%)	38	32
1	BBB	111/106 (105%)	109 (98%)	2 (2%)	51	50
1	CCC	108/106 (102%)	105 (97%)	3 (3%)	38	32
All	All	325/318 (102%)	317 (98%)	8 (2%)	42	37

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	CCC	124	SER
1	CCC	107	TRP
1	BBB	107	TRP
1	BBB	98	SER
1	CCC	98	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLA	A	1	2	12,12,12	1.05	1 (8%)	17,17,17	1.58	5 (29%)
2	NGA	A	2	2	14,14,15	1.04	1 (7%)	17,19,21	2.28	8 (47%)
2	GAL	A	3	2	11,11,12	1.23	2 (18%)	15,15,17	1.58	5 (33%)
2	FUC	A	4	2	10,10,11	1.66	3 (30%)	14,14,16	1.06	1 (7%)
3	GAL	B	1	3	12,12,12	0.84	0	17,17,17	1.36	4 (23%)
3	GLA	B	2	3	11,11,12	1.16	1 (9%)	15,15,17	3.44	10 (66%)
3	NGA	B	3	3	14,14,15	0.85	0	17,19,21	1.49	4 (23%)
3	GAL	B	4	3	11,11,12	1.08	1 (9%)	15,15,17	1.95	6 (40%)
3	FUC	B	5	3	10,10,11	1.47	2 (20%)	14,14,16	1.53	4 (28%)
3	GAL	C	1	3	12,12,12	1.11	1 (8%)	17,17,17	1.85	4 (23%)
3	GLA	C	2	3	11,11,12	1.17	1 (9%)	15,15,17	1.92	4 (26%)
3	NGA	C	3	3	14,14,15	1.10	1 (7%)	17,19,21	2.18	4 (23%)
3	GAL	C	4	3	11,11,12	1.23	1 (9%)	15,15,17	1.46	1 (6%)
3	FUC	C	5	3	10,10,11	1.39	1 (10%)	14,14,16	1.31	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLA	A	1	2	-	0/2/22/22	0/1/1/1
2	NGA	A	2	2	-	1/6/23/26	0/1/1/1
2	GAL	A	3	2	-	2/2/19/22	0/1/1/1
2	FUC	A	4	2	-	-	0/1/1/1
3	GAL	B	1	3	-	1/2/22/22	0/1/1/1
3	GLA	B	2	3	-	0/2/19/22	0/1/1/1
3	NGA	B	3	3	-	2/6/23/26	0/1/1/1
3	GAL	B	4	3	-	1/2/19/22	0/1/1/1
3	FUC	B	5	3	-	-	0/1/1/1
3	GAL	C	1	3	-	1/2/22/22	0/1/1/1
3	GLA	C	2	3	-	2/2/19/22	0/1/1/1
3	NGA	C	3	3	-	0/6/23/26	0/1/1/1
3	GAL	C	4	3	-	1/2/19/22	0/1/1/1
3	FUC	C	5	3	-	-	0/1/1/1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	5	FUC	O5-C5	-3.82	1.36	1.43
2	A	4	FUC	C2-C3	-3.72	1.46	1.52
3	C	2	GLA	C2-C3	3.11	1.57	1.52
3	B	5	FUC	O5-C5	-3.10	1.37	1.43
3	B	2	GLA	C2-C3	2.89	1.56	1.52

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	GLA	O4-C4-C3	-5.35	97.78	110.38
3	B	2	GLA	O2-C2-C3	5.25	121.03	110.15
3	C	2	GLA	O3-C3-C4	-5.12	98.31	110.38
3	B	2	GLA	O4-C4-C5	4.90	121.38	109.32
3	C	3	NGA	C1-O5-C5	-4.66	105.94	112.19

There are no chirality outliers.


5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	3	NGA	O5-C5-C6-O6
3	C	2	GLA	C4-C5-C6-O6
3	C	2	GLA	O5-C5-C6-O6
3	B	3	NGA	C4-C5-C6-O6
3	B	4	GAL	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	134/134 (100%)	0.11	4 (2%) 52 56	11, 22, 49, 63	2 (1%)
1	BBB	134/134 (100%)	0.15	1 (0%) 84 86	12, 21, 47, 64	6 (4%)
1	CCC	134/134 (100%)	0.16	3 (2%) 62 66	11, 22, 50, 63	4 (2%)
All	All	402/402 (100%)	0.14	8 (1%) 65 69	11, 22, 50, 64	12 (2%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	131	ALA	3.0
1	CCC	131	ALA	2.9
1	BBB	98	SER	2.9
1	AAA	31	ALA	2.6
1	AAA	98	SER	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLA	A	1	12/12	-	-	36,46,50,50	0
2	NGA	A	2	14/15	-	-	25,30,46,50	0
2	GAL	A	3	11/12	-	-	21,29,41,47	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FUC	A	4	10/11	-	-	17,19,21,22	0
3	GAL	B	1	12/12	-	-	50,53,65,66	0
3	GLA	B	2	11/12	-	-	49,59,66,67	0
3	NGA	B	3	14/15	-	-	29,34,43,54	0
3	GAL	B	4	11/12	-	-	23,31,43,44	0
3	FUC	B	5	10/11	-	-	17,20,22,22	0
3	GAL	C	1	12/12	-	-	41,48,53,57	0
3	GLA	C	2	11/12	-	-	39,49,58,61	0
3	NGA	C	3	14/15	-	-	28,32,47,52	0
3	GAL	C	4	11/12	-	-	24,32,44,45	0
3	FUC	C	5	10/11	-	-	17,19,22,22	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.