



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:43 AM UTC

PDB ID : 1TJ1 / pdb_00001tj1
Title : Crystal structure of E. coli PutA proline dehydrogenase domain (residues 86-669) complexed with L-lactate
Authors : Tanner, J.J.; Zhang, M.; White, T.A.; Schuermann, J.P.; Baban, B.A.; Becker, D.F.
Deposited on : 2004-06-03
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	602	12% 67% 7% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FAD	A	2001	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional putA protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	450	3474	2206	611	640	17	0	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	670	SER	-	expression tag	UNP P09546
A	671	SER	-	expression tag	UNP P09546
A	672	SER	-	expression tag	UNP P09546
A	673	VAL	-	expression tag	UNP P09546
A	674	ASP	-	expression tag	UNP P09546
A	675	LYS	-	expression tag	UNP P09546
A	676	LEU	-	expression tag	UNP P09546
A	677	ALA	-	expression tag	UNP P09546
A	678	ALA	-	expression tag	UNP P09546
A	679	ALA	-	expression tag	UNP P09546
A	680	LEU	-	expression tag	UNP P09546
A	681	GLU	-	expression tag	UNP P09546
A	682	HIS	-	expression tag	UNP P09546
A	683	HIS	-	expression tag	UNP P09546
A	684	HIS	-	expression tag	UNP P09546
A	685	HIS	-	expression tag	UNP P09546
A	686	HIS	-	expression tag	UNP P09546
A	687	HIS	-	expression tag	UNP P09546

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is (2S)-2-HYDROXYPROPANOIC ACID (CCD ID: 20P) (formula: $C_3H_6O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	194	Total 194	O 194	0	0

- Molecule 1: Bifunctional putA protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	72.81Å 141.12Å 146.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.49 – 2.00 40.49 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.0 (40.49-2.00) 97.8 (40.49-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.214 , 0.255 0.221 , 0.263	Depositor DCC
R_{free} test set	2492 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.798	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3727	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2OP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/3538	0.92	1/4793 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	ARG	N-CA-C	5.71	117.50	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3474	0	3467	24	0
2	A	53	0	30	3	0
3	A	6	0	5	0	0
4	A	194	0	0	2	0
All	All	3727	0	3502	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ARG:NH2	1:A:382:ASP:OD2	2.23	0.68
1:A:503:TYR:OH	1:A:505:PRO:HA	1.98	0.63
1:A:409:ARG:HD2	4:A:2004:HOH:O	2.00	0.61
1:A:409:ARG:CD	4:A:2004:HOH:O	2.50	0.60
1:A:101:ARG:NH1	1:A:109:MET:HE1	2.24	0.52
1:A:111:LEU:HD22	1:A:585:VAL:HG13	1.93	0.51
1:A:261:THR:HG21	1:A:541:ALA:HB1	1.96	0.47
1:A:410:CYS:HB3	1:A:411:PRO:HD3	1.98	0.46
1:A:396:TRP:CH2	1:A:398:GLY:HA3	2.51	0.45
1:A:261:THR:HG22	1:A:514:HIS:CG	2.50	0.45
1:A:400:GLY:HA2	1:A:429:MET:O	2.17	0.44
2:A:2001:FAD:H9	2:A:2001:FAD:H1'1	1.73	0.44
1:A:570:ILE:HD12	1:A:580:LEU:CD2	2.47	0.44
1:A:570:ILE:HD12	1:A:580:LEU:HD21	1.99	0.44
1:A:518:GLU:HB2	1:A:519:PRO:HD3	2.00	0.43
1:A:98:ALA:O	1:A:99:TYR:C	2.62	0.43
1:A:396:TRP:CZ2	1:A:398:GLY:HA3	2.54	0.42
1:A:110:LEU:HD13	1:A:465:SER:HA	2.02	0.42
1:A:396:TRP:O	1:A:426:ARG:NH1	2.53	0.42
1:A:540:TYR:HB2	2:A:2001:FAD:HM72	2.01	0.41
1:A:503:TYR:CZ	1:A:505:PRO:HA	2.55	0.41
1:A:377:LEU:HD23	1:A:409:ARG:CZ	2.51	0.41
2:A:2001:FAD:H1'1	2:A:2001:FAD:H4'	1.87	0.41
1:A:135:LEU:HD11	1:A:519:PRO:HB2	2.03	0.40
1:A:137:ASN:ND2	1:A:142:SER:CB	2.84	0.40
1:A:431:ARG:HA	1:A:483:GLN:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/602 (74%)	435 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	347/490 (71%)	330 (95%)	17 (5%)	22	20

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	LEU
1	A	163	VAL
1	A	168	LEU
1	A	254	LEU
1	A	263	GLU
1	A	281	ARG
1	A	344	ARG
1	A	384	LEU
1	A	406	TYR
1	A	409	ARG
1	A	426	ARG
1	A	443	LYS
1	A	495	ILE
1	A	501	GLN
1	A	547	GLU
1	A	563	ASN
1	A	573	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	137	ASN
1	A	271	ASN
1	A	298	GLN
1	A	305	GLN
1	A	316	ASN
1	A	497	GLN
1	A	501	GLN
1	A	534	ASN
1	A	597	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	2001	-	58,58,58	1.68	9 (15%)	85,89,89	1.68	19 (22%)
3	2OP	A	2002	-	4,5,5	1.01	0	2,6,6	1.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	2001	-	1/1/9/9	5/34/50/50	0/6/6/6
3	2OP	A	2002	-	-	1/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	P-O3P	6.08	1.66	1.59
2	A	2001	FAD	C4X-N5	4.50	1.40	1.30
2	A	2001	FAD	O4B-C4B	-4.04	1.36	1.45
2	A	2001	FAD	C5'-C4'	3.59	1.56	1.51
2	A	2001	FAD	C2A-N1A	3.50	1.40	1.33
2	A	2001	FAD	C5A-N7A	-2.86	1.33	1.39
2	A	2001	FAD	C10-N1	2.71	1.38	1.33
2	A	2001	FAD	O4B-C1B	2.60	1.48	1.42
2	A	2001	FAD	C4A-N9A	-2.45	1.32	1.37

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	O4B-C4B-C5B	5.07	125.58	109.33
2	A	2001	FAD	N3A-C2A-N1A	-4.29	122.09	128.58
2	A	2001	FAD	O3P-PA-O1A	-4.11	98.33	110.70
2	A	2001	FAD	N9A-C8A-N7A	-3.89	108.42	113.94
2	A	2001	FAD	C4X-C10-N10	3.34	121.26	116.48
2	A	2001	FAD	O2A-PA-O3P	3.26	116.09	107.27
2	A	2001	FAD	C4A-N9A-C8A	3.02	108.91	105.74
2	A	2001	FAD	O4B-C4B-C3B	3.01	111.12	105.15
2	A	2001	FAD	C10-C4X-N5	-2.76	119.17	124.81
2	A	2001	FAD	C5A-C4A-N3A	-2.66	123.05	126.72
2	A	2001	FAD	C5A-N7A-C8A	2.58	107.51	103.45
2	A	2001	FAD	C5X-C9A-N10	2.40	120.14	117.97
2	A	2001	FAD	C2A-N3A-C4A	2.35	117.56	111.83
2	A	2001	FAD	C9-C9A-N10	-2.34	118.71	121.85
2	A	2001	FAD	C4-N3-C2	-2.18	121.77	125.64
2	A	2001	FAD	C5B-C4B-C3B	2.11	122.79	115.21
2	A	2001	FAD	O5B-C5B-C4B	2.08	116.08	108.99
2	A	2001	FAD	O4B-C1B-N9A	-2.07	104.12	108.09
2	A	2001	FAD	C4-C4X-N5	2.05	121.04	118.21

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	2001	FAD	C4B

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2001	FAD	N10-C1'-C2'-O2'
2	A	2001	FAD	N10-C1'-C2'-C3'
2	A	2001	FAD	C1'-C2'-C3'-C4'
2	A	2001	FAD	O4B-C4B-C5B-O5B
2	A	2001	FAD	C3B-C4B-C5B-O5B
3	A	2002	2OP	O-C-CA-OHN

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/602 (74%)	0.92	75 (16%) 4 4	17, 29, 56, 76	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	147	MET	7.8
1	A	145	ALA	6.1
1	A	245	ARG	5.2
1	A	163	VAL	5.0
1	A	183	ALA	4.5
1	A	143	GLY	4.5
1	A	165	LEU	4.3
1	A	138	GLN	4.2
1	A	169	ALA	4.2
1	A	574	SER	4.1
1	A	244	ILE	4.0
1	A	164	ALA	3.9
1	A	167	CYS	3.8
1	A	184	LEU	3.8
1	A	246	LYS	3.7
1	A	142	SER	3.5
1	A	546	HIS	3.4
1	A	529	ALA	3.4
1	A	296	ASP	3.3
1	A	146	GLY	3.3
1	A	141	ALA	3.3
1	A	573	THR	3.2
1	A	168	LEU	3.2
1	A	144	ARG	3.1
1	A	251	ALA	3.1
1	A	575	LEU	3.1
1	A	140	ASN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	162	GLY	3.0
1	A	294	ALA	3.0
1	A	166	MET	3.0
1	A	253	ARG	3.0
1	A	182	ASP	2.9
1	A	118	GLN	2.8
1	A	172	LEU	2.8
1	A	250	MET	2.7
1	A	93	ALA	2.7
1	A	173	LEU	2.7
1	A	443	LYS	2.7
1	A	545	THR	2.6
1	A	249	ASP	2.6
1	A	576	PRO	2.5
1	A	177	ASP	2.5
1	A	571	ALA	2.5
1	A	247	GLY	2.5
1	A	547	GLU	2.5
1	A	298	GLN	2.4
1	A	610	ASP	2.4
1	A	272	ALA	2.4
1	A	137	ASN	2.4
1	A	134	LYS	2.4
1	A	171	ALA	2.4
1	A	572	ASP	2.4
1	A	444	ARG	2.4
1	A	97	ALA	2.3
1	A	262	GLY	2.3
1	A	248	VAL	2.3
1	A	293	THR	2.3
1	A	109	MET	2.3
1	A	263	GLU	2.3
1	A	179	ALA	2.3
1	A	270	ALA	2.3
1	A	175	ILE	2.2
1	A	448	ASP	2.2
1	A	383	LEU	2.2
1	A	314	ALA	2.1
1	A	592	ALA	2.1
1	A	447	MET	2.1
1	A	257	GLU	2.1
1	A	563	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	306	GLN	2.1
1	A	451	GLU	2.0
1	A	254	LEU	2.0
1	A	271	ASN	2.0
1	A	607	LEU	2.0
1	A	256	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	2OP	A	2002	6/6	0.95	0.10	27,28,29,29	0
2	FAD	A	2001	53/53	0.97	0.07	23,27,31,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.