



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 11:16 AM UTC

PDB ID : 7TL7 / pdb_00007tl7
Title : 1.90Å resolution structure of independent phosphoglycerate mutase from *C. elegans* in complex with a macrocyclic peptide inhibitor (Sa-D2)
Authors : Liu, L.; Lovell, S.; Battaile, K.P.; Dranchak, P.; Queme, B.; Aitha, M.; van Neer, R.H.P.; Kimura, H.; Katho, T.; Suga, H.; Inglese, J.
Deposited on : 2022-01-18
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	93% 6%
1	B	534	2% 90% 8%
1	C	534	% 91% 7%
1	D	534	2% 91% 7%
2	a	15	73% 20% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	b	15	73% 20% 7%
2	c	15	73% 13% 13%
2	d	15	87% 7% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 17471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-independent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	531	Total	C	N	O	S	0	1	0
			4039	2551	700	759	29			
1	B	526	Total	C	N	O	S	0	2	0
			3994	2525	688	752	29			
1	C	522	Total	C	N	O	S	0	2	0
			3977	2513	684	750	30			
1	D	521	Total	C	N	O	S	0	1	0
			3917	2482	675	731	29			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	540	LYS	-	expression tag	UNP G5EFZ1
A	541	LEU	-	expression tag	UNP G5EFZ1
A	542	ALA	-	expression tag	UNP G5EFZ1
A	543	ALA	-	expression tag	UNP G5EFZ1
A	544	ALA	-	expression tag	UNP G5EFZ1
A	545	LEU	-	expression tag	UNP G5EFZ1
A	546	GLU	-	expression tag	UNP G5EFZ1
A	547	HIS	-	expression tag	UNP G5EFZ1
A	548	HIS	-	expression tag	UNP G5EFZ1
A	549	HIS	-	expression tag	UNP G5EFZ1
A	550	HIS	-	expression tag	UNP G5EFZ1
A	551	HIS	-	expression tag	UNP G5EFZ1
A	552	HIS	-	expression tag	UNP G5EFZ1
B	540	LYS	-	expression tag	UNP G5EFZ1
B	541	LEU	-	expression tag	UNP G5EFZ1
B	542	ALA	-	expression tag	UNP G5EFZ1
B	543	ALA	-	expression tag	UNP G5EFZ1
B	544	ALA	-	expression tag	UNP G5EFZ1
B	545	LEU	-	expression tag	UNP G5EFZ1
B	546	GLU	-	expression tag	UNP G5EFZ1
B	547	HIS	-	expression tag	UNP G5EFZ1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	548	HIS	-	expression tag	UNP G5EFZ1
B	549	HIS	-	expression tag	UNP G5EFZ1
B	550	HIS	-	expression tag	UNP G5EFZ1
B	551	HIS	-	expression tag	UNP G5EFZ1
B	552	HIS	-	expression tag	UNP G5EFZ1
C	540	LYS	-	expression tag	UNP G5EFZ1
C	541	LEU	-	expression tag	UNP G5EFZ1
C	542	ALA	-	expression tag	UNP G5EFZ1
C	543	ALA	-	expression tag	UNP G5EFZ1
C	544	ALA	-	expression tag	UNP G5EFZ1
C	545	LEU	-	expression tag	UNP G5EFZ1
C	546	GLU	-	expression tag	UNP G5EFZ1
C	547	HIS	-	expression tag	UNP G5EFZ1
C	548	HIS	-	expression tag	UNP G5EFZ1
C	549	HIS	-	expression tag	UNP G5EFZ1
C	550	HIS	-	expression tag	UNP G5EFZ1
C	551	HIS	-	expression tag	UNP G5EFZ1
C	552	HIS	-	expression tag	UNP G5EFZ1
D	540	LYS	-	expression tag	UNP G5EFZ1
D	541	LEU	-	expression tag	UNP G5EFZ1
D	542	ALA	-	expression tag	UNP G5EFZ1
D	543	ALA	-	expression tag	UNP G5EFZ1
D	544	ALA	-	expression tag	UNP G5EFZ1
D	545	LEU	-	expression tag	UNP G5EFZ1
D	546	GLU	-	expression tag	UNP G5EFZ1
D	547	HIS	-	expression tag	UNP G5EFZ1
D	548	HIS	-	expression tag	UNP G5EFZ1
D	549	HIS	-	expression tag	UNP G5EFZ1
D	550	HIS	-	expression tag	UNP G5EFZ1
D	551	HIS	-	expression tag	UNP G5EFZ1
D	552	HIS	-	expression tag	UNP G5EFZ1

- Molecule 2 is a protein called peptide Sa-D2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	a	15	Total	C	N	O	S	0	0	1
			127	86	16	23	2			
2	b	15	Total	C	N	O	S	0	0	1
			127	86	16	23	2			
2	c	15	Total	C	N	O	S	0	0	1
			123	83	15	23	2			
2	d	15	Total	C	N	O	S	0	0	1
			124	84	15	23	2			


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Zn 2 2	0	0
3	B	2	Total Zn 2 2	0	0
3	C	2	Total Zn 2 2	0	0
3	D	2	Total Zn 2 2	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0
4	B	1	Total Na 1 1	0	0
4	C	1	Total Na 1 1	0	0
4	D	1	Total Na 1 1	0	0

- Molecule 5 is IMIDAZOLE (CCD ID: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	N	0	0
			5	3	2		
5	D	1	Total	C	N	0	0
			5	3	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	273	Total	O	0	0
			273	273		
6	B	253	Total	O	0	0
			253	253		
6	C	269	Total	O	0	0
			269	269		
6	D	174	Total	O	0	0
			174	174		
6	a	13	Total	O	0	0
			13	13		
6	b	14	Total	O	0	0
			14	14		
6	c	14	Total	O	0	0
			14	14		
6	d	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase



- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase




- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase




- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase







● Molecule 2: peptide Sa-D2





● Molecule 2: peptide Sa-D2



● Molecule 2: peptide Sa-D2



● Molecule 2: peptide Sa-D2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.00Å 87.62Å 151.34Å 90.00° 97.13° 90.00°	Depositor
Resolution (Å)	39.84 – 1.90 39.84 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.84-1.90) 99.7 (39.84-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.20rc2_4402	Depositor
R, R_{free}	0.171 , 0.216 0.178 , 0.221	Depositor DCC
R_{free} test set	7975 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17471	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9809e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, ACE, NH2, DTY, NA, ZN, YNM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.18	0/4133	0.42	0/5608
1	B	0.17	0/4086	0.42	0/5546
1	C	0.18	0/4072	0.44	0/5524
1	D	0.16	0/4009	0.40	0/5445
2	a	0.19	0/85	0.43	0/109
2	b	0.20	0/85	0.40	0/109
2	c	0.19	0/81	0.50	0/105
2	d	0.23	0/82	0.44	0/106
All	All	0.17	0/16633	0.42	0/22552

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4039	0	3893	18	0
1	B	3994	0	3838	25	0
1	C	3977	0	3833	21	0
1	D	3917	0	3747	19	0
2	a	127	0	107	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	b	127	0	106	2	0
2	c	123	0	95	2	1
2	d	124	0	97	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	C	5	0	5	3	0
5	D	5	0	5	1	0
6	A	273	0	0	0	0
6	B	253	0	0	1	0
6	C	269	0	0	1	0
6	D	174	0	0	2	0
6	a	13	0	0	0	0
6	b	14	0	0	1	0
6	c	14	0	0	0	0
6	d	11	0	0	0	0
All	All	17471	0	15726	89	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:214:MET:HB3	1:D:287:ARG:HB3	1.72	0.72
1:C:21:MET:HE3	1:C:348:LYS:HA	1.80	0.63
1:C:141:VAL:HG22	1:C:153[A]:MET:HE2	1.83	0.61
1:D:50:ILE:HD12	1:D:472:GLU:HB3	1.85	0.58
1:A:265:LEU:O	1:A:271:ARG:NH2	2.36	0.58

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:a:2:DTY:OH	2:c:5:GLU:OE1[2_655]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/534 (99%)	520 (98%)	8 (2%)	2 (0%)	30	22
1	B	526/534 (98%)	516 (98%)	9 (2%)	1 (0%)	43	36
1	C	522/534 (98%)	513 (98%)	8 (2%)	1 (0%)	43	36
1	D	520/534 (97%)	507 (98%)	11 (2%)	2 (0%)	30	22
2	a	10/15 (67%)	9 (90%)	1 (10%)	0	100	100
2	b	10/15 (67%)	9 (90%)	1 (10%)	0	100	100
2	c	10/15 (67%)	8 (80%)	2 (20%)	0	100	100
2	d	10/15 (67%)	8 (80%)	2 (20%)	0	100	100
All	All	2138/2196 (97%)	2090 (98%)	42 (2%)	6 (0%)	36	29

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	389	ALA
1	A	468	HIS
1	B	470	ASN
1	C	470	ASN
1	D	470	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/432 (96%)	414 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	406/432 (94%)	404 (100%)	2 (0%)	81	84
1	C	409/432 (95%)	407 (100%)	2 (0%)	81	84
1	D	393/432 (91%)	393 (100%)	0	100	100
2	a	9/10 (90%)	9 (100%)	0	100	100
2	b	9/10 (90%)	9 (100%)	0	100	100
2	c	8/10 (80%)	8 (100%)	0	100	100
2	d	8/10 (80%)	8 (100%)	0	100	100
All	All	1656/1768 (94%)	1652 (100%)	4 (0%)	87	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	116	VAL
1	B	254	ASP
1	C	153[A]	MET
1	C	153[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	456	GLN
1	D	320	GLN
1	B	508	HIS
1	D	409	GLN
1	D	112	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DTY	d	2	2	11,12,13	0.38	0	10,15,17	0.19	0
2	DTY	b	2	2	11,12,13	0.37	0	10,15,17	0.19	0
2	YNM	c	9	2	12,13,14	0.59	0	15,16,18	1.26	3 (20%)
2	DTY	a	2	2	11,12,13	0.41	0	10,15,17	0.22	0
2	YNM	d	9	2	12,13,14	0.61	0	15,16,18	1.20	3 (20%)
2	YNM	b	9	2	12,13,14	0.63	0	15,16,18	1.08	2 (13%)
2	YNM	b	6	2	12,13,14	0.57	0	15,16,18	1.17	2 (13%)
2	YNM	d	6	2	12,13,14	0.56	0	15,16,18	1.10	0
2	YNM	c	6	2	12,13,14	0.61	0	15,16,18	1.12	2 (13%)
2	YNM	a	9	2	12,13,14	0.64	0	15,16,18	1.39	3 (20%)
2	DTY	c	2	2	11,12,13	0.40	0	10,15,17	0.18	0
2	YNM	a	6	2	12,13,14	0.61	0	15,16,18	1.17	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DTY	d	2	2	-	0/5/6/8	0/1/1/1
2	DTY	b	2	2	-	0/5/6/8	0/1/1/1
2	YNM	c	9	2	-	0/5/8/10	0/1/1/1
2	DTY	a	2	2	-	0/5/6/8	0/1/1/1
2	YNM	d	9	2	-	0/5/8/10	0/1/1/1
2	YNM	b	9	2	-	0/5/8/10	0/1/1/1
2	YNM	b	6	2	-	0/5/8/10	0/1/1/1
2	YNM	d	6	2	-	0/5/8/10	0/1/1/1
2	YNM	c	6	2	-	0/5/8/10	0/1/1/1
2	YNM	a	9	2	-	0/5/8/10	0/1/1/1
2	DTY	c	2	2	-	0/5/6/8	0/1/1/1
2	YNM	a	6	2	-	0/5/8/10	0/1/1/1

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	c	9	YNM	CG-CB-CA	-3.19	108.97	113.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	a	9	YNM	CG-CB-CA	-2.91	109.37	113.51
2	a	9	YNM	CM-N-CA	2.47	121.09	113.70
2	d	9	YNM	CM-N-CA	2.47	121.08	113.70
2	d	9	YNM	CB-CA-C	-2.33	107.17	111.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	c	9	YNM	1	0
2	a	2	DTY	0	1
2	d	9	YNM	1	0
2	b	6	YNM	1	0
2	c	6	YNM	1	0
2	a	9	YNM	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 12 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	IMD	D	601	-	5,5,5	0.64	0	5,5,5	0.45	0
5	IMD	C	601	-	5,5,5	0.64	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	IMD	D	601	-	-	-	0/1/1/1
5	IMD	C	601	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	601	IMD	1	0
5	C	601	IMD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/534 (99%)	-0.12	2 (0%) 88 90	14, 26, 40, 66	1 (0%)
1	B	526/534 (98%)	0.10	10 (1%) 66 70	13, 28, 48, 83	2 (0%)
1	C	522/534 (97%)	-0.15	3 (0%) 85 87	14, 26, 41, 54	2 (0%)
1	D	521/534 (97%)	0.21	10 (1%) 66 70	17, 32, 49, 76	1 (0%)
2	a	10/15 (66%)	-0.23	0 100 100	21, 24, 38, 41	0
2	b	10/15 (66%)	-0.30	0 100 100	21, 26, 35, 36	0
2	c	10/15 (66%)	-0.05	0 100 100	25, 31, 40, 40	0
2	d	10/15 (66%)	0.26	0 100 100	28, 32, 39, 40	0
All	All	2140/2196 (97%)	0.01	25 (1%) 76 79	13, 28, 45, 83	6 (0%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	389	ALA	4.1
1	B	393	LEU	3.6
1	B	389	ALA	3.5
1	B	543	ALA	3.5
1	B	436	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DTY	c	2	12/13	0.89	0.15	31,59,73,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	DTY	d	2	12/13	0.91	0.10	28,38,45,54	0
2	YNM	c	9	13/14	0.91	0.10	26,32,39,41	0
2	DTY	a	2	12/13	0.92	0.11	24,37,50,57	0
2	YNM	c	6	13/14	0.93	0.09	25,33,43,45	0
2	YNM	d	9	13/14	0.93	0.08	28,33,38,40	0
2	YNM	d	6	13/14	0.94	0.08	23,31,39,41	0
2	YNM	a	9	13/14	0.94	0.08	21,24,33,40	0
2	YNM	b	6	13/14	0.95	0.07	22,26,34,36	0
2	YNM	b	9	13/14	0.95	0.07	20,26,31,36	0
2	YNM	a	6	13/14	0.96	0.07	21,23,32,32	0
2	DTY	b	2	12/13	0.96	0.07	19,30,39,46	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NA	B	603	1/1	0.86	0.12	44,44,44,44	0
4	NA	D	604	1/1	0.94	0.06	48,48,48,48	0
5	IMD	D	601	5/5	0.94	0.07	19,20,22,24	0
5	IMD	C	601	5/5	0.95	0.06	18,18,22,23	0
4	NA	A	603	1/1	0.98	0.06	28,28,28,28	0
3	ZN	B	601	1/1	0.98	0.04	28,28,28,28	0
4	NA	C	604	1/1	0.98	0.05	27,27,27,27	0
3	ZN	B	602	1/1	0.99	0.05	34,34,34,34	0
3	ZN	D	602	1/1	0.99	0.02	27,27,27,27	0
3	ZN	C	602	1/1	1.00	0.01	19,19,19,19	0
3	ZN	C	603	1/1	1.00	0.04	25,25,25,25	0
3	ZN	A	601	1/1	1.00	0.01	23,23,23,23	0
3	ZN	D	603	1/1	1.00	0.05	32,32,32,32	0
3	ZN	A	602	1/1	1.00	0.06	28,28,28,28	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ZN B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.