



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:27 PM UTC

PDB ID : 8TLN / pdb_00008tlh
Title : STRUCTURAL COMPARISON SUGGESTS THAT THERMOLYSIN AND
RELATED NEUTRAL PROTEASES UNDERGO HINGE-BENDING MO-
TION DURING CATALYSIS
Authors : Tronrud, D.; Matthews, B.W.
Deposited on : 1993-09-01
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	316	2% 72% 22% 5%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THERMOLYSIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	316	2438	1530	411	495	2	0	2	0

- Molecule 2 is VALINE (CCD ID: VAL) (formula: $C_5H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	E	1	7	5	1	1	0	0

- Molecule 3 is LYSINE (CCD ID: LYS) (formula: $C_6H_{15}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	E	1	Total	C	N	O	0	0
			10	6	2	2		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	4	Total	Ca	0	0
			4	4		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	1	Total	Zn	0	0
			1	1		

- Molecule 6 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	E	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	157	Total	O	0	0
			157	157		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THERMOLYSIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	94.10Å 94.10Å 131.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.60 20.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.60) 76.2 (20.00-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 1.60Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.165 , (Not available) 0.177 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	12.4	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 70.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2621	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DMS, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	1.79	36/2508 (1.4%)	1.99	63/3413 (1.8%)

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	142	HIS	CE1-NE2	9.59	1.42	1.32
1	E	131	ILE	CA-CB	-7.32	1.44	1.54
1	E	258	ILE	CA-CB	-7.02	1.47	1.54
1	E	226	ASP	CA-C	-6.55	1.44	1.53
1	E	299	THR	CA-C	-6.43	1.44	1.52
1	E	276	THR	C-O	-6.31	1.18	1.24
1	E	52	GLY	CA-C	6.17	1.58	1.52
1	E	194	THR	C-N	-6.12	1.26	1.33
1	E	213	ASP	C-N	-6.00	1.25	1.33
1	E	224	THR	C-N	-5.99	1.25	1.33
1	E	226	ASP	C-N	-5.92	1.25	1.33
1	E	49	THR	CB-OG1	-5.89	1.34	1.43
1	E	197	ILE	CA-CB	-5.88	1.47	1.54
1	E	272	THR	CA-C	-5.86	1.44	1.52
1	E	9	VAL	C-O	5.84	1.29	1.23
1	E	302	GLU	C-N	-5.77	1.26	1.33
1	E	196	GLY	CA-C	5.76	1.59	1.51
1	E	242	TYR	CA-C	5.71	1.60	1.52
1	E	93	TYR	CA-C	-5.65	1.44	1.52
1	E	303	VAL	N-CA	-5.60	1.39	1.46
1	E	68	ALA	N-CA	5.52	1.51	1.46
1	E	74	HIS	ND1-CE1	5.49	1.38	1.32
1	E	23	THR	CA-C	-5.43	1.45	1.52
1	E	32	ASP	C-N	5.32	1.41	1.33
1	E	194	THR	CA-C	-5.30	1.46	1.52
1	E	83	TYR	CA-CB	5.26	1.61	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	211	TYR	CA-C	-5.24	1.45	1.52
1	E	307	LYS	N-CA	5.21	1.52	1.46
1	E	8	GLY	N-CA	5.18	1.49	1.44
1	E	175	LEU	N-CA	5.17	1.52	1.46
1	E	258	ILE	N-CA	5.13	1.51	1.46
1	E	56	ALA	CA-C	-5.08	1.46	1.52
1	E	150	ASP	CG-OD2	5.06	1.34	1.25
1	E	260	ARG	CA-C	-5.06	1.46	1.52
1	E	170	ASP	CG-OD1	5.06	1.34	1.25
1	E	20	ILE	C-O	5.03	1.29	1.23

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	68	ALA	CA-C-O	8.86	126.78	118.63
1	E	183	ASN	CB-CA-C	-7.72	94.97	110.17
1	E	172	PHE	CA-C-N	-7.16	111.13	120.22
1	E	172	PHE	C-N-CA	-7.16	111.13	120.22
1	E	52	GLY	CA-C-O	-7.13	115.19	122.25
1	E	215	ASP	CA-CB-CG	-6.99	105.61	112.60
1	E	231	HIS	N-CA-CB	6.87	121.25	110.46
1	E	21	ASN	CA-CB-CG	-6.86	105.74	112.60
1	E	112	ASN	OD1-CG-ND2	-6.83	115.77	122.60
1	E	197	ILE	N-CA-CB	-6.81	101.98	111.25
1	E	172	PHE	CA-CB-CG	-6.56	107.24	113.80
1	E	316	LYS	CB-CG-CD	6.55	126.36	111.30
1	E	151	TYR	CA-CB-CG	-6.54	102.14	113.90
1	E	308	GLN	CB-CA-C	-6.45	100.08	110.79
1	E	181	ASN	N-CA-CB	-6.40	102.47	112.13
1	E	128	GLN	CB-CA-C	-6.35	99.95	110.37
1	E	260	ARG	N-CA-C	6.23	118.15	111.36
1	E	256	VAL	CB-CA-C	-6.12	102.82	111.21
1	E	213	ASP	CA-CB-CG	-6.05	106.55	112.60
1	E	225	GLN	CA-CB-CG	6.02	126.14	114.10
1	E	298	SER	N-CA-C	6.01	119.72	112.38
1	E	148	VAL	O-C-N	-5.94	116.09	121.91
1	E	251	TYR	N-CA-CB	5.92	120.67	111.84
1	E	89	ASN	CA-CB-CG	-5.92	106.68	112.60
1	E	117	GLY	N-CA-C	-5.88	107.69	114.69
1	E	309	ALA	O-C-N	-5.85	115.91	122.12
1	E	86	ASN	CA-CB-CG	-5.83	106.77	112.60
1	E	58	ALA	N-CA-C	5.79	119.53	112.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	21	ASN	OD1-CG-ND2	-5.77	116.83	122.60
1	E	120	MET	CA-CB-CG	-5.74	102.62	114.10
1	E	234	SER	N-CA-C	-5.70	105.59	112.54
1	E	260	ARG	CB-CA-C	-5.64	101.26	110.85
1	E	169	SER	O-C-N	-5.63	116.16	122.12
1	E	47	ARG	CB-CG-CD	5.58	124.13	111.30
1	E	128	GLN	CA-C-O	5.53	126.27	119.81
1	E	224	THR	CA-C-N	-5.46	111.22	120.95
1	E	224	THR	C-N-CA	-5.46	111.22	120.95
1	E	1	ILE	O-C-N	-5.45	114.28	123.00
1	E	89	ASN	OD1-CG-ND2	5.43	128.03	122.60
1	E	225	GLN	N-CA-CB	-5.42	102.92	110.29
1	E	200	ASP	CB-CG-OD2	-5.40	105.99	118.40
1	E	194	THR	CA-C-N	5.39	125.22	119.28
1	E	194	THR	C-N-CA	5.39	125.22	119.28
1	E	150	ASP	CB-CA-C	-5.38	100.67	110.63
1	E	194	THR	O-C-N	5.37	127.49	121.32
1	E	97	ASN	N-CA-C	5.36	119.10	112.24
1	E	32	ASP	CA-CB-CG	5.29	117.89	112.60
1	E	49	THR	OG1-CB-CG2	-5.28	98.74	109.30
1	E	182	LYS	N-CA-CB	-5.25	103.49	110.57
1	E	159	ASN	CB-CA-C	-5.24	109.55	115.79
1	E	190	GLU	N-CA-C	5.24	117.67	111.33
1	E	133	LEU	N-CA-C	5.23	117.73	111.71
1	E	89	ASN	N-CA-CB	-5.23	104.35	112.30
1	E	66	TYR	N-CA-CB	-5.22	101.73	110.39
1	E	195	PRO	N-CA-C	-5.21	106.62	113.65
1	E	299	THR	N-CA-C	-5.21	106.24	112.59
1	E	224	THR	N-CA-C	5.17	120.58	113.97
1	E	124	ASP	CA-C-N	-5.13	115.49	121.85
1	E	124	ASP	C-N-CA	-5.13	115.49	121.85
1	E	249	THR	CA-C-N	-5.07	114.89	122.39
1	E	249	THR	C-N-CA	-5.07	114.89	122.39
1	E	316	LYS	CB-CA-C	5.06	119.72	110.10
1	E	88	HIS	O-C-N	5.06	128.17	122.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2438	0	2273	30	0
2	E	7	0	8	0	0
3	E	10	0	13	1	0
4	E	4	0	0	0	0
5	E	1	0	0	0	0
6	E	4	0	6	0	0
7	E	157	0	0	0	0
All	All	2621	0	2300	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:1:ILE:CD1	1:E:29:TYR:CE2	2.58	0.86
1:E:1:ILE:CD1	1:E:29:TYR:CD2	2.67	0.77
1:E:311:ASP:OD1	1:E:316:LYS:HE3	1.84	0.76
1:E:1:ILE:HD11	1:E:29:TYR:CE2	2.24	0.72
1:E:1:ILE:HD11	1:E:29:TYR:CZ	2.26	0.70
1:E:1:ILE:HD12	1:E:29:TYR:CD2	2.32	0.64
1:E:1:ILE:HD12	1:E:29:TYR:CG	2.41	0.55
1:E:31:GLN:HG3	1:E:40:PHE:CE2	2.45	0.52
1:E:1:ILE:HD13	1:E:29:TYR:CE2	2.42	0.51
1:E:128:GLN:O	1:E:195:PRO:HD2	2.12	0.48
1:E:1:ILE:HD13	1:E:29:TYR:CD2	2.47	0.47
1:E:194:THR:HA	1:E:195:PRO:HD2	1.39	0.46
1:E:88:HIS:O	1:E:89:ASN:CB	2.64	0.45
1:E:11[A]:ARG:NH1	1:E:11[A]:ARG:HG3	2.31	0.45
1:E:131:ILE:O	1:E:132:PRO:C	2.56	0.45
1:E:131:ILE:HB	1:E:132:PRO:CD	2.48	0.44
1:E:1:ILE:HD13	1:E:1:ILE:HG21	1.71	0.44
1:E:255:VAL:HG22	1:E:308:GLN:HB3	1.99	0.44
1:E:194:THR:N	1:E:195:PRO:CD	2.76	0.43
1:E:175:LEU:HA	1:E:175:LEU:HD23	1.71	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:226:ASP:O	1:E:227:ASN:HB2	2.19	0.43
1:E:83:TYR:CG	1:E:176:VAL:HG22	2.54	0.42
1:E:29:TYR:CE1	1:E:56:ALA:HB2	2.54	0.42
1:E:142:HIS:CG	1:E:169:SER:HB3	2.54	0.42
1:E:32:ASP:O	1:E:38:GLY:HA2	2.20	0.41
1:E:202:LEU:HD21	3:E:1323:LYS:HB3	2.02	0.41
1:E:35:ARG:HH11	1:E:35:ARG:HD2	1.72	0.41
1:E:1:ILE:HB	1:E:31:GLN:HE22	1.86	0.41
1:E:194:THR:N	1:E:195:PRO:HD3	2.36	0.40
1:E:300:SER:HB2	1:E:302:GLU:OE1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	316/316 (100%)	304 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	254/252 (101%)	242 (95%)	12 (5%)	23	6

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1	ILE
1	E	18	LYS
1	E	20	ILE
1	E	45	LYS
1	E	89	ASN
1	E	156	ILE
1	E	161[A]	SER
1	E	161[B]	SER
1	E	182	LYS
1	E	185	ASP
1	E	260	ARG
1	E	316	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	31	GLN
1	E	33	ASN
1	E	97	ASN
1	E	290	GLN
1	E	301	GLN
1	E	308	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	DMS	E	324	-	3,3,3	0.37	0	3,3,3	0.82	0
2	VAL	E	1322	3	4,6,7	1.02	0	6,7,9	1.45	2 (33%)
3	LYS	E	1323	2	8,9,9	1.23	1 (12%)	7,10,10	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VAL	E	1322	3	-	1/5/6/8	-
3	LYS	E	1323	2	-	1/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1323	LYS	O-C	3.01	1.31	1.22

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1322	VAL	CB-CA-C	-2.22	109.84	112.87
2	E	1322	VAL	O-C-CA	-2.01	119.61	124.77

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1322	VAL	O-C-CA-CB
3	E	1323	LYS	CE-CD-CG-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1323	LYS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	316/316 (100%)	0.00	5 (1%) 70 74	10, 18, 41, 62	2 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	2	THR	3.2
1	E	196	GLY	2.6
1	E	157	TYR	2.4
1	E	1	ILE	2.3
1	E	308	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	LYS	E	1323	10/10	0.86	0.15	24,35,74,100	0
6	DMS	E	324	4/4	0.92	0.12	40,56,57,62	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	VAL	E	1322	7/8	0.93	0.09	18,24,31,68	0
4	CA	E	319	1/1	0.98	0.04	15,15,15,15	0
4	CA	E	317	1/1	0.99	0.03	14,14,14,14	0
4	CA	E	320	1/1	0.99	0.03	20,20,20,20	0
5	ZN	E	321	1/1	0.99	0.02	16,16,16,16	0
4	CA	E	318	1/1	0.99	0.03	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.