



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:53 AM UTC

PDB ID : 1TMN / pdb_00001tmn
Title : Binding of n-carboxymethyl dipeptide inhibitors to thermolysin determined by x-ray crystallography. a novel class of transition-state analogues for zinc peptidases
Authors : Monzingo, A.F.; Matthews, B.W.
Deposited on : 1987-06-29
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	316	69% 24% 5% •


i

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THERMOLYSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	316	Total	C	N	O	S	0	0	0
			2432	1528	408	494	2			

- Molecule 2 is N-[(1R)-1-carboxy-3-phenylpropyl]-L-leucyl-L-tryptophan (CCD ID: 0ZN) (formula: $C_{27}H_{33}N_3O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	E	1	Total	C	N	O	0	0
			35	27	3	5		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	4	Total Ca 4 4	0	0

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total 1	Zn 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	144	Total 144	O 144	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THERMOLYSIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	94.20Å 94.20Å 131.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	14.90 – 1.90 14.90 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (14.90-1.90) 68.9 (14.90-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.90 (at 1.70Å)	Xtriage
Refinement program	EREF	Depositor
R, R_{free}	0.171 , (Not available) 0.161 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	15.0	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 69.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2616	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, 0ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	1.50	25/2491 (1.0%)	1.77	42/3391 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	2	24

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	231	HIS	CE1-NE2	10.76	1.43	1.32
1	E	88	HIS	ND1-CE1	10.61	1.43	1.32
1	E	142	HIS	CE1-NE2	10.26	1.42	1.32
1	E	250	HIS	ND1-CE1	9.61	1.42	1.32
1	E	231	HIS	ND1-CE1	9.21	1.41	1.32
1	E	55	TRP	NE1-CE2	-9.08	1.27	1.37
1	E	216	HIS	ND1-CE1	9.03	1.41	1.32
1	E	250	HIS	CE1-NE2	8.63	1.41	1.32
1	E	183	ASN	N-CA	8.62	1.52	1.46
1	E	88	HIS	CE1-NE2	8.13	1.40	1.32
1	E	105	HIS	ND1-CE1	7.96	1.40	1.32
1	E	186	TRP	NE1-CE2	-7.62	1.29	1.37
1	E	115	TRP	NE1-CE2	-7.56	1.29	1.37
1	E	142	HIS	ND1-CE1	7.42	1.40	1.32
1	E	216	HIS	CE1-NE2	7.42	1.40	1.32
1	E	105	HIS	CE1-NE2	7.14	1.39	1.32
1	E	74	HIS	CE1-NE2	6.71	1.39	1.32
1	E	74	HIS	ND1-CE1	6.69	1.39	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	146	HIS	ND1-CE1	6.30	1.38	1.32
1	E	146	HIS	CE1-NE2	6.20	1.38	1.32
1	E	183	ASN	C-O	5.91	1.26	1.23
1	E	297	GLY	C-N	-5.89	1.26	1.34
1	E	88	HIS	CD2-NE2	5.76	1.44	1.37
1	E	157	TYR	CA-CB	-5.34	1.46	1.52
1	E	231	HIS	CG-ND1	5.08	1.43	1.38

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	183	ASN	CB-CA-C	-11.75	100.78	111.00
1	E	89	ASN	CA-CB-CG	-9.70	102.90	112.60
1	E	96	ASN	CB-CG-ND2	-9.21	102.59	116.40
1	E	96	ASN	CA-CB-CG	-8.54	104.06	112.60
1	E	119	GLU	N-CA-CB	-8.50	98.39	111.56
1	E	181	ASN	CA-CB-CG	-8.14	104.46	112.60
1	E	216	HIS	CA-CB-CG	-7.92	105.88	113.80
1	E	260	ARG	N-CA-C	7.65	122.38	112.89
1	E	308	GLN	CG-CD-NE2	-7.54	105.10	116.40
1	E	101	ARG	CD-NE-CZ	7.46	134.85	124.40
1	E	185	ASP	CA-CB-CG	-7.02	105.58	112.60
1	E	124	ASP	CA-CB-CG	-6.75	105.85	112.60
1	E	172	PHE	CA-CB-CG	-6.56	107.24	113.80
1	E	96	ASN	CB-CA-C	-6.46	100.25	110.85
1	E	108	GLN	O-C-N	6.24	130.72	123.30
1	E	191	ASP	CA-CB-CG	-6.23	106.37	112.60
1	E	293	THR	O-C-N	5.67	127.91	122.07
1	E	256	VAL	CB-CA-C	-5.66	103.30	111.34
1	E	159	ASN	CA-CB-CG	-5.61	106.99	112.60
1	E	197	ILE	CA-CB-CG1	5.59	119.90	110.40
1	E	198	SER	N-CA-CB	-5.58	101.32	110.41
1	E	45	LYS	CB-CA-C	-5.53	103.72	111.80
1	E	214	PRO	N-CA-CB	5.52	108.22	103.31
1	E	291	SER	O-C-N	5.49	127.73	122.07
1	E	233	ASN	CA-CB-CG	-5.41	107.19	112.60
1	E	53	SER	CA-C-O	-5.40	114.81	120.80
1	E	209	ALA	CA-C-N	5.39	128.25	120.38
1	E	209	ALA	C-N-CA	5.39	128.25	120.38
1	E	37	ASP	CB-CA-C	-5.37	101.51	110.81
1	E	220	ARG	CB-CA-C	-5.33	101.84	109.84
1	E	88	HIS	O-C-N	5.32	128.47	122.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	26	THR	CA-CB-CG2	5.31	119.52	110.50
1	E	261	ASP	CB-CA-C	-5.27	102.57	110.90
1	E	128	GLN	OE1-CD-NE2	5.25	127.85	122.60
1	E	133	LEU	N-CA-C	5.23	118.76	112.38
1	E	69	PRO	N-CA-CB	5.21	109.01	103.39
1	E	273	GLN	CB-CA-C	-5.19	101.29	109.34
1	E	264	GLY	CA-C-N	-5.16	112.47	122.06
1	E	264	GLY	C-N-CA	-5.16	112.47	122.06
1	E	119	GLU	CA-C-O	-5.14	115.73	121.23
1	E	166	GLU	CB-CG-CD	-5.03	104.04	112.60
1	E	97	ASN	CA-CB-CG	-5.01	107.58	112.60

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	E	197	ILE	CB
1	E	278	THR	CB

All (24) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	115	TRP	Mainchain
1	E	119	GLU	Sidechain
1	E	124	ASP	Sidechain
1	E	158	GLN	Sidechain
1	E	164	ILE	Mainchain
1	E	166	GLU	Sidechain
1	E	181	ASN	Sidechain
1	E	220	ARG	Mainchain
1	E	223	GLY	Peptide,Mainchain
1	E	26	THR	Mainchain
1	E	261	ASP	Sidechain
1	E	265	LYS	Mainchain
1	E	269	ARG	Sidechain
1	E	273	GLN	Sidechain,Mainchain
1	E	276	THR	Mainchain
1	E	282	SER	Mainchain
1	E	315	VAL	Mainchain
1	E	32	ASP	Sidechain
1	E	37	ASP	Sidechain
1	E	61	GLN	Sidechain
1	E	72	ASP	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	E	96	ASN	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2432	0	2267	33	5
2	E	35	0	31	4	0
3	E	4	0	0	0	0
4	E	1	0	0	0	0
5	E	144	0	0	8	14
All	All	2616	0	2298	36	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:317:0ZN:HD1	2:E:317:0ZN:CD11	0.97	1.08
2:E:317:0ZN:HD1	2:E:317:0ZN:CG2	2.05	0.87
1:E:285:ARG:HD3	1:E:316:LYS:HD3	1.62	0.80
1:E:131:ILE:HG23	1:E:195:PRO:HG3	1.65	0.77
1:E:4:THR:CG2	1:E:24:TYR:HB3	2.29	0.63
1:E:308:GLN:HG2	5:E:388:HOH:O	1.98	0.63
2:E:317:0ZN:HD1	2:E:317:0ZN:NE1	2.12	0.63
1:E:221:TYR:OH	1:E:225:GLN:HG2	1.99	0.63
1:E:280:ASN:HD22	1:E:280:ASN:C	2.09	0.61
1:E:1:ILE:HD13	1:E:29:TYR:CE2	2.40	0.57
1:E:280:ASN:ND2	1:E:283:GLN:H	2.07	0.53
1:E:131:ILE:HB	1:E:132:PRO:HD2	1.90	0.52
1:E:111:ASN:O	2:E:317:0ZN:HE11	2.11	0.50
1:E:50:LEU:HD12	5:E:336:HOH:O	2.12	0.50
1:E:100:ILE:HD11	5:E:451:HOH:O	2.12	0.50
1:E:4:THR:HG23	1:E:24:TYR:HB3	1.94	0.49
1:E:81:TYR:HD1	5:E:451:HOH:O	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:31:GLN:HG3	1:E:40:PHE:CE1	2.49	0.48
1:E:218:SER:HB3	5:E:435:HOH:O	2.13	0.48
1:E:273:GLN:HB3	5:E:372:HOH:O	2.13	0.47
1:E:92:SER:HB2	1:E:97:ASN:HA	1.98	0.46
1:E:76:TYR:OH	1:E:182:LYS:HE3	2.16	0.46
1:E:216:HIS:CE1	5:E:435:HOH:O	2.69	0.46
1:E:282:SER:HB2	1:E:316:LYS:HD2	1.97	0.46
1:E:150:ASP:HA	5:E:379:HOH:O	2.15	0.45
1:E:4:THR:HG22	1:E:24:TYR:HB3	1.97	0.45
1:E:217:TYR:O	1:E:220:ARG:HB2	2.16	0.45
1:E:296:TYR:O	1:E:300:SER:HB3	2.18	0.44
1:E:29:TYR:CE1	1:E:56:ALA:HB2	2.54	0.43
1:E:131:ILE:HB	1:E:132:PRO:CD	2.50	0.42
1:E:300:SER:HB2	1:E:302:GLU:OE1	2.20	0.42
1:E:142:HIS:CG	1:E:169:SER:HB3	2.55	0.41
1:E:137:ILE:HG21	1:E:182:LYS:HD2	2.03	0.41
1:E:131:ILE:CG2	1:E:195:PRO:HG3	2.44	0.41
1:E:97:ASN:HA	1:E:97:ASN:HD22	1.47	0.40
1:E:197:ILE:N	1:E:197:ILE:HD13	2.37	0.40

All (14) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:333:HOH:O	5:E:337:HOH:O[10_664]	0.94	1.26
5:E:372:HOH:O	5:E:463:HOH:O[5_564]	1.52	0.68
5:E:485:HOH:O	5:E:485:HOH:O[7_555]	1.59	0.61
5:E:379:HOH:O	5:E:396:HOH:O[10_664]	1.65	0.55
1:E:116:ASN:O	5:E:336:HOH:O[10_664]	1.66	0.54
5:E:414:HOH:O	5:E:449:HOH:O[8_565]	1.67	0.53
5:E:332:HOH:O	5:E:336:HOH:O[10_664]	1.84	0.36
1:E:273:GLN:CG	5:E:409:HOH:O[5_564]	1.85	0.35
5:E:418:HOH:O	5:E:418:HOH:O[12_565]	1.85	0.35
1:E:6:THR:CB	5:E:473:HOH:O[12_565]	1.87	0.33
5:E:372:HOH:O	5:E:409:HOH:O[5_564]	1.92	0.28
5:E:369:HOH:O	5:E:407:HOH:O[5_564]	1.96	0.24
1:E:273:GLN:CD	5:E:409:HOH:O[5_564]	1.98	0.22
1:E:273:GLN:OE1	5:E:409:HOH:O[5_564]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	314/316 (99%)	302 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	252/252 (100%)	237 (94%)	15 (6%)	17	9

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1	ILE
1	E	14	LEU
1	E	20	ILE
1	E	26	THR
1	E	89	ASN
1	E	96	ASN
1	E	97	ASN
1	E	118	SER
1	E	144	LEU
1	E	197	ILE
1	E	198	SER
1	E	224	THR
1	E	280	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	282	SER
1	E	316	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	21	ASN
1	E	89	ASN
1	E	97	ASN
1	E	280	ASN
1	E	290	GLN
1	E	301	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	0ZN	E	317	4	37,37,37	1.61	5 (13%)	50,50,50	1.52	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0ZN	E	317	4	-	4/33/33/33	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	317	0ZN	CE21-NE1	-6.19	1.27	1.37
2	E	317	0ZN	CD21-CG2	-2.71	1.39	1.44
2	E	317	0ZN	CD11-NE1	2.65	1.41	1.37
2	E	317	0ZN	CD11-CG2	-2.15	1.33	1.36
2	E	317	0ZN	OG2-CB2	-2.13	1.23	1.30

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	317	0ZN	CE3-CD21-CE21	3.56	122.53	118.84
2	E	317	0ZN	CB1-CA-CB2	-2.75	103.83	110.35
2	E	317	0ZN	C-CA1-N	2.73	116.48	110.67
2	E	317	0ZN	CG1-CB-CA1	-2.70	108.15	115.40
2	E	317	0ZN	CG-CB1-CA	-2.67	107.81	113.21
2	E	317	0ZN	CB2-CA-N	2.34	116.95	109.40
2	E	317	0ZN	CA2-CB3-CG2	-2.26	109.16	113.44
2	E	317	0ZN	CZ3-CE3-CD21	-2.20	115.81	119.80
2	E	317	0ZN	C1-CA2-N1	2.04	115.29	110.57
2	E	317	0ZN	OG2-CB2-OG1	-2.03	119.47	124.08

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	317	0ZN	CB2-CA-CB1-CG
2	E	317	0ZN	N-CA-CB1-CG
2	E	317	0ZN	CA2-CB3-CG2-CD11
2	E	317	0ZN	CA2-CB3-CG2-CD21

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	317	0ZN	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	316/316 (100%)	-0.66	0 100 100	7, 14, 24, 33	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	0ZN	E	317	35/35	0.91	0.09	15,23,30,30	0
3	CA	E	320	1/1	0.98	0.05	12,12,12,12	0
3	CA	E	321	1/1	0.98	0.02	16,16,16,16	0
3	CA	E	318	1/1	0.99	0.02	13,13,13,13	0
3	CA	E	319	1/1	1.00	0.01	14,14,14,14	0
4	ZN	E	322	1/1	1.00	0.03	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.

Electron density around 0ZN E 317:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.