



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:04 AM UTC

PDB ID : 1TN6 / pdb_00001tn6
Title : Protein Farnesyltransferase Complexed with a Rap2a Peptide Substrate and a FPP Analog at 1.8Å Resolution
Authors : Reid, T.S.; Terry, K.L.; Casey, P.J.; Beese, L.S.
Deposited on : 2004-06-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	
2	B	437	
3	C	11	
4	D	2	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein farnesyltransferase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2683	1711	467	500	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	380	GLU	-	insertion	UNP P49354
A	381	GLU	-	insertion	UNP P49354
A	382	PHE	-	insertion	UNP P49354


- Molecule 2 is a protein called Protein farnesyltransferase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	410	Total	C	N	O	S	0	0	0
			3228	2065	552	589	22			

- Molecule 3 is a protein called peptide derived from the C-terminus of Rap2a.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	11	Total	C	N	O	S	0	0	0
			78	44	13	20	1			

- Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is ACETIC ACID (CCD ID: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is [(3,7,11-TRIMETHYL-DODECA-2,6,10-TRIENYLOXYCARBAMOYL)-METHYL]-PHOSPHONIC ACID (CCD ID: FII) (formula: $C_{17}H_{30}NO_5P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	N	O	P	0	0
			24	17	1	5	1		


- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Zn	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	301	Total	O	0	0
			301	301		
8	B	348	Total	O	0	0
			348	348		
8	C	10	Total	O	0	0
			10	10		

- Molecule 1: Protein farnesyltransferase alpha subunit



Chain D:

100%

GLU1
FRU2

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	178.47Å 178.47Å 64.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.61 – 1.80 49.61 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.5 (49.61-1.80) 92.7 (49.61-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 1.81Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.179 , 0.199 0.180 , 0.200	Depositor DCC
R_{free} test set	5027 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6704	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, GLC, ACY, ZN, FII

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2750	0.83	10/3735 (0.3%)
2	B	0.38	0/3317	0.87	9/4508 (0.2%)
3	C	0.41	0/78	0.85	0/105
All	All	0.37	0/6145	0.85	19/8348 (0.2%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLU	N-CA-C	9.16	120.95	110.97
1	A	285	GLN	N-CA-C	7.43	119.38	111.28
2	B	875	HIS	N-CA-C	7.25	120.23	108.41
2	B	795	LEU	N-CA-C	6.38	119.96	110.52
2	B	865	TYR	N-CA-C	6.32	119.15	111.82
1	A	85	ASN	CA-C-N	5.99	125.95	119.78
1	A	85	ASN	C-N-CA	5.99	125.95	119.78
1	A	183	SER	N-CA-C	5.94	119.36	111.75
2	B	801	SER	N-CA-C	-5.91	104.92	111.71
2	B	851	LEU	N-CA-C	5.89	116.87	108.74
2	B	664	GLY	N-CA-C	5.61	122.97	114.61
1	A	348	LYS	N-CA-C	5.52	119.74	112.89
1	A	270	VAL	CA-C-N	5.47	125.81	119.47
1	A	270	VAL	C-N-CA	5.47	125.81	119.47
2	B	828	TRP	N-CA-C	-5.39	103.27	110.55
1	A	217	ASP	N-CA-C	5.37	122.23	110.80
1	A	180	ARG	N-CA-C	-5.33	105.42	112.24
2	B	903	ILE	N-CA-C	-5.32	100.07	108.90
2	B	692	LEU	N-CA-C	-5.01	103.06	110.48

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2683	0	2601	19	0
2	B	3228	0	3147	33	0
3	C	78	0	65	2	0
4	D	23	0	21	0	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	B	24	0	28	1	0
7	B	1	0	0	0	0
8	A	301	0	0	4	0
8	B	348	0	0	4	0
8	C	10	0	0	0	0
All	All	6704	0	5868	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:634:GLN:HE22	2:B:673:ASN:H	1.25	0.83
2:B:547:GLU:HG3	8:B:1969:HOH:O	1.84	0.74
2:B:918:LYS:HE3	8:B:1730:HOH:O	1.93	0.68
1:A:135:HIS:HD2	8:A:1875:HOH:O	1.76	0.67
1:A:109:ARG:HG3	8:A:1256:HOH:O	1.97	0.65
1:A:330:LYS:HE2	1:A:367:HIS:O	1.99	0.63
1:A:330:LYS:HE3	1:A:367:HIS:HB3	1.81	0.63
2:B:577:LYS:HD3	2:B:846:PRO:O	2.02	0.60
2:B:801:SER:O	2:B:805:ALA:HB3	2.02	0.59
2:B:634:GLN:NE2	2:B:673:ASN:H	1.99	0.58
2:B:880:ALA:C	2:B:881:MET:HE2	2.29	0.58
1:A:316:VAL:O	1:A:320:GLU:HG3	2.03	0.57
1:A:303:GLN:N	1:A:304:PRO:HD2	2.20	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:GLN:O	1:A:302:LEU:HD23	2.05	0.56
1:A:335:ASN:O	1:A:339:GLU:HG3	2.06	0.56
2:B:771:LYS:HD2	2:B:772:SER:N	2.20	0.56
2:B:739:ILE:HB	2:B:752:THR:HA	1.89	0.55
2:B:881:MET:O	2:B:882:LEU:HD12	2.06	0.55
2:B:881:MET:C	2:B:882:LEU:HD12	2.32	0.54
2:B:574:GLN:HB3	2:B:577:LYS:HD2	1.91	0.52
2:B:908:VAL:O	2:B:912:THR:HG23	2.09	0.51
2:B:701:VAL:HG12	2:B:751:TYR:CD2	2.46	0.51
2:B:753:PHE:HA	2:B:807:LEU:HD21	1.93	0.50
2:B:885:VAL:HG23	2:B:885:VAL:O	2.13	0.49
1:A:114:GLU:OE2	1:A:118:LYS:HE3	2.13	0.49
2:B:635:SER:OG	2:B:637:GLU:HG2	2.14	0.48
2:B:516:SER:HB2	2:B:517:PRO:HD2	1.96	0.48
2:B:852:ASP:C	2:B:852:ASP:OD1	2.58	0.46
2:B:526:LEU:HD13	2:B:563:LYS:HB2	1.98	0.46
2:B:696:GLY:HA2	8:B:1901:HOH:O	2.15	0.46
1:A:311:LEU:C	1:A:311:LEU:HD23	2.41	0.45
2:B:692:LEU:HD23	2:B:699:VAL:CG2	2.46	0.45
1:A:58:LEU:HD12	1:A:125:GLU:HB3	1.99	0.45
1:A:135:HIS:CD2	8:A:1875:HOH:O	2.61	0.45
2:B:858:ARG:O	3:C:1999:ASP:HA	2.17	0.44
1:A:330:LYS:NZ	1:A:369:THR:OG1	2.48	0.43
2:B:905:PRO:HD2	8:B:1546:HOH:O	2.19	0.43
2:B:589:LEU:CD1	2:B:603:LEU:HD13	2.48	0.43
2:B:652:PRO:HD3	3:C:2009:GLN:OE1	2.18	0.43
1:A:328:ASP:O	1:A:329:ASN:HB2	2.18	0.43
1:A:84:PRO:C	1:A:86:PRO:HD3	2.44	0.43
2:B:551:VAL:O	2:B:555:ILE:HG12	2.19	0.43
1:A:112:ARG:HA	1:A:140:LEU:CD2	2.49	0.42
1:A:225:GLN:HG2	8:A:1286:HOH:O	2.20	0.42
2:B:806:GLY:O	2:B:809:PRO:HD2	2.20	0.41
2:B:808:LEU:HD13	2:B:808:LEU:HA	1.93	0.41
1:A:96:PHE:HA	1:A:126:LEU:HD13	2.03	0.41
1:A:77:PRO:HB2	1:A:101:ASP:HB3	2.03	0.41
6:B:1:FII:H352	6:B:1:FII:H301	1.85	0.41
2:B:651:ALA:HB3	2:B:652:PRO:CD	2.51	0.40
2:B:701:VAL:HG12	2:B:751:TYR:HD2	1.86	0.40
2:B:797:ASP:HB3	2:B:800:TYR:CD1	2.56	0.40
2:B:850:LEU:HB2	2:B:863:THR:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/382 (82%)	302 (96%)	11 (4%)	0	100	100
2	B	408/437 (93%)	400 (98%)	8 (2%)	0	100	100
3	C	9/11 (82%)	8 (89%)	1 (11%)	0	100	100
All	All	730/830 (88%)	710 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/344 (86%)	293 (100%)	1 (0%)	86	86
2	B	346/370 (94%)	343 (99%)	3 (1%)	70	67
3	C	9/9 (100%)	9 (100%)	0	100	100
All	All	649/723 (90%)	645 (99%)	4 (1%)	78	77

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	342	GLU
2	B	620	GLN
2	B	763	LYS
2	B	851	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	GLN
1	A	234	ASN
1	A	278	ASN
1	A	329	ASN
1	A	364	GLN
2	B	588	GLN
2	B	634	GLN
2	B	775	GLN
2	B	827	HIS
2	B	917	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	D	1	4	11,11,12	3.01	4 (36%)	15,15,17	1.48	2 (13%)
4	FRU	D	2	4	11,12,12	1.46	2 (18%)	10,18,18	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	D	1	4	-	0/2/19/22	0/1/1/1
4	FRU	D	2	4	-	0/5/24/24	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1	GLC	C2-C3	8.68	1.65	1.52
4	D	2	FRU	O2-C2	3.76	1.47	1.40
4	D	1	GLC	C4-C5	2.69	1.58	1.53
4	D	1	GLC	O5-C5	2.65	1.48	1.43
4	D	1	GLC	O5-C1	2.03	1.47	1.43
4	D	2	FRU	C1-C2	2.00	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	GLC	C1-O5-C5	4.16	117.76	112.19
4	D	1	GLC	O3-C3-C2	-2.15	105.66	110.05


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACY	A	3004	-	3,3,3	1.28	0	3,3,3	1.66	1 (33%)
5	ACY	B	3001	-	3,3,3	1.28	1 (33%)	3,3,3	1.65	1 (33%)
6	FII	B	1	-	23,23,23	1.56	5 (21%)	27,29,29	1.24	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FII	B	1	-	-	0/23/24/24	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1	FII	P46-O50	-4.54	1.44	1.55
6	B	1	FII	P46-O51	-3.23	1.47	1.55
6	B	1	FII	C22-C15	2.23	1.38	1.33
6	B	1	FII	P46-O49	-2.21	1.45	1.50
6	B	1	FII	P46-C45	2.07	1.83	1.79
5	B	3001	ACY	O-C	2.02	1.31	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1	FII	C45-C43-N42	-2.79	112.54	115.19
5	A	3004	ACY	O-C-CH3	-2.26	113.27	122.53
5	B	3001	ACY	O-C-CH3	-2.24	113.33	122.53
6	B	1	FII	O50-P46-C45	2.23	111.51	106.84

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1	FII	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	315/382 (82%)	0.69	32 (10%)	12 10	14, 23, 38, 54	0
2	B	410/437 (93%)	0.18	27 (6%)	24 23	12, 18, 30, 42	0
3	C	11/11 (100%)	2.91	8 (72%)	0 0	21, 45, 56, 57	0
All	All	736/830 (88%)	0.44	67 (9%)	15 13	12, 20, 35, 57	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	2005	ALA	6.3
1	A	302	LEU	6.2
1	A	369	THR	6.1
1	A	55	PHE	6.0
3	C	2003	ALA	5.2
1	A	56	VAL	4.7
1	A	368	SER	4.5
1	A	305	SER	4.5
2	B	924	GLU	4.4
3	C	2002	THR	4.4
1	A	304	PRO	4.3
1	A	328	ASP	4.3
2	B	515	SER	3.9
1	A	326	GLN	3.9
2	B	766	ARG	3.8
1	A	307	SER	3.7
3	C	2001	PRO	3.7
1	A	59	ASP	3.6
3	C	2004	SER	3.6
1	A	300	LEU	3.6
1	A	303	GLN	3.5
2	B	819	GLY	3.4
1	A	306	HIS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	879	GLY	3.3
1	A	301	ASP	3.3
1	A	143	SER	3.3
2	B	923	PHE	3.3
2	B	516	SER	3.3
1	A	109	ARG	3.1
1	A	287	ARG	3.1
2	B	917	GLN	3.1
2	B	618	ILE	3.0
2	B	881	MET	3.0
2	B	620	GLN	2.9
1	A	329	ASN	2.8
2	B	880	ALA	2.8
1	A	327	CYS	2.8
1	A	325	ASN	2.7
2	B	617	PRO	2.7
2	B	720	ASP	2.6
1	A	58	LEU	2.6
2	B	576	GLU	2.5
2	B	882	LEU	2.5
2	B	587	ARG	2.5
1	A	268	LYS	2.5
1	A	228	LYS	2.4
1	A	57	SER	2.4
1	A	297	ASN	2.4
2	B	885	VAL	2.4
3	C	2006	CYS	2.3
1	A	150	GLU	2.3
1	A	331	GLU	2.3
1	A	347	GLU	2.2
2	B	553	GLU	2.2
2	B	825	MET	2.2
1	A	299	LEU	2.2
2	B	719	PRO	2.2
2	B	813	ARG	2.2
2	B	622	VAL	2.2
1	A	86	PRO	2.2
2	B	588	GLN	2.1
2	B	817	ALA	2.1
3	C	1999	ASP	2.1
2	B	590	THR	2.1
1	A	147	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	2000	ASP	2.0
2	B	878	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GLC	D	1	11/12	0.94	0.09	24,25,26,26	0
4	FRU	D	2	12/12	0.94	0.09	21,23,24,24	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	FII	B	1	24/24	0.85	0.16	16,25,29,31	0
5	ACY	B	3001	4/4	0.88	0.18	37,37,38,38	0
5	ACY	A	3004	4/4	0.95	0.10	23,23,24,24	0
7	ZN	B	1001	1/1	1.00	0.02	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.