



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:26 PM UTC

PDB ID : 4TNA / pdb_00004tna
Title : FURTHER REFINEMENT OF THE STRUCTURE OF YEAST T-RNA-PHE
Authors : Hingerty, B.E.; Brown, R.S.; Jack, A.
Deposited on : 1978-04-12
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6492 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)
RNA backbone	3983	1003 (2.78-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	7% 51% 30% 12%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called TRNAPHE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	76	Total	C	N	O	P	0	0	0
			1652	746	294	536	76			


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Mg	0	0
			4	4		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRNAPHE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.30Å 33.40Å 63.00Å 90.00° 90.25° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.50 63.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50) 94.7 (63.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.00Å)	Xtriage
Refinement program	JACK-LEVITT	Depositor
R, R_{free}	(Not available) , (Not available) 0.300 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.710	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.081 for h,-k,-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	1656	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M2G, MG, 1MA, 5MU, PSU, 7MG, 5MC, 2MG, OMG, YG, H2U, OMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.37	2/1487 (0.1%)	1.78	27/2315 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	44

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	G	OP3-P	6.21	1.60	1.48
1	A	19	G	C6-N1	-6.10	1.27	1.39

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	A	C4'-C3'-C2'	-7.48	95.12	102.60
1	A	3	G	C4'-C3'-C2'	-6.58	96.02	102.60
1	A	13	C	C4'-C3'-C2'	-6.40	96.20	102.60
1	A	3	G	P-O5'-C5'	-6.31	111.44	120.90
1	A	7	U	C3'-C2'-O2'	-6.08	105.48	114.60
1	A	68	U	C4'-C3'-C2'	-6.08	96.52	102.60
1	A	5	A	C4'-C3'-C2'	-6.07	96.53	102.60
1	A	25	C	N1-C1'-C2'	-6.05	102.93	112.00
1	A	3	G	C5'-C4'-C3'	-5.96	107.06	116.00
1	A	33	U	C4'-C3'-C2'	-5.83	96.77	102.60
1	A	22	G	P-O5'-C5'	-5.83	112.16	120.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	U	O5'-C5'-C4'	-5.67	102.99	111.50
1	A	64	A	C4'-C3'-C2'	-5.58	97.02	102.60
1	A	74	C	C4'-C3'-C2'	-5.54	97.06	102.60
1	A	65	G	N9-C1'-C2'	-5.50	103.75	112.00
1	A	57	G	C2'-C3'-O3'	-5.48	105.47	113.70
1	A	24	G	O5'-C5'-C4'	-5.36	103.46	111.50
1	A	70	C	C4'-C3'-C2'	-5.27	97.33	102.60
1	A	75	C	C2'-C3'-O3'	-5.26	105.82	113.70
1	A	67	A	O5'-C5'-C4'	-5.25	103.62	111.50
1	A	6	U	C4'-C3'-C2'	-5.22	97.38	102.60
1	A	42	G	C4'-C3'-C2'	-5.20	97.40	102.60
1	A	66	A	O5'-C5'-C4'	-5.19	103.72	111.50
1	A	11	C	O5'-C5'-C4'	-5.18	103.73	111.50
1	A	22	G	C4'-C3'-C2'	-5.16	97.44	102.60
1	A	43	G	O5'-C5'-C4'	-5.10	103.85	111.50
1	A	5	A	O5'-C5'-C4'	-5.10	103.86	111.50

There are no chirality outliers.

All (44) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	G	Sidechain
1	A	12	U	Sidechain
1	A	13	C	Sidechain
1	A	15	G	Sidechain
1	A	18	G	Sidechain
1	A	19	G	Sidechain
1	A	2	C	Sidechain
1	A	20	G	Sidechain
1	A	21	A	Sidechain
1	A	23	A	Sidechain
1	A	24	G	Sidechain
1	A	25	C	Sidechain
1	A	27	C	Sidechain
1	A	28	C	Sidechain
1	A	29	A	Sidechain
1	A	3	G	Sidechain
1	A	30	G	Sidechain
1	A	33	U	Sidechain
1	A	38	A	Sidechain
1	A	41	U	Sidechain
1	A	42	G	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	43	G	Sidechain
1	A	44	A	Sidechain
1	A	47	U	Sidechain
1	A	5	A	Sidechain
1	A	50	U	Sidechain
1	A	51	G	Sidechain
1	A	53	G	Sidechain
1	A	56	C	Sidechain
1	A	57	G	Sidechain
1	A	59	U	Sidechain
1	A	6	U	Sidechain
1	A	60	C	Sidechain
1	A	61	C	Sidechain
1	A	62	A	Sidechain
1	A	64	A	Sidechain
1	A	68	U	Sidechain
1	A	71	G	Sidechain
1	A	72	C	Sidechain
1	A	73	A	Sidechain
1	A	74	C	Sidechain
1	A	75	C	Sidechain
1	A	76	A	Sidechain
1	A	8	U	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1652	0	859	13	0
2	A	4	0	0	0	0
All	All	1656	0	859	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:C:H2'	1:A:75:C:H6	1.57	0.69
1:A:74:C:H2'	1:A:75:C:C6	2.27	0.68
1:A:28:C:H2'	1:A:29:A:C8	2.48	0.49
1:A:16:H2U:HN3	1:A:60:C:H1'	1.78	0.48
1:A:16:H2U:P	1:A:16:H2U:H51	2.57	0.45
1:A:32:OMC:H2'	1:A:33:U:C6	2.53	0.43
1:A:16:H2U:H51	1:A:16:H2U:OP2	2.18	0.43
1:A:31:A:H2'	1:A:32:OMC:O4'	2.19	0.43
1:A:16:H2U:H2'	1:A:16:H2U:H62	1.51	0.43
1:A:17:H2U:H2'	1:A:17:H2U:H62	1.73	0.43
1:A:16:H2U:H4'	1:A:17:H2U:OP1	2.19	0.43
1:A:14:A:H2'	1:A:15:G:O4'	2.21	0.41
1:A:3:G:H2'	1:A:4:G:C8	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	73/76 (96%)	16 (21%)	4 (5%)

All (16) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	9	A
1	A	13	C
1	A	16	H2U
1	A	17	H2U
1	A	18	G
1	A	19	G

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	21	A
1	A	22	G
1	A	46	7MG
1	A	47	U
1	A	48	C
1	A	49	5MC
1	A	57	G
1	A	58	1MA
1	A	59	U
1	A	76	A

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	18	G
1	A	21	A
1	A	46	7MG
1	A	58	1MA

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

14 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	M2G	A	26	1	24,27,28	1.12	3 (12%)	33,40,43	1.80	6 (18%)
1	H2U	A	16	1	18,21,22	1.32	3 (16%)	19,30,33	1.70	3 (15%)
1	5MC	A	40	1	19,22,23	1.71	4 (21%)	26,32,35	1.32	5 (19%)
1	2MG	A	10	1	23,26,27	1.12	2 (8%)	33,38,41	2.15	9 (27%)
1	7MG	A	46	1	23,26,27	3.45	3 (13%)	27,39,42	2.36	10 (37%)
1	PSU	A	55	1	18,21,22	1.66	4 (22%)	21,30,33	2.01	5 (23%)
1	OMG	A	34	1	23,26,27	0.96	3 (13%)	32,38,41	1.91	9 (28%)
1	1MA	A	58	1	21,25,26	1.04	0	30,37,40	1.84	7 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	YG	A	37	1	38,42,43	1.70	6 (15%)	45,62,65	2.06	14 (31%)
1	5MC	A	49	1	19,22,23	2.26	4 (21%)	26,32,35	1.39	4 (15%)
1	PSU	A	39	1	18,21,22	1.55	3 (16%)	21,30,33	1.93	5 (23%)
1	5MU	A	54	1	19,22,23	1.55	4 (21%)	27,32,35	2.03	7 (25%)
1	OMC	A	32	1	19,22,23	1.03	2 (10%)	25,31,34	1.05	1 (4%)
1	H2U	A	17	1	18,21,22	1.21	2 (11%)	19,30,33	1.60	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	M2G	A	26	1	-	1/11/29/30	0/3/3/3
1	H2U	A	16	1	-	4/7/38/39	0/2/2/2
1	5MC	A	40	1	-	0/7/25/26	0/2/2/2
1	2MG	A	10	1	-	0/9/27/28	0/3/3/3
1	7MG	A	46	1	-	2/7/37/38	0/3/3/3
1	PSU	A	55	1	-	2/7/25/26	0/2/2/2
1	OMG	A	34	1	-	0/9/27/28	0/3/3/3
1	1MA	A	58	1	-	1/7/25/26	0/3/3/3
1	YG	A	37	1	-	5/24/42/43	0/4/4/4
1	5MC	A	49	1	-	2/7/25/26	0/2/2/2
1	PSU	A	39	1	-	0/7/25/26	0/2/2/2
1	5MU	A	54	1	-	0/7/25/26	0/2/2/2
1	OMC	A	32	1	-	0/9/27/28	0/2/2/2
1	H2U	A	17	1	-	4/7/38/39	0/2/2/2

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	46	7MG	C8-N9	-15.41	1.35	1.45
1	A	49	5MC	C5-C4	-7.49	1.38	1.44
1	A	37	YG	O23-C21	-5.71	1.24	1.34
1	A	40	5MC	C5-C4	-5.26	1.40	1.44
1	A	37	YG	C6-N1	-4.73	1.31	1.42
1	A	39	PSU	C6-C5	4.26	1.40	1.35
1	A	55	PSU	C6-C5	4.18	1.39	1.35
1	A	49	5MC	C6-C5	3.79	1.40	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	37	YG	O18-C16	-3.46	1.24	1.33
1	A	54	5MU	C4-C5	-3.44	1.39	1.44
1	A	46	7MG	C5-N7	3.33	1.39	1.35
1	A	40	5MC	C6-C5	3.31	1.40	1.34
1	A	10	2MG	C6-N1	-3.21	1.32	1.38
1	A	17	H2U	C4-N3	-3.09	1.32	1.37
1	A	16	H2U	C2-N1	-3.08	1.31	1.35
1	A	55	PSU	C4-N3	-3.05	1.33	1.38
1	A	16	H2U	C4-N3	-3.01	1.32	1.37
1	A	49	5MC	C2-N1	-2.99	1.33	1.40
1	A	49	5MC	C6-N1	-2.94	1.33	1.38
1	A	16	H2U	C2-N3	-2.92	1.32	1.38
1	A	46	7MG	C6-N1	-2.92	1.33	1.38
1	A	17	H2U	C2-N3	-2.73	1.33	1.38
1	A	32	OMC	C2-N1	-2.68	1.34	1.40
1	A	54	5MU	C6-C5	2.66	1.38	1.34
1	A	39	PSU	C4-N3	-2.65	1.33	1.38
1	A	54	5MU	C4-N3	-2.64	1.33	1.38
1	A	40	5MC	C6-N1	-2.55	1.33	1.38
1	A	26	M2G	C6-N1	-2.55	1.34	1.38
1	A	34	OMG	C6-N1	-2.49	1.34	1.38
1	A	40	5MC	C2-N1	-2.41	1.35	1.40
1	A	26	M2G	C4-N3	2.36	1.39	1.34
1	A	37	YG	C15-N20	2.31	1.50	1.45
1	A	54	5MU	C6-N1	-2.29	1.34	1.38
1	A	37	YG	C2-N2	2.23	1.37	1.33
1	A	39	PSU	C2-N3	-2.20	1.33	1.37
1	A	55	PSU	C4-C5	-2.18	1.38	1.44
1	A	34	OMG	C8-N7	2.15	1.38	1.32
1	A	32	OMC	C6-C5	2.12	1.40	1.35
1	A	26	M2G	C8-N7	2.08	1.38	1.32
1	A	55	PSU	C2-N1	-2.02	1.34	1.36
1	A	34	OMG	C4-N3	2.01	1.38	1.34
1	A	10	2MG	C2-N1	-2.01	1.33	1.36
1	A	37	YG	C5-C6	-2.00	1.40	1.45

All (88) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	2MG	C2-N3-C4	6.50	120.13	112.00
1	A	10	2MG	C5-C4-N3	-5.68	119.36	128.39
1	A	26	M2G	C5-C4-N3	-5.59	119.50	128.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	58	1MA	C5-C4-N3	-5.32	119.44	127.27
1	A	37	YG	O23-C21-O22	-5.28	116.93	124.62
1	A	39	PSU	N1-C2-N3	5.23	120.69	115.17
1	A	34	OMG	C5-C4-N3	-5.09	120.29	128.39
1	A	46	7MG	N9-C4-N3	4.84	132.56	125.46
1	A	55	PSU	N1-C2-N3	4.84	120.27	115.17
1	A	37	YG	C24-O23-C21	-4.64	110.26	115.63
1	A	46	7MG	C5-C6-N1	4.57	118.99	110.94
1	A	54	5MU	C4-N3-C2	-4.57	121.35	127.34
1	A	54	5MU	C5-C4-N3	4.55	119.28	115.32
1	A	34	OMG	C2-N3-C4	4.42	119.92	112.30
1	A	46	7MG	C2-N3-C4	4.32	119.74	112.30
1	A	16	H2U	O2-C2-N1	-4.22	118.03	123.10
1	A	54	5MU	N3-C2-N1	4.18	120.33	114.89
1	A	37	YG	O23-C21-N20	4.08	117.64	110.77
1	A	17	H2U	N3-C2-N1	4.07	120.74	116.65
1	A	54	5MU	O4-C4-C5	-4.06	120.27	124.92
1	A	46	7MG	C5-C4-N3	-4.04	120.55	128.13
1	A	10	2MG	N9-C4-N3	3.90	133.74	125.95
1	A	55	PSU	O2-C2-N1	-3.83	118.83	122.79
1	A	26	M2G	N9-C4-N3	3.83	133.61	125.95
1	A	58	1MA	N1-C2-N3	-3.76	121.53	126.00
1	A	26	M2G	C2-N3-C4	3.69	119.33	112.51
1	A	55	PSU	C4-N3-C2	-3.67	121.31	126.37
1	A	58	1MA	C2-N3-C4	3.65	119.70	112.53
1	A	39	PSU	C4-N3-C2	-3.61	121.40	126.37
1	A	49	5MC	C5-C6-N1	-3.45	119.57	123.31
1	A	37	YG	O18-C16-C15	3.42	120.18	111.49
1	A	16	H2U	C5-C4-N3	3.39	120.30	116.69
1	A	37	YG	N3-C2-N2	-3.34	122.02	126.62
1	A	39	PSU	O2-C2-N1	-3.31	119.37	122.79
1	A	34	OMG	N9-C4-N3	3.29	132.53	125.95
1	A	46	7MG	O4'-C1'-N9	3.27	113.76	109.30
1	A	58	1MA	N9-C4-N3	3.24	134.29	126.90
1	A	37	YG	C14-C13-C12	-3.23	106.75	113.36
1	A	16	H2U	N3-C2-N1	3.20	119.86	116.65
1	A	46	7MG	O6-C6-C5	-3.19	119.78	127.62
1	A	34	OMG	C2'-C1'-N9	-3.13	108.31	114.24
1	A	40	5MC	C5-C6-N1	-3.09	119.96	123.31
1	A	46	7MG	N9-C8-N7	3.09	107.75	103.37
1	A	37	YG	C14-C15-C16	-3.08	101.76	110.38
1	A	17	H2U	C5-C4-N3	3.03	119.91	116.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	5MU	C5-C6-N1	-2.97	120.08	123.31
1	A	58	1MA	O4'-C4'-C5'	-2.89	100.09	109.33
1	A	26	M2G	N9-C8-N7	-2.86	108.10	113.40
1	A	34	OMG	N9-C8-N7	-2.86	108.10	113.40
1	A	37	YG	C5'-C4'-C3'	-2.82	105.07	115.21
1	A	40	5MC	O2-C2-N3	-2.79	117.94	122.33
1	A	37	YG	C19-O18-C16	-2.78	109.60	115.92
1	A	46	7MG	C6-C5-C4	-2.76	117.55	122.40
1	A	10	2MG	O6-C6-C5	-2.73	119.32	126.53
1	A	37	YG	C8-N9-C4	2.71	109.15	106.54
1	A	46	7MG	C5'-C4'-C3'	-2.69	105.53	115.21
1	A	58	1MA	N9-C8-N7	-2.68	108.44	113.40
1	A	40	5MC	C5-C4-N3	-2.62	119.07	121.75
1	A	10	2MG	C5-C6-N1	2.60	119.88	113.25
1	A	40	5MC	C2'-C3'-C4'	-2.60	97.59	102.61
1	A	17	H2U	O2-C2-N1	-2.57	120.01	123.10
1	A	34	OMG	C5-C6-N1	2.54	119.72	113.25
1	A	55	PSU	C2'-C3'-C4'	-2.54	97.71	102.61
1	A	10	2MG	N9-C8-N7	-2.53	108.71	113.40
1	A	37	YG	N9-C8-N7	-2.49	108.78	113.40
1	A	10	2MG	C2-N1-C6	-2.47	121.56	124.55
1	A	34	OMG	O6-C6-C5	-2.47	120.02	126.53
1	A	49	5MC	C2'-C3'-C4'	-2.46	97.86	102.61
1	A	26	M2G	C5-C6-N1	2.45	119.49	113.25
1	A	26	M2G	O6-C6-C5	-2.41	120.17	126.53
1	A	49	5MC	C5-C4-N3	-2.37	119.32	121.75
1	A	49	5MC	O2-C2-N3	-2.37	118.59	122.33
1	A	37	YG	C10-C11-N2	2.35	124.04	119.32
1	A	46	7MG	C4-C5-N7	2.34	108.14	105.38
1	A	39	PSU	C5'-C4'-C3'	-2.33	106.84	115.21
1	A	32	OMC	CM2-O2'-C2'	-2.21	108.79	114.47
1	A	34	OMG	CM2-O2'-C2'	-2.21	108.81	114.47
1	A	10	2MG	N1-C2-N2	2.19	118.80	116.56
1	A	39	PSU	C6-N1-C2	-2.19	120.66	122.69
1	A	55	PSU	O2'-C2'-C3'	-2.18	104.84	111.82
1	A	58	1MA	O3'-C3'-C2'	-2.16	104.88	111.82
1	A	54	5MU	O2-C2-N1	-2.14	120.01	122.80
1	A	10	2MG	O3'-C3'-C4'	-2.14	104.95	111.08
1	A	37	YG	C5-C4-N3	-2.09	122.30	123.99
1	A	37	YG	C2-N1-C12	-2.08	106.19	109.94
1	A	40	5MC	CM5-C5-C6	-2.04	120.09	122.85
1	A	34	OMG	C2-N1-C6	-2.04	121.42	125.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	5MU	C5M-C5-C6	-2.03	120.11	122.85

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	16	H2U	O4'-C1'-N1-C6
1	A	16	H2U	C2'-C1'-N1-C6
1	A	17	H2U	O4'-C1'-N1-C6
1	A	17	H2U	C2'-C1'-N1-C6
1	A	46	7MG	C2'-C1'-N9-C8
1	A	49	5MC	O4'-C4'-C5'-O5'
1	A	58	1MA	O4'-C4'-C5'-O5'
1	A	37	YG	C12-C13-C14-C15
1	A	16	H2U	C2'-C1'-N1-C2
1	A	46	7MG	O4'-C4'-C5'-O5'
1	A	37	YG	O23-C21-N20-C15
1	A	37	YG	N20-C15-C16-O18
1	A	26	M2G	O4'-C4'-C5'-O5'
1	A	49	5MC	C3'-C4'-C5'-O5'
1	A	16	H2U	O4'-C1'-N1-C2
1	A	17	H2U	C2'-C1'-N1-C2
1	A	37	YG	C16-C15-N20-C21
1	A	55	PSU	O4'-C1'-C5-C4
1	A	17	H2U	O4'-C1'-N1-C2
1	A	55	PSU	O4'-C1'-C5-C6
1	A	37	YG	N20-C15-C16-O17

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	16	H2U	5	0
1	A	32	OMC	2	0
1	A	17	H2U	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	62/76 (81%)	-1.37	1 (1%) 70 67	0, 0, 0, 0	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	76	A	2.5

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	YG	A	37	39/40	0.94	0.09	0,0,0,0	0
1	H2U	A	17	20/21	0.96	0.10	0,0,0,0	0
1	H2U	A	16	20/21	0.96	0.13	0,0,0,0	0
1	OMC	A	32	21/22	0.97	0.06	0,0,0,0	0
1	OMG	A	34	24/25	0.98	0.06	0,0,0,0	0
1	M2G	A	26	25/26	0.98	0.06	0,0,0,0	0
1	PSU	A	39	20/21	0.98	0.05	0,0,0,0	0
1	5MC	A	40	21/22	0.98	0.05	0,0,0,0	0
1	2MG	A	10	24/25	0.99	0.04	0,0,0,0	0
1	7MG	A	46	24/25	0.99	0.04	0,0,0,0	0
1	5MC	A	49	21/22	0.99	0.06	0,0,0,0	0
1	5MU	A	54	21/22	0.99	0.05	0,0,0,0	0
1	PSU	A	55	20/21	0.99	0.04	0,0,0,0	0
1	1MA	A	58	23/24	0.99	0.06	0,0,0,0	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	77	1/1	0.98	0.06	0,0,0,0	0
2	MG	A	78	1/1	0.98	0.03	0,0,0,0	0
2	MG	A	80	1/1	0.98	0.04	0,0,0,0	0
2	MG	A	79	1/1	0.99	0.12	0,0,0,0	0

6.5 Other polymers [i](#)

There are no such residues in this entry.