



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:04 AM UTC

PDB ID : 6TNB / pdb_00006tnb
Title : X-RAY STRUCTURE OF MPS1 IN COMPLEX WITH COMPOUND 41
Authors : Holton, S.J.; Schulze, V.K.; Klar, U.; Kosemund, D.; Siemeister, G.; Bader, B.; Prechtel, S.; Briem, H.; Marquardt, T.; Schirok, H.; Bohlmann, R.; Nguyen, D.; Fernandez-Montalvan, A.; Boemer, U.; Eberspaecher, U.; Brands, M.; Nussbaum, F.; Koppitz, M.
Deposited on : 2019-12-06
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	2% 52% 27% 7% 14%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dual specificity protein kinase TTK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	252	2059	1325	342	379	1	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	513	GLY	-	expression tag	UNP P33981
A	514	SER	-	expression tag	UNP P33981

- Molecule 2 is (2 {R})-2-(4-fluorophenyl)- {N}-[4-[2-[(2-methoxy-4-methylsulfonyl-phenyl)amino]-[1,2,4]triazolo[1,5-a]pyridin-6-yl]phenyl]propanamide (CCD ID: 8QW) (formula: C₂₉H₂₆FN₅O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
2	A	1	40	29	1	5	4	1	0	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total 24	O 24	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity protein kinase TTK



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	106.93Å 111.67Å 71.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.15 – 2.65 35.15 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.7 (35.15-2.65) 98.7 (35.15-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.212 , 0.262 0.216 , 0.255	Depositor DCC
R_{free} test set	633 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	67.2	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.033 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2124	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, CL, 8QW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	0/2092	1.44	3/2823 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	620	ASP	CB-CA-C	5.46	117.85	109.50
1	A	561	ASP	CA-CB-CG	5.36	117.95	112.60
1	A	608	ASP	CA-CB-CG	5.19	117.79	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2059	0	2076	59	0
2	A	40	0	0	2	0
3	A	1	0	0	0	0
4	A	24	0	0	1	0
All	All	2124	0	2076	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:649:LYS:NZ	1:A:686:TPO:HG23	1.92	0.85
1:A:594:THR:HG22	1:A:596:GLN:H	1.55	0.70
1:A:531:ILE:HD11	1:A:541:GLN:HB2	1.80	0.64
1:A:564:THR:HG22	1:A:669:ASN:HD21	1.63	0.63
1:A:564:THR:HG22	1:A:669:ASN:ND2	2.14	0.62
1:A:764:LEU:HD12	1:A:764:LEU:O	2.01	0.61
1:A:525:TYR:HB3	1:A:542:VAL:HG13	1.83	0.60
1:A:649:LYS:HZ2	1:A:686:TPO:HG23	1.67	0.59
1:A:584:LYS:HE2	1:A:630:ASN:OD1	2.04	0.58
1:A:529:LYS:HE3	1:A:541:GLN:HE21	1.70	0.57
1:A:633:GLU:O	1:A:637:THR:HG23	2.04	0.57
1:A:579:GLN:HE22	1:A:587:ARG:HA	1.70	0.56
1:A:579:GLN:NE2	1:A:588:LEU:H	2.04	0.55
1:A:695:ILE:HA	1:A:698:MET:HE2	1.89	0.55
1:A:653:PHE:HB3	1:A:660:LEU:HD22	1.88	0.54
1:A:584:LYS:HE3	1:A:659:MET:SD	2.49	0.53
1:A:649:LYS:HZ1	1:A:686:TPO:HG23	1.69	0.53
1:A:602:MET:HE1	2:A:901:8QW:C21	2.38	0.52
1:A:525:TYR:HB3	1:A:542:VAL:CG1	2.40	0.51
1:A:771:CYS:O	1:A:779:ARG:HD2	2.11	0.51
1:A:528:LEU:HD11	1:A:543:LEU:HG	1.93	0.51
1:A:609:LEU:HD23	1:A:650:PRO:HB3	1.93	0.50
1:A:761:GLU:HG2	4:A:1012:HOH:O	2.11	0.50
1:A:607:ILE:HD11	1:A:611:SER:HB3	1.95	0.48
1:A:788:HIS:CG	1:A:789:PRO:HD2	2.47	0.48
1:A:531:ILE:HD11	1:A:541:GLN:CB	2.42	0.48
1:A:627:TYR:HE1	1:A:658:GLY:O	1.97	0.48
1:A:659:MET:HE3	1:A:661:LYS:HE2	1.95	0.48
1:A:639:HIS:CE1	1:A:713:PRO:HB2	2.49	0.48
1:A:576:ASN:O	1:A:579:GLN:HG2	2.13	0.48
1:A:602:MET:SD	2:A:901:8QW:C21	3.02	0.47
1:A:528:LEU:HD21	1:A:543:LEU:CD1	2.44	0.47
1:A:639:HIS:CE1	1:A:713:PRO:CB	2.98	0.47
1:A:687:VAL:HG22	1:A:687:VAL:O	2.15	0.46
1:A:554:TYR:HD1	1:A:599:TYR:CE1	2.33	0.46
1:A:649:LYS:CE	1:A:686:TPO:HG23	2.45	0.46
1:A:575:LEU:HD13	1:A:588:LEU:HD22	1.99	0.45
1:A:530:GLN:HE21	1:A:530:GLN:HB2	1.60	0.44
1:A:773:LYS:O	1:A:779:ARG:HD3	2.18	0.44
1:A:579:GLN:HE22	1:A:588:LEU:H	1.66	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:697:ASP:C	1:A:698:MET:CG	2.91	0.43
1:A:625:LYS:HG3	1:A:790:TYR:CZ	2.53	0.43
1:A:620:ASP:HA	1:A:621:PRO:HD3	1.93	0.43
1:A:770:CYS:HA	1:A:773:LYS:HE3	1.99	0.43
1:A:563:GLN:O	1:A:566:ASP:N	2.51	0.42
1:A:628:TRP:CE3	1:A:724:LEU:HD13	2.55	0.42
1:A:586:ILE:HB	1:A:663:ILE:HG22	2.01	0.42
1:A:782:ILE:HB	1:A:783:PRO:HD3	2.02	0.41
1:A:615:LYS:O	1:A:616:LYS:C	2.62	0.41
1:A:724:LEU:O	1:A:728:THR:HG23	2.20	0.41
1:A:780:ILE:HD13	1:A:785:LEU:CD2	2.51	0.41
1:A:586:ILE:O	1:A:586:ILE:HG23	2.20	0.41
1:A:759:ILE:HB	1:A:760:PRO:CD	2.51	0.41
1:A:608:ASP:OD1	1:A:610:ASN:N	2.54	0.41
1:A:736:GLN:HG2	1:A:737:ILE:N	2.35	0.41
1:A:540:PHE:O	1:A:551:ALA:HA	2.21	0.40
1:A:554:TYR:CD1	1:A:599:TYR:CE1	3.08	0.40
1:A:589:TYR:HB2	1:A:601:VAL:HG22	2.02	0.40
1:A:607:ILE:HG12	1:A:608:ASP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/294 (83%)	221 (90%)	23 (9%)	1 (0%)	30	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	656	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	230/269 (86%)	190 (83%)	40 (17%)	2 2

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	520	VAL
1	A	521	LYS
1	A	526	SER
1	A	527	ILE
1	A	530	GLN
1	A	533	SER
1	A	541	GLN
1	A	542	VAL
1	A	545	GLU
1	A	546	LYS
1	A	558	GLU
1	A	562	ASN
1	A	563	GLN
1	A	575	LEU
1	A	584	LYS
1	A	587	ARG
1	A	594	THR
1	A	601	VAL
1	A	608	ASP
1	A	616	LYS
1	A	618	SER
1	A	619	ILE
1	A	637	THR
1	A	648	LEU
1	A	649	LYS
1	A	660	LEU
1	A	669	ASN
1	A	727	MET
1	A	728	THR
1	A	731	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	740	GLN
1	A	741	ILE
1	A	753	GLU
1	A	755	GLU
1	A	759	ILE
1	A	762	LYS
1	A	764	LEU
1	A	777	LYS
1	A	779	ARG
1	A	791	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	541	GLN
1	A	579	GLN
1	A	610	ASN
1	A	652	ASN
1	A	669	ASN
1	A	752	HIS
1	A	765	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	686	1	8,10,11	1.09	1 (12%)	10,14,16	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	686	1	-	5/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	686	TPO	P-OG1	2.73	1.64	1.59

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	686	TPO	N-CA-CB-OG1
1	A	686	TPO	C-CA-CB-CG2
1	A	686	TPO	N-CA-CB-CG2
1	A	686	TPO	CB-OG1-P-O3P
1	A	686	TPO	O-C-CA-CB

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	686	TPO	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	8QW	A	901	-	44,44,44	2.34	11 (25%)	61,64,64	2.16	13 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8QW	A	901	-	-	1/28/28/28	0/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	8QW	N5-N7	-11.00	1.24	1.37
2	A	901	8QW	O29-S27	3.95	1.53	1.44
2	A	901	8QW	C1-N5	3.36	1.44	1.38
2	A	901	8QW	C10-C4	-3.15	1.42	1.48
2	A	901	8QW	C9-N7	3.12	1.36	1.33
2	A	901	8QW	C6-N5	2.78	1.41	1.36
2	A	901	8QW	C12-N11	-2.61	1.36	1.41
2	A	901	8QW	C3-C1	-2.37	1.37	1.41
2	A	901	8QW	O28-S27	2.28	1.49	1.44
2	A	901	8QW	C21-C10	2.14	1.42	1.39
2	A	901	8QW	C38-C36	2.01	1.41	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	8QW	O28-S27-C15	-7.57	102.34	108.23
2	A	901	8QW	C30-S27-C15	6.05	111.22	104.55
2	A	901	8QW	O29-S27-C15	5.05	112.16	108.23
2	A	901	8QW	C9-N7-N5	4.85	105.89	100.67
2	A	901	8QW	O28-S27-O29	-4.37	110.58	117.99
2	A	901	8QW	C6-N5-N7	4.27	130.86	125.20
2	A	901	8QW	O33-C24-C25	-3.99	117.01	122.17
2	A	901	8QW	C25-C24-N23	3.46	119.97	114.84
2	A	901	8QW	N7-C9-N8	-3.25	112.34	116.67
2	A	901	8QW	C39-C25-C26	-3.14	105.04	112.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	8QW	C6-N5-C1	-2.67	119.16	122.53
2	A	901	8QW	C35-C26-C37	2.24	121.08	118.30
2	A	901	8QW	C38-C37-C26	-2.01	119.17	121.18

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	901	8QW	C39-C25-C26-C37

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	8QW	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/294 (85%)	0.10	7 (2%) 55 48	47, 82, 132, 154	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	794	GLN	3.9
1	A	619	ILE	2.6
1	A	741	ILE	2.5
1	A	711	ILE	2.4
1	A	622	TRP	2.4
1	A	790	TYR	2.2
1	A	737	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	686	11/12	0.82	0.12	136,148,176,176	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	8QW	A	901	40/40	0.93	0.09	47,66,94,100	0
3	CL	A	902	1/1	0.94	0.15	94,94,94,94	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.