



Full wwPDB EM Validation Report ⓘ

Mar 28, 2026 – 03:26 AM UTC

PDB ID : 7TOS / pdb_00007tos
EMDB ID : EMD-26037
Title : E. coli 70S ribosomes bound with the ALS/FTD-associated dipeptide repeat protein PR20
Authors : Loveland, A.B.; Svidritskiy, E.; Susorov, D.; Lee, S.; Park, A.; Zvornicanin, S.; Demo, G.; Gao, F.B.; Korostelev, A.A.
Deposited on : 2022-01-24
Resolution : 2.90 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : **FAILED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

i

ELECTRON MICROSCOPY

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Q-score	25397	13054 (2.40 - 3.40)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 56 unique types of molecules in this entry. The entry contains 145427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	L02	271	Total	C	N	O	S	0	0
			2083	1288	423	365	7		

- Molecule 2 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	L03	209	Total	C	N	O	S	0	0
			1565	979	288	294	4		

- Molecule 3 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	L04	201	Total	C	N	O	S	0	0
			1552	974	283	290	5		

- Molecule 4 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	L05	177	Total	C	N	O	S	0	0
			1411	899	249	257	6		

- Molecule 5 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	L06	176	Total	C	N	O	S	0	0
			1323	832	243	246	2		

- Molecule 6 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	L09	149	Total	C	N	O	S	0	0
			1111	699	197	214	1		

- Molecule 7 is a protein called 50S ribosomal protein L10.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	L10	131	Total	C	N	O	S	0	0
			989	625	175	184	5		

- Molecule 8 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L11	141	Total	C	N	O	S	0	0
			1032	651	179	196	6		

- Molecule 9 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	L13	142	Total	C	N	O	S	0	0
			1129	714	212	199	4		

- Molecule 10 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	L14	122	Total	C	N	O	S	0	0
			939	587	180	166	6		

- Molecule 11 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	L15	143	Total	C	N	O	S	0	0
			1045	649	206	189	1		

- Molecule 12 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L16	136	Total	C	N	O	S	0	0
			1074	686	205	177	6		

- Molecule 13 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	L17	120	Total	C	N	O	S	0	0
			961	593	196	167	5		

- Molecule 14 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	L18	116	Total	C	N	O	0	0
			892	552	178	162		

- Molecule 15 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	L19	114	Total	C	N	O	S	0	0
			917	574	179	163	1		

- Molecule 16 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms				AltConf	Trace
16	L20	117	Total	C	N	O	0	0
			947	604	192	151		

- Molecule 17 is a protein called Ribosomal protein L21.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	L21	103	Total	C	N	O	S	0	0
			816	516	153	145	2		

- Molecule 18 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	L22	110	Total	C	N	O	S	0	0
			857	532	166	156	3		

- Molecule 19 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	L23	93	Total	C	N	O	S	0	0
			739	466	139	132	2		

- Molecule 20 is a protein called Ribosomal protein L24.

Mol	Chain	Residues	Atoms				AltConf	Trace
20	L24	102	Total	C	N	O	0	0
			780	492	146	142		

- Molecule 21 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	L25	94	Total	C	N	O	S	0	0
			753	479	137	134	3		

- Molecule 22 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	L27	75	Total	C	N	O	S	0	0
			575	356	116	102	1		

- Molecule 23 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	L28	77	Total	C	N	O	S	0	0
			625	388	129	106	2		

- Molecule 24 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	L29	63	Total	C	N	O	S	0	0
			509	313	99	95	2		

- Molecule 25 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	L30	58	Total	C	N	O	S	0	0
			449	281	87	79	2		

- Molecule 26 is a protein called 50S ribosomal protein L31.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	L31	66	Total	C	N	O	S	0	0
			523	323	99	95	6		

- Molecule 27 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	L32	56	Total	C	N	O	S	0	0
			444	269	94	80	1		

- Molecule 28 is a protein called 50S ribosomal protein L33.

Mol	Chain	Residues	Atoms				AltConf	Trace
28	L33	50	Total	C	N	O	0	0
			410	263	75	72		

- Molecule 29 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	L34	46	Total	C	N	O	S	0	0
			377	228	90	57	2		

- Molecule 30 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	L35	64	Total	C	N	O	S	0	0
			504	323	105	74	2		

- Molecule 31 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	L36	38	Total	C	N	O	S	0	0
			302	185	65	48	4		

- Molecule 32 is a protein called 30S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	S02	225	Total	C	N	O	S	0	0
			1757	1111	315	323	8		

- Molecule 33 is a protein called 30S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	S03	206	Total	C	N	O	S	0	0
			1625	1028	305	289	3		

- Molecule 34 is a protein called 30S ribosomal protein S4.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	S04	205	Total	C	N	O	S	0	0
			1643	1026	315	298	4		

- Molecule 35 is a protein called 30S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	S05	157	Total	C	N	O	S	0	0
			1157	719	218	214	6		

- Molecule 36 is a protein called 30S ribosomal protein S6, non-modified isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	S06	100	Total	C	N	O	S	0	0
			818	515	148	149	6		

- Molecule 37 is a protein called 30S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
37	S07	151	Total	C	N	O	S	0	0
			1182	735	227	216	4		

- Molecule 38 is a protein called 30S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
38	S08	129	Total	C	N	O	S	0	0
			979	616	173	184	6		

- Molecule 39 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
39	S09	127	Total	C	N	O	S	0	0
			1022	634	206	179	3		

- Molecule 40 is a protein called 30S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
40	S10	98	Total	C	N	O	S	0	0
			787	493	150	143	1		

- Molecule 41 is a protein called 30S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
41	S11	116	Total	C	N	O	S	0	0
			869	534	173	159	3		

- Molecule 42 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues	Atoms					AltConf	Trace
42	S12	123	Total	C	N	O	S	0	0
			955	590	196	165	4		

- Molecule 43 is a protein called 30S ribosomal protein S13.

Mol	Chain	Residues	Atoms					AltConf	Trace
43	S13	114	Total	C	N	O	S	0	0
			884	546	178	157	3		

- Molecule 44 is a protein called 30S ribosomal protein S14.

Mol	Chain	Residues	Atoms					AltConf	Trace
44	S14	100	Total	C	N	O	S	0	0
			805	499	164	139	3		

- Molecule 45 is a protein called 30S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
45	S15	88	Total	C	N	O	S	0	0
			714	439	144	130	1		

- Molecule 46 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
46	S16	82	Total	C	N	O	S	0	0
			649	406	128	114	1		

- Molecule 47 is a protein called 30S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
47	S17	80	Total	C	N	O	S	0	0
			649	411	121	114	3		

- Molecule 48 is a protein called 30S ribosomal protein S18.

Mol	Chain	Residues	Atoms					AltConf	Trace
48	S18	65	Total	C	N	O	S	0	0
			536	339	100	96	1		

- Molecule 49 is a protein called 30S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
49	S19	79	Total	C	N	O	S	0	0
			638	408	120	108	2		

- Molecule 50 is a protein called 30S ribosomal protein S20.

Mol	Chain	Residues	Atoms					AltConf	Trace
50	S20	85	Total	C	N	O	S	0	0
			665	411	137	114	3		

- Molecule 51 is a protein called Ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
51	S21	65	Total	C	N	O	S	0	0
			545	335	117	92	1		

- Molecule 52 is a protein called 50S ribosomal protein L1.

Mol	Chain	Residues	Atoms					AltConf	Trace
52	L1	134	Total	C	N	O	S	0	0
			1027	645	186	194	2		

- Molecule 53 is a RNA chain called 16S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
53	16S	1532	Total	C	N	O	P	0	0
			32870	14661	6031	10647	1531		

- Molecule 54 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
54	23S	2903	Total	C	N	O	P	0	0
			62317	27801	11468	20146	2902		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
23S	747	C	U	variant	GB 1036415628

- Molecule 55 is a RNA chain called 5S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
55	5S	120	Total	C	N	O	P	0	0
			2568	1145	471	833	119		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
5S	120	A	-	insertion	GB 1370526515

- Molecule 56 is a protein called PR20, ALS/FTD dipeptide repeat protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
56	PR	15	Total	C	N	O	0	0
			133	82	36	15		

MolProbity failed to run properly - this section is therefore empty.

3 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	49219	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	35	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	26.145	Depositor
Minimum map value	-6.229	Depositor
Average map value	0.018	Depositor
Map value standard deviation	2.086	Depositor
Recommended contour level	3	Depositor
Map size ($\text{\AA}$)	375.84, 375.84, 375.84	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.87, 0.87, 0.87	Depositor

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-26037. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

5.1 Orthogonal projections [i](#)


5.1.1 Primary map



X




Y




Z


5.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

5.2 Central slices [i](#)


5.2.1 Primary map



X Index: 216




Y Index: 216




Z Index: 216


5.2.2 Raw map



X Index: 216



Y Index: 216




Z Index: 216


The images above show central slices of the map in three orthogonal directions.

5.3 Largest variance slices [i](#)


5.3.1 Primary map



X Index: 225




Y Index: 223




Z Index: 202


5.3.2 Raw map



X Index: 224



Y Index: 223




Z Index: 196


The images above show the largest variance slices of the map in three orthogonal directions.

5.4 Orthogonal standard-deviation projections (False-color) [i](#)


5.4.1 Primary map



X




Y




Z


5.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


5.5 Orthogonal surface views [i](#)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

5.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


6.2 Volume estimate [i](#)



The volume at the recommended contour level is 2732 nm³; this corresponds to an approximate mass of 2468 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

6.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.345 $\text{\AA}^{-1}$

7 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

7.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹

7.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.90	-	-
Author-provided FSC curve	2.93	3.18	2.96
Unmasked-calculated*	3.18	3.82	3.22

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-26037 and PDB model 7TOS. Per-residue inclusion information can be found in section ?? on page ??.


8.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3).

8.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9680	 0.4280
16S	 0.9930	 0.4370
23S	 0.9940	 0.4500
5S	 0.9970	 0.4130
L02	 0.9900	 0.5030
L03	 0.9820	 0.4860
L04	 0.9340	 0.4240
L05	 0.9710	 0.3380
L06	 0.9700	 0.3980
L09	 0.3800	 0.2900
L1	 0.4910	 0.0640
L10	 0.4360	 0.0890
L11	 0.7170	 0.0450
L13	 0.9740	 0.4680
L14	 0.9660	 0.4790
L15	 0.9620	 0.4500
L16	 0.9650	 0.4730
L17	 0.9940	 0.4640
L18	 0.9750	 0.4010
L19	 0.9730	 0.4690
L20	 0.9860	 0.4730
L21	 0.9420	 0.4560
L22	 0.9550	 0.4590
L23	 0.9710	 0.4360
L24	 0.9490	 0.4020
L25	 0.9470	 0.4300
L27	 0.9700	 0.4890
L28	 0.9880	 0.4710
L29	 0.9620	 0.3560
L30	 0.9570	 0.4540
L31	 0.9080	 0.2990
L32	 0.9670	 0.4650
L33	 0.7260	 0.4150
L34	 0.9780	 0.4840
L35	 0.9940	 0.5000



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
L36	 1.0000	 0.4610
PR	 0.9500	 0.4130
S02	 0.9080	 0.3740
S03	 0.8900	 0.4010
S04	 0.9420	 0.3900
S05	 0.9570	 0.4350
S06	 0.9500	 0.3820
S07	 0.9590	 0.3370
S08	 0.9670	 0.4500
S09	 0.9530	 0.3410
S10	 0.8150	 0.3450
S11	 0.9910	 0.4240
S12	 0.9510	 0.4510
S13	 0.9590	 0.3260
S14	 0.9430	 0.3740
S15	 0.9830	 0.4080
S16	 0.9510	 0.4210
S17	 0.9720	 0.4260
S18	 0.9940	 0.4170
S19	 0.9520	 0.3490
S20	 0.9750	 0.3920
S21	 0.9050	 0.2590