



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:44 AM UTC

PDB ID : 1TPY / pdb_00001tpy
Title : Structure of the cyclopropane synthase MmaA2 from Mycobacterium tuberculosis
Authors : Smith, C.V.; Sacchettini, J.C.
Deposited on : 2004-06-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	287	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called methoxy mycolic acid synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	2292	1463	397	422	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q79FX6

- Molecule 2 is CARBONATE ION (CCD ID: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	1	3	0	0

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 4 is CETYL-TRIMETHYL-AMMONIUM (CCD ID: 16A) (formula: $C_{19}H_{42}N$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			20	19	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	192	Total 192	O 192	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: methoxy mycolic acid synthase 2



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.59Å 58.42Å 133.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20	Depositor
% Data completeness (in resolution range)	93.6 (50.00-2.20)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	0.04	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.194 , 0.232	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2534	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, SAH, 16A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2348	0.85	4/3183 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	GLY	N-CA-C	7.49	125.33	115.36
1	A	207	THR	N-CA-C	-6.67	100.59	110.46
1	A	89	ASN	N-CA-C	-5.65	102.93	110.55
1	A	173	THR	N-CA-C	-5.20	104.02	110.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2292	0	2241	56	0
2	A	4	0	0	0	0
3	A	26	0	19	2	0
4	A	20	0	42	3	0
5	A	192	0	0	4	0
All	All	2534	0	2302	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:LEU:HD13	1:A:124:GLU:HB2	1.63	0.80
1:A:207:THR:HG22	1:A:210:MET:HG3	1.67	0.74
1:A:207:THR:HG22	1:A:210:MET:CG	2.19	0.72
1:A:198:GLU:HB3	1:A:261:ARG:HD2	1.77	0.66
1:A:236:LEU:HD21	4:A:1901:16A:H13	1.76	0.66
1:A:225:ARG:HH11	1:A:281:ASN:ND2	1.94	0.65
1:A:3:ASN:OD1	1:A:4:ASP:N	2.25	0.64
1:A:144:HIS:H	1:A:204:GLN:NE2	1.95	0.64
1:A:49:GLN:O	1:A:53:ILE:HG12	1.99	0.63
1:A:4:ASP:CG	1:A:5:LEU:H	2.06	0.62
1:A:43:MET:HG2	1:A:47:GLU:HB2	1.80	0.62
1:A:85:GLN:HG2	5:A:1952:HOH:O	2.01	0.60
1:A:207:THR:CG2	1:A:210:MET:HG3	2.33	0.59
1:A:7:PRO:HB2	1:A:9:PHE:CE1	2.39	0.57
1:A:165:LEU:C	1:A:165:LEU:HD13	2.29	0.57
1:A:35:CYS:HB2	1:A:232:TYR:CD2	2.39	0.56
1:A:207:THR:HG23	1:A:210:MET:H	1.70	0.56
1:A:207:THR:HG21	5:A:1929:HOH:O	2.05	0.55
1:A:33:TYR:O	4:A:1901:16A:H171	2.08	0.54
1:A:117:ARG:NH1	1:A:128:GLU:OE1	2.33	0.54
1:A:73:CYS:HB2	1:A:79:MET:HE2	1.91	0.51
1:A:114:ARG:HA	1:A:114:ARG:HE	1.75	0.51
1:A:35:CYS:HB2	1:A:232:TYR:CE2	2.46	0.50
1:A:7:PRO:HG3	1:A:124:GLU:HG2	1.95	0.49
1:A:173:THR:OG1	1:A:176:GLN:HG3	2.12	0.49
1:A:191:PHE:CZ	1:A:268:GLY:HA3	2.48	0.49
1:A:4:ASP:O	1:A:5:LEU:HD12	2.14	0.48
1:A:126:PHE:CZ	1:A:128:GLU:HB2	2.49	0.48
1:A:72:GLY:HA2	3:A:1900:SAH:H1'	1.96	0.48
1:A:52:LYS:C	1:A:52:LYS:HD2	2.39	0.48
1:A:214:GLN:HE21	1:A:214:GLN:HA	1.78	0.48
1:A:130:VAL:O	1:A:159:PRO:HD3	2.15	0.47
3:A:1900:SAH:SD	4:A:1901:16A:H173	2.54	0.47
1:A:155:HIS:CE1	1:A:287:LYS:HE3	2.50	0.47
1:A:265:TYR:C	1:A:265:TYR:CD1	2.93	0.47
1:A:167:HIS:C	1:A:167:HIS:CD2	2.93	0.47
1:A:12:VAL:HG21	1:A:95:LEU:HD12	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:ILE:HG23	1:A:167:HIS:CG	2.50	0.46
1:A:40:ARG:O	1:A:43:MET:HB2	2.16	0.45
1:A:87:ASP:OD1	1:A:114:ARG:HD2	2.15	0.45
1:A:4:ASP:CG	1:A:5:LEU:N	2.73	0.45
1:A:53:ILE:HD12	1:A:78:THR:OG1	2.16	0.45
1:A:5:LEU:HD12	1:A:5:LEU:O	2.17	0.44
1:A:62:LEU:HD23	1:A:132:ARG:HG3	1.98	0.44
1:A:57:LEU:HD21	1:A:82:ALA:HB2	1.99	0.44
1:A:159:PRO:O	1:A:287:LYS:HD3	2.18	0.43
1:A:49:GLN:HA	1:A:49:GLN:OE1	2.19	0.43
1:A:59:LYS:NZ	1:A:282:GLN:HE21	2.17	0.43
1:A:144:HIS:CE1	1:A:210:MET:HE1	2.54	0.43
1:A:174:ARG:NH2	5:A:1978:HOH:O	2.51	0.42
1:A:18:LEU:HD11	1:A:254:GLN:NE2	2.34	0.42
1:A:198:GLU:OE2	1:A:261:ARG:HD3	2.19	0.42
1:A:213:GLU:O	1:A:217:LYS:HG3	2.19	0.42
1:A:225:ARG:HH11	1:A:281:ASN:HD21	1.66	0.41
1:A:121:ALA:HB1	1:A:125:GLN:HE21	1.85	0.41
1:A:249:GLU:OE2	1:A:249:GLU:N	2.46	0.41
1:A:111:ASP:C	5:A:2012:HOH:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	242/244 (99%)	235 (97%)	7 (3%)	37 51

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LEU
1	A	114	ARG
1	A	166	LEU
1	A	182	LEU
1	A	192	LEU
1	A	232	TYR
1	A	265	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	37	HIS
1	A	99	GLN
1	A	125	GLN
1	A	204	GLN
1	A	214	GLN
1	A	254	GLN
1	A	281	ASN
1	A	282	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	16A	A	1901	-	19,19,19	1.32	1 (5%)	21,21,21	1.55	4 (19%)
2	CO3	A	902	-	3,3,3	4.01	1 (33%)	2,3,3	0.14	0
3	SAH	A	1900	-	27,28,28	0.94	2 (7%)	36,40,40	0.83	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	16A	A	1901	-	-	2/17/17/17	-
3	SAH	A	1900	-	-	0/15/31/31	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	902	CO3	O1-C	6.94	1.49	1.25
4	A	1901	16A	C17-N1	-5.24	1.34	1.50
3	A	1900	SAH	C1'-N9	2.29	1.52	1.46
3	A	1900	SAH	C4-N3	2.22	1.38	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1901	16A	C17-N1-C16	4.24	126.78	109.91
4	A	1901	16A	C19-N1-C17	-3.04	101.00	108.98
4	A	1901	16A	C15-C16-N1	3.03	127.07	115.97
4	A	1901	16A	C18-N1-C17	-2.19	103.21	108.98
3	A	1900	SAH	C1'-N9-C8	2.02	131.58	127.09

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1901	16A	C10-C11-C12-C13

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1901	16A	C6-C7-C8-C9


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1901	16A	3	0
3	A	1900	SAH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.