



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:05 PM UTC

PDB ID : 5TPW / pdb_00005tpw
Title : Crystal structure of amino terminal domains of the NMDA receptor subunit GluN1 and GluN2A in complex with zinc at the GluN2A
Authors : Romero-Hernandez, A.; Simorowski, N.; Karakas, E.; Furukawa, H.
Deposited on : 2016-10-21
Resolution : 2.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	3% 81% 11% 7%
2	B	360	4% 78% 13% 8%
3	H	221	% 79% 15% 5%
4	L	214	2% 86% 10% .

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 8358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NMDA glutamate receptor subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	0	0
			2734	1745	467	511	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLN	ASN	engineered mutation	UNP Q91977
A	371	GLN	ASN	engineered mutation	UNP Q91977
A	409	GLY	-	expression tag	UNP Q91977
A	410	THR	-	expression tag	UNP Q91977
A	411	LEU	-	expression tag	UNP Q91977
A	412	VAL	-	expression tag	UNP Q91977

- Molecule 2 is a protein called Glutamate receptor ionotropic, NMDA 2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	331	Total	C	N	O	S	0	0	0
			2472	1586	405	469	12			

- Molecule 3 is a protein called FAB, HEAVY CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	211	Total	C	N	O	S	0	0	0
			1542	993	244	298	7			

- Molecule 4 is a protein called FAB, LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	L	206	Total	C	N	O	S	0	0	0
			1500	945	258	290	7			


- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Zn	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	19	Total 19	O 19	0	0
9	B	7	Total 7	O 7	0	0
9	H	5	Total 5	O 5	0	0
9	L	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NMDA glutamate receptor subunit




- Molecule 2: Glutamate receptor ionotropic, NMDA 2A



- Molecule 3: FAB, HEAVY CHAIN



● Molecule 4: FAB, LIGHT CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.56Å 118.40Å 155.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.56 – 2.91 47.56 – 2.91	Depositor EDS
% Data completeness (in resolution range)	88.5 (47.56-2.91) 94.4 (47.56-2.91)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.200 , 0.245 (Not available) , 0.255	Depositor DCC
R_{free} test set	3180 reflections (9.45%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8358	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, GOL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2787	0.33	0/3791
2	B	0.11	0/2527	0.28	0/3447
3	H	0.13	0/1590	0.33	0/2188
4	L	0.13	0/1536	0.37	2/2096 (0.1%)
All	All	0.12	0/8440	0.32	2/11522 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	H	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	66	GLY	CA-C-N	-5.18	111.65	121.54
4	L	66	GLY	C-N-CA	-5.18	111.65	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	H	148	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2734	0	2683	29	1
2	B	2472	0	2285	23	1
3	H	1542	0	1376	16	0
4	L	1500	0	1341	12	0
5	A	28	0	26	2	0
6	A	15	0	0	1	0
6	B	15	0	0	1	0
7	A	12	0	16	3	0
8	B	1	0	0	0	0
9	A	19	0	0	1	0
9	B	7	0	0	0	0
9	H	5	0	0	0	0
9	L	8	0	0	0	0
All	All	8358	0	7727	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:71:SER:HB3	3:H:80:TYR:HB2	1.64	0.79
3:H:149:LYS:O	3:H:182:THR:OG1	2.05	0.71
2:B:188:LYS:HG3	2:B:200:MET:HE1	1.74	0.69
2:B:174:THR:HG22	2:B:230:TYR:HB3	1.75	0.67
2:B:159:LEU:HB3	2:B:196:VAL:HG11	1.78	0.64
3:H:12:LYS:HG3	3:H:18:VAL:HG22	1.80	0.63
1:A:38:HIS:NE2	6:A:504:SO4:O1	2.27	0.63
3:H:149:LYS:HA	3:H:182:THR:HG23	1.82	0.61
2:B:167:TRP:HB3	2:B:226:VAL:HG21	1.82	0.60
1:A:389:ASN:OD1	5:A:502:NAG:N2	2.35	0.60
1:A:67:HIS:CE1	1:A:93:SER:O	2.55	0.59
3:H:91:THR:HG23	3:H:116:THR:HA	1.85	0.58
1:A:167:VAL:HG12	1:A:243:SER:HB3	1.84	0.58
1:A:262:THR:HG23	1:A:284:PRO:HB3	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:ASN:HB2	1:A:394:ILE:HG13	1.85	0.58
3:H:38:LYS:NZ	3:H:94:TYR:OH	2.36	0.57
1:A:216:LEU:HB3	1:A:228:LEU:HD11	1.88	0.56
1:A:297:ASN:OD1	5:A:501:NAG:H83	2.05	0.56
3:H:161:ASN:HB3	3:H:164:SER:HB3	1.87	0.56
2:B:134:ILE:HG12	2:B:355:TYR:HB3	1.88	0.55
4:L:61:ARG:NH2	4:L:82:ASP:OD1	2.40	0.55
2:B:308:GLU:OE1	2:B:308:GLU:N	2.40	0.54
1:A:68:ARG:NH2	9:A:603:HOH:O	2.38	0.54
1:A:36:LYS:H	7:A:506:GOL:H32	1.72	0.53
1:A:275:ILE:HD13	1:A:289:GLY:HA3	1.90	0.52
2:B:89:LEU:HB3	2:B:95:ILE:HD12	1.92	0.52
1:A:36:LYS:HE2	3:H:103:GLU:HB2	1.91	0.51
2:B:74:MET:HE2	2:B:76:ARG:HD2	1.93	0.51
3:H:129:PRO:O	4:L:121:SER:OG	2.23	0.50
1:A:35:THR:HB	7:A:506:GOL:H32	1.93	0.49
1:A:129:SER:OG	1:A:139:ARG:NH2	2.46	0.49
3:H:154:GLU:HG2	3:H:181:TYR:CZ	2.48	0.49
2:B:139:ASP:OD1	2:B:141:THR:OG1	2.25	0.48
1:A:363:GLU:H	1:A:363:GLU:CD	2.21	0.48
3:H:202:ASN:ND2	3:H:213:ASP:OD2	2.40	0.48
4:L:136:LEU:HD13	4:L:175:MET:HE3	1.95	0.48
2:B:277:ILE:HG12	2:B:365:VAL:HG12	1.95	0.48
3:H:55:THR:O	3:H:57:GLU:N	2.45	0.48
3:H:130:LEU:HD21	3:H:147:LEU:HB2	1.95	0.47
3:H:97:ALA:HB1	3:H:106:PHE:HB3	1.97	0.47
4:L:37:GLN:HB2	4:L:47:LEU:HD11	1.97	0.46
4:L:149:LYS:HB2	4:L:193:THR:OG1	2.15	0.46
2:B:298:ILE:HA	2:B:341:VAL:HG11	1.96	0.46
1:A:86:GLN:HB3	1:A:325:PRO:HG2	1.97	0.46
4:L:33:LEU:HG	4:L:71:PHE:CD1	2.49	0.46
1:A:287:ILE:HG22	1:A:377:LEU:HB3	1.98	0.46
2:B:333:THR:OG1	6:B:404:SO4:O3	2.34	0.45
1:A:67:HIS:HE1	1:A:93:SER:O	2.00	0.45
1:A:34:SER:HB3	1:A:94:HIS:O	2.16	0.44
4:L:175:MET:HE2	4:L:175:MET:HB3	1.73	0.44
3:H:13:LYS:N	3:H:16:GLU:OE1	2.45	0.44
4:L:83:VAL:HG11	4:L:166:GLN:HB3	2.00	0.44
2:B:307:LEU:HD23	2:B:307:LEU:HA	1.75	0.44
2:B:206:LEU:HD13	2:B:239:ILE:HD12	2.00	0.44
2:B:285:ASP:OD2	2:B:374:LYS:NZ	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:PRO:HA	1:A:358:ARG:HA	1.99	0.43
2:B:158:MET:HE3	2:B:256:ILE:HG22	1.99	0.43
1:A:298:GLU:O	1:A:298:GLU:HG2	2.18	0.43
1:A:264:ALA:HA	1:A:403:PRO:O	2.19	0.43
2:B:229:LEU:HB3	2:B:257:VAL:HG12	2.00	0.42
4:L:86:TYR:HB2	4:L:102:THR:HG23	2.01	0.42
1:A:27:VAL:HG13	1:A:88:TYR:CD1	2.54	0.42
1:A:36:LYS:HG3	7:A:506:GOL:H12	2.00	0.42
2:B:37:ALA:HB2	2:B:95:ILE:HD13	2.01	0.42
4:L:32:TYR:HB2	4:L:92:HIS:HB2	2.00	0.42
1:A:113:PHE:CZ	2:B:106:GLN:HG2	2.54	0.42
1:A:103:THR:O	1:A:106:PRO:HD2	2.20	0.42
2:B:93:ALA:O	2:B:95:ILE:HG13	2.20	0.42
2:B:363:VAL:HB	2:B:376:GLY:HA3	2.02	0.42
2:B:204:ILE:HG23	2:B:218:GLN:HG3	2.02	0.41
2:B:145:PHE:CE2	2:B:338:MET:HE3	2.55	0.41
1:A:268:TRP:HB3	1:A:287:ILE:HG13	2.03	0.41
4:L:31:ASP:OD1	4:L:31:ASP:N	2.53	0.41
1:A:241:ILE:HA	1:A:269:LEU:O	2.21	0.41
1:A:402:TRP:HE1	1:A:408:GLU:CB	2.34	0.40
4:L:39:LYS:HB3	4:L:40:SER:H	1.79	0.40
3:H:56:GLY:O	3:H:58:PRO:HD3	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:406:GLU:OE1	2:B:225:SER:OG[3_454]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/389 (91%)	338 (96%)	16 (4%)	0	100	100
2	B	325/360 (90%)	311 (96%)	14 (4%)	0	100	100
3	H	207/221 (94%)	193 (93%)	13 (6%)	1 (0%)	24	54
4	L	202/214 (94%)	192 (95%)	10 (5%)	0	100	100
All	All	1088/1184 (92%)	1034 (95%)	53 (5%)	1 (0%)	48	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	H	56	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/334 (85%)	281 (99%)	4 (1%)	59	85
2	B	251/318 (79%)	241 (96%)	10 (4%)	28	62
3	H	151/188 (80%)	143 (95%)	8 (5%)	20	52
4	L	146/188 (78%)	143 (98%)	3 (2%)	47	77
All	All	833/1028 (81%)	808 (97%)	25 (3%)	36	70

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	270	VAL
1	A	338	THR
1	A	406	GLU
1	A	412	VAL
2	B	48	GLU
2	B	72	LEU
2	B	127	ILE
2	B	134	ILE
2	B	210	PHE
2	B	229	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	312	TYR
2	B	339	VAL
2	B	362	VAL
2	B	386	ARG
3	H	3	LYS
3	H	45	LEU
3	H	122	THR
3	H	143	THR
3	H	154	GLU
3	H	158	VAL
3	H	176	LEU
3	H	177	GLN
4	L	43	SER
4	L	78	VAL
4	L	102	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	HIS
1	A	73	GLN
1	A	161	ASN
1	A	217	GLN
2	B	356	GLN
3	H	39	GLN
4	L	38	GLN
4	L	92	HIS
4	L	156	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	502	1	14,14,15	0.43	0	17,19,21	0.49	0
6	SO4	A	503	-	4,4,4	0.24	0	6,6,6	0.08	0
5	NAG	A	501	1	14,14,15	0.78	1 (7%)	17,19,21	1.24	1 (5%)
6	SO4	B	404	-	4,4,4	0.24	0	6,6,6	0.09	0
6	SO4	B	403	-	4,4,4	0.23	0	6,6,6	0.07	0
6	SO4	A	504	-	4,4,4	0.37	0	6,6,6	0.20	0
7	GOL	A	507	-	5,5,5	0.37	0	5,5,5	0.23	0
6	SO4	A	505	-	4,4,4	0.24	0	6,6,6	0.09	0
7	GOL	A	506	-	5,5,5	0.38	0	5,5,5	0.34	0
6	SO4	B	402	-	4,4,4	0.24	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	502	1	-	2/6/23/26	0/1/1/1
5	NAG	A	501	1	-	4/6/23/26	0/1/1/1
7	GOL	A	506	-	-	2/4/4/4	-
7	GOL	A	507	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	501	NAG	C1-C2	2.75	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	501	NAG	C1-O5-C5	4.37	118.04	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	506	GOL	O1-C1-C2-C3
5	A	502	NAG	O5-C5-C6-O6
5	A	501	NAG	O5-C5-C6-O6
5	A	501	NAG	C4-C5-C6-O6
5	A	502	NAG	C4-C5-C6-O6
5	A	501	NAG	C8-C7-N2-C2
5	A	501	NAG	O7-C7-N2-C2
7	A	506	GOL	O1-C1-C2-O2
7	A	507	GOL	O1-C1-C2-O2
7	A	507	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	502	NAG	1	0
5	A	501	NAG	1	0
6	B	404	SO4	1	0
6	A	504	SO4	1	0
7	A	506	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/389 (93%)	-0.09	12 (3%) 49 40	12, 34, 69, 98	0
2	B	331/360 (91%)	0.49	14 (4%) 40 32	22, 56, 89, 104	0
3	H	211/221 (95%)	0.31	3 (1%) 73 65	24, 52, 78, 94	0
4	L	206/214 (96%)	0.32	4 (1%) 66 58	23, 51, 80, 100	0
All	All	1110/1184 (93%)	0.23	33 (2%) 52 43	12, 48, 82, 104	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	196	VAL	4.0
4	L	59	PRO	3.9
1	A	408	GLU	3.4
2	B	260	LEU	3.2
3	H	149	LYS	3.2
2	B	194	SER	3.2
1	A	411	LEU	3.1
2	B	380	ASN	3.1
1	A	53	HIS	3.1
2	B	195	PHE	3.0
2	B	360	ARG	3.0
4	L	52	SER	2.9
1	A	412	VAL	2.9
2	B	334	LEU	2.7
1	A	237	ALA	2.7
2	B	337	PHE	2.7
1	A	258	MET	2.6
1	A	24	PRO	2.6
2	B	381	GLN	2.5
1	A	93	SER	2.5
2	B	286	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	410	THR	2.4
2	B	48	GLU	2.4
3	H	50	TRP	2.4
2	B	34	LEU	2.3
1	A	57	LYS	2.3
4	L	109	ALA	2.2
2	B	167	TRP	2.2
1	A	236	GLU	2.2
4	L	51	ALA	2.1
2	B	280	SER	2.1
3	H	75	SER	2.0
1	A	396	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	502	14/15	0.72	0.16	70,86,101,103	0
6	SO4	B	404	5/5	0.75	0.16	83,85,103,106	0
7	GOL	A	507	6/6	0.82	0.15	43,57,77,88	0
5	NAG	A	501	14/15	0.84	0.15	63,77,93,93	0
6	SO4	A	505	5/5	0.84	0.10	65,74,100,106	0
7	GOL	A	506	6/6	0.88	0.11	41,55,60,63	0
6	SO4	B	402	5/5	0.92	0.11	39,71,78,83	0
8	ZN	B	401	1/1	0.92	0.10	58,58,58,58	1
6	SO4	B	403	5/5	0.93	0.09	44,46,56,67	0
6	SO4	A	503	5/5	0.95	0.08	61,67,71,81	0
6	SO4	A	504	5/5	0.97	0.09	36,39,60,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.