



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:19 PM UTC

PDB ID : 7TP4 / pdb_00007tp4
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with neutralizing antibody K398.22
Authors : Yuan, M.; Zhu, X.; Wilson, I.A.
Deposited on : 2022-01-24
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Z	205	16% 81% 14% 5%
2	H	227	7% 87% 9% .
3	L	214	4% 92% 7% .
4	A	3	100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Z	194	Total	C	N	O	S	0	0	0
			1536	985	256	287	8			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Z	531	GLY	-	expression tag	UNP P0DTC2
Z	532	HIS	-	expression tag	UNP P0DTC2
Z	533	HIS	-	expression tag	UNP P0DTC2
Z	534	HIS	-	expression tag	UNP P0DTC2
Z	535	HIS	-	expression tag	UNP P0DTC2
Z	536	HIS	-	expression tag	UNP P0DTC2
Z	537	HIS	-	expression tag	UNP P0DTC2


- Molecule 2 is a protein called K398.22 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	219	Total	C	N	O	S	0	0	0
			1649	1048	275	320	6			

- Molecule 3 is a protein called K398.22 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	211	Total	C	N	O	S	0	0	0
			1558	970	261	321	6			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	A	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	Z	31	Total	O	0	0
			31	31		
6	H	138	Total	O	0	0
			138	138		
6	L	136	Total	O	0	0
			136	136		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 2: K398.22 heavy chain



- Molecule 3: K398.22 light chain



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	68.64Å 277.27Å 91.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.21 – 1.95 46.21 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.21-1.95) 99.7 (46.21-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.205 , 0.231 0.219 , 0.239	Depositor DCC
R_{free} test set	3304 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtriage
Anisotropy	0.643	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5090	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	Z	0.39	0/1580	0.53	0/2151
2	H	0.63	1/1688 (0.1%)	0.77	3/2297 (0.1%)
3	L	0.56	0/1595	0.71	0/2177
All	All	0.54	1/4863 (0.0%)	0.68	3/6625 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	221	ARG	CD-NE	-5.17	1.39	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	220	LYS	CA-C-N	-5.70	112.32	122.07
2	H	220	LYS	C-N-CA	-5.70	112.32	122.07
2	H	221	ARG	CB-CG-CD	-5.65	98.31	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Z	1536	0	1452	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1649	0	1618	18	0
3	L	1558	0	1501	14	0
4	A	38	0	34	2	0
5	L	4	0	6	2	0
6	H	138	0	0	2	1
6	L	136	0	0	4	1
6	Z	31	0	0	5	0
All	All	5090	0	4611	53	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:214:SER:O	2:H:216:THR:N	2.00	0.95
1:Z:358:ILE:HB	1:Z:395:VAL:HG23	1.65	0.78
3:L:168:LYS:NZ	6:L:401:HOH:O	2.16	0.74
2:H:168:GLY:O	6:H:301:HOH:O	2.06	0.73
1:Z:375:SER:OG	6:Z:601:HOH:O	2.10	0.69
1:Z:514:SER:OG	6:Z:602:HOH:O	2.11	0.68
1:Z:458:LYS:HE3	1:Z:473:TYR:CE1	2.35	0.61
2:H:87:ARG:NH2	6:H:304:HOH:O	2.25	0.60
2:H:152:LEU:HG	2:H:154:LYS:HG3	1.85	0.58
1:Z:358:ILE:HB	1:Z:395:VAL:CG2	2.34	0.56
3:L:200:GLU:O	6:L:402:HOH:O	2.18	0.55
1:Z:467:ASP:OD1	1:Z:469:SER:HB3	2.06	0.55
2:H:206:ILE:CD1	2:H:221:ARG:HB3	2.37	0.54
2:H:83:MET:HE2	2:H:86:LEU:HD21	1.89	0.54
2:H:221:ARG:HD2	2:H:221:ARG:H	1.74	0.52
3:L:8:ARG:HD3	3:L:147:THR:HG22	1.92	0.51
1:Z:334:ASN:N	6:Z:604:HOH:O	2.43	0.50
2:H:221:ARG:HD2	2:H:221:ARG:N	2.27	0.50
3:L:80:LEU:HG	3:L:107:VAL:HG22	1.94	0.50
1:Z:408:ARG:NH2	6:Z:605:HOH:O	2.44	0.49
4:A:1:NAG:H61	4:A:2:NAG:H82	1.94	0.49
3:L:112:LYS:NZ	6:L:403:HOH:O	2.45	0.49
2:H:47:TRP:CZ2	5:L:301:EDO:H22	2.48	0.48
3:L:104:ARG:HH21	3:L:104:ARG:HB2	1.79	0.48
3:L:33:ASN:O	3:L:52:ALA:HA	2.14	0.47
1:Z:425:LEU:O	6:Z:603:HOH:O	2.21	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1:NAG:H4	4:A:3:FUC:H5	1.98	0.45
1:Z:357:ARG:HG3	1:Z:396:TYR:CE2	2.52	0.45
3:L:151:LYS:HB2	3:L:194:SER:HB2	1.99	0.45
1:Z:337:PRO:O	1:Z:340:GLU:HG2	2.17	0.45
3:L:200:GLU:OE1	6:L:403:HOH:O	2.21	0.45
2:H:206:ILE:HD13	2:H:221:ARG:CB	2.48	0.44
1:Z:383:SER:OG	1:Z:385:THR:HG22	2.18	0.44
2:H:214:SER:O	2:H:214:SER:OG	2.31	0.44
2:H:2:VAL:HG13	2:H:27:PHE:CD1	2.53	0.43
1:Z:444:LYS:HG2	1:Z:448:ASN:HB2	1.99	0.43
1:Z:359:SER:HA	1:Z:524:VAL:CG2	2.48	0.43
2:H:87:ARG:O	2:H:88:ALA:C	2.61	0.43
2:H:24:ALA:HB1	2:H:27:PHE:CE1	2.53	0.43
1:Z:447:GLY:HA2	1:Z:498:GLN:HG2	2.01	0.43
3:L:158:LYS:HB3	3:L:158:LYS:HE3	1.86	0.43
3:L:8:ARG:HD3	3:L:147:THR:CG2	2.49	0.43
1:Z:378:LYS:HA	1:Z:378:LYS:HD2	1.85	0.42
3:L:150:TRP:C	3:L:151:LYS:HD3	2.44	0.42
2:H:50:VAL:HG11	5:L:301:EDO:H11	2.01	0.42
1:Z:383:SER:O	1:Z:387:LEU:HG	2.19	0.42
2:H:206:ILE:HD12	2:H:221:ARG:HB3	2.02	0.41
3:L:105:LEU:C	3:L:105:LEU:HD23	2.45	0.41
2:H:130:PRO:HB3	2:H:156:TYR:HB3	2.01	0.41
1:Z:440:ASN:OD1	1:Z:440:ASN:N	2.53	0.41
1:Z:336:CYS:SG	1:Z:363:ALA:HB2	2.62	0.40
2:H:126:SER:O	2:H:127:THR:C	2.65	0.40
3:L:150:TRP:O	3:L:151:LYS:HD3	2.20	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:H:405:HOH:O	6:H:405:HOH:O[3_555]	1.82	0.38
6:L:527:HOH:O	6:L:527:HOH:O[3_455]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Z	192/205 (94%)	183 (95%)	9 (5%)	0	100	100
2	H	215/227 (95%)	211 (98%)	3 (1%)	1 (0%)	24	16
3	L	209/214 (98%)	200 (96%)	9 (4%)	0	100	100
All	All	616/646 (95%)	594 (96%)	21 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	215	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Z	167/177 (94%)	167 (100%)	0	100	100
2	H	185/192 (96%)	185 (100%)	0	100	100
3	L	174/177 (98%)	174 (100%)	0	100	100
All	All	526/546 (96%)	526 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	Z	334	ASN
2	H	39	GLN
2	H	104	GLN
3	L	39	GLN
3	L	40	GLN
3	L	81	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1	4,1	14,14,15	0.59	0	17,19,21	1.54	3 (17%)
4	NAG	A	2	4	14,14,15	0.40	0	17,19,21	1.19	3 (17%)
4	FUC	A	3	4	10,10,11	0.23	0	14,14,16	1.05	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	A	2	4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FUC	A	3	4	-	-	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	NAG	C1-O5-C5	4.29	117.94	112.19
4	A	3	FUC	C3-C4-C5	-2.74	105.65	109.81
4	A	1	NAG	O5-C1-C2	-2.59	107.28	111.29
4	A	2	NAG	C2-N2-C7	-2.56	119.47	122.90
4	A	2	NAG	O5-C1-C2	-2.47	107.48	111.29
4	A	1	NAG	C1-C2-N2	-2.22	106.94	110.43
4	A	2	NAG	C3-C4-C5	-2.08	106.46	110.23

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1	NAG	C4-C5-C6-O6
4	A	1	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3	FUC	1	0
4	A	2	NAG	1	0
4	A	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	L	301	-	3,3,3	0.57	0	2,2,2	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	L	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	301	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	Z	194/205 (94%)	1.14	32 (16%) 4 4	45, 62, 91, 102	0
2	H	219/227 (96%)	0.67	16 (7%) 21 24	38, 47, 59, 82	0
3	L	211/214 (98%)	0.46	8 (3%) 44 50	35, 44, 67, 92	0
All	All	624/646 (96%)	0.74	56 (8%) 15 17	35, 50, 79, 102	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	1	GLN	4.8
2	H	221	ARG	4.3
3	L	2	ALA	4.3
1	Z	449	TYR	3.9
2	H	215	ASN	3.5
2	H	145	GLY	3.5
1	Z	382	VAL	3.4
1	Z	527	PRO	3.3
1	Z	522	ALA	3.0
2	H	138	SER	2.9
1	Z	395	VAL	2.9
1	Z	478	THR	2.9
2	H	169	ALA	2.9
2	H	181	LEU	2.8
1	Z	371	SER	2.7
2	H	214	SER	2.7
2	H	225	LYS	2.7
2	H	202	THR	2.7
1	Z	368	LEU	2.7
2	H	204	THR	2.6
1	Z	432	CYS	2.6
2	H	187	TYR	2.5
1	Z	519	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	Z	479	PRO	2.5
1	Z	344	ALA	2.5
1	Z	369	TYR	2.5
1	Z	367	VAL	2.5
1	Z	365	TYR	2.4
1	Z	517	LEU	2.4
1	Z	486	PHE	2.4
1	Z	372	ALA	2.3
1	Z	381	GLY	2.3
1	Z	392	PHE	2.3
1	Z	526	GLY	2.3
2	H	212	LYS	2.2
2	H	164	SER	2.2
3	L	159	ALA	2.2
1	Z	525	CYS	2.2
1	Z	470	THR	2.2
1	Z	336	CYS	2.2
3	L	209	ALA	2.2
1	Z	464	PHE	2.1
2	H	157	PHE	2.1
3	L	96	THR	2.1
1	Z	480	CYS	2.1
1	Z	338	PHE	2.1
1	Z	359	SER	2.1
1	Z	390	LEU	2.1
2	H	206	ILE	2.1
2	H	224	PRO	2.1
1	Z	363	ALA	2.1
3	L	188	LYS	2.0
3	L	211	THR	2.0
1	Z	350	VAL	2.0
3	L	30	GLY	2.0
1	Z	341	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	1	14/15	-	-	30,30,30,30	0
4	NAG	A	2	14/15	-	-	30,30,30,30	0
4	FUC	A	3	10/11	-	-	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	L	301	4/4	0.87	0.17	51,53,55,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.